

POLISH ACADEMY OF SCIENCES  
INSTITUTE OF ZOOLOGY

JANUSZ NAST

**PALAEARCTIC AUCHENORRHYNCHA**

**(HOMOPTERA)**

**AN ANNOTATED CHECK LIST**

WARSZAWA 1972  
POLISH SCIENTIFIC PUBLISHERS

Genus: *Asiraca* LATREILLE, 1796.

Type-species: *Cicada clavicornis* FABRICIUS, 1796, by subsequent designation by LATREILLE, 1810.

*Asiraca clavicornis* (FABRICIUS, 1794).

*Cicada clavicornis* FABRICIUS, 1794.

*Asiraca clavicornis divisa* REY, 1891.

Distribution: Afghanistan, Albania, Austria, Belgium, Bulgaria, China (Manchuria, Sinkiang), Cyprus, Czechoslovakia (Bohemia, Moravia, Slovakia), France, German D.R., German F.R., Great Britain (England), Hungary, Ireland, Israel, Italy (also Sardinia and Sicily), Morocco, Netherlands, Poland, Portugal, Romania, Spain, Switzerland, Tunisia, U.S.S.R. (Armenia, Azerbaijan, Georgia, Kazakhstan, Kirghizia, Latvia, Moldavia, s.Russia, Tadzhikistan, Turkmenia, Ukraine, Uzbekistan), Yugoslavia.

Subfamily: *Kelisiinae* WAGNER, 1963.

Genus: *Kelisia* FIEBER, 1866.

Type-species: *Delphax guttula* GERMAR, 1818, by monotypy.

*Kelisia bajancogta* DLABOLA, 1965.

Distribution: Mongolia.

*Kelisia bispinifera* DLABOLA, 1967.

Distribution: Mongolia.

*Kelisia brucki* FIEBER, 1878.

Distribution: Albania, Cyprus, Czechoslovakia (Slovakia), France, Greece, Hungary, Israel, Italy (also Sardinia), U.S.S.R. (s.Russia), Yugoslavia.

*Kelisia confusa* LINNAVUORI, 1957.

Distribution: Italy.

*Kelisia guttula* (GERMAR, 1818).

*Delphax guttula* GERMAR, 1818.

*Kelisia pascuorum* RIBAUT, 1934.

*Kelisia guttula distincta* METCALF, 1943.

*Kelisia guttula fulva* METCALF, 1943.

Distribution: Albania, Algeria, Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Hungary, Ireland, Italy (also Sardinia and Sicily), Norway, Poland, Portugal, Romania, Spain, Sweden, Switzerland, Tunisia, U.S.S.R. (Azerbaijan, Estonia, Latvia, n.Russia, m.Siberia, Tadzhikistan, Ukraine), Yugoslavia.

*Kelisia guttulifera* (KIRSCHBAUM, 1868).

*Delphax guttulifera* KIRSCHBAUM, 1868.

*Kelisia fallax* RIBAUT, 1934.

Distribution: Austria, Czechoslovakia (Moravia), France, German D.R., German F.R., Great Britain (England, Wales), Netherlands, Poland, Yugoslavia.

*Kelisia haupti* WAGNER, 1939.

Distribution: Austria, German D.R., German F.R., Poland.

*Kelisia henschii* HORVÁTH, 1897.

Distribution: Hungary.

*Kelisia irregulata* HAUPT, 1935.

Distribution: Austria, Czechoslovakia (Slovakia), France, German D.R., German F.R.

*Kelisia melanops* FIEBER, 1878.

Distribution: Czechoslovakia (Slovakia), France, Hungary, Italy, Romania, Turkey (Anatolia), Yugoslavia.

*Kelisia melanura* VILBASTE, 1968.

Distribution: U.S.S.R. (Maritime Territory).

*Kelisia minima* RIBAUT, 1934.

Distribution: France.

*Kelisia monoceros* RIBAUT, 1934.

Distribution: Austria, Czechoslovakia (Moravia, Slovakia), Finland, France, German D.R., German F.R., Italy, Poland, Romania, Sweden, U.S.S.R. (Estonia, Latvia, m.Russia).

*Kelisia orchonica* DLABOLA, 1970.

Distribution: Mongolia.

*Kelisia pallidula* (BOHEMAN, 1847).

*Delphax pallidula* BOHEMAN, 1847.

*Delphax raniceps* BOHEMAN, 1847.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), Hungary, Netherlands, Norway, Poland, Romania, Sweden, U.S.S.R. (Estonia, Kazakhstan, Latvia, Ukraine, Uzbekistan), Yugoslavia.

*Kelisia pannonica* MATSUMURA, 1910.

Distribution: German D.R., Hungary.

*Kelisia perrieri* RIBAUT, 1934.

Distribution: Cyprus, France, Italy.

*Kelisia praecox* HAUPT, 1935.

Distribution: Bulgaria, Czechoslovakia (Bohemia, Moravia), German D.R., German F.R., Mongolia, Poland, U.S.S.R. (Estonia, Georgia, Kazakhstan, s.Russia).

***Kelisia punctulum* (KIRSCHBAUM, 1868).**

*Delphax punctulum* KIRSCHBAUM, 1868.

*Kelisia tarda* HAUPT, 1935.

Distribution: Austria, Belgium, Czechoslovakia (Moravia), France, German D.R., German F.R., Great Britain (England), Ireland, Netherlands, U.S.S.R. (n.Russia).

***Kelisia putoni* A. COSTA, 1888.**

Distribution: Italy (also Sardinia), Morocco.

***Kelisia ribauti* WAGNER, 1938.**

Distribution: Afghanistan, Austria, Azores, Canary Is., Cyprus, Czechoslovakia (Bohemia, Moravia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), Iraq, Madeira Archipelago, Morocco, Poland, Portugal, Romania, Sweden, Turkey (Anatolia), U.S.S.R. (Altai Mts., Estonia, Georgia, Kazakhstan, Kirghizia, Latvia, n.Russia, Tadzhikistan, Uzbekistan).

***Kelisia sabulicola* WAGNER, 1952.**

Distribution: German D.R., German F.R., Great Britain (England, Scotland, Wales), Ireland, Netherlands, Poland, Sweden.

***Kelisia sima* RIBAUT, 1934.**

*Kelisia pascuorum sima* RIBAUT, 1934.

Distribution: Austria, France, German F.R.

***Kelisia sulcata* RIBAUT, 1934.**

Distribution: France, Italy (Sardinia).

***Kelisia vittipennis* (J. SAHLBERG, 1868).**

*Delphax vittipennis* J. SAHLBERG, 1868.

Distribution: Algeria, Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Finland, France, German D.R., German F.R., Great Britain (England, Scotland, ? Wales), Hungary, Ireland, Italy, Netherlands, Norway, Poland, Portugal, Romania, Spain, Sweden, Switzerland, U.S.S.R. (Estonia, Latvia, m.Siberia), Yugoslavia.

***Kelisia xiphura* VILBASTE, 1968.**

Distribution: U.S.S.R. (Maritime Territory).

***Kelisia yarkonensis* LINNAVUORI, 1962.**

Distribution: Israel.

Genus: *Anakelisia* WAGNER, 1963.

Type-species: *Ditropis fasciata* KIRSCHBAUM, 1868, by original designation.

***Anakelisia fasciata* (KIRSCHBAUM, 1868).**

*Ditropis fasciata* KIRSCHBAUM, 1868.

*Delphax scotti* SCOTT, 1870.

*Kelisia bicarinata* HAUPT, 1935.  
*Kelisia scotti nigricollis* KUNTZE, 1937.  
*Kelisia scotti quadrimaculata* KUNTZE, 1937.  
*Kelisia fasciata nigra* METCALF, 1943.  
*Kelisia fasciata triangularis* METCALF, 1943.

Distribution: Austria, Czechoslovakia (Bohemia), Denmark, France, German D.R., German F.R., Great Britain (England), Hungary, Poland, Romania, Sweden.

*Anakelisia perspicillata* (BOHEMAN, 1845).

*Delphax perspicillata* BOHEMAN, 1845.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia), Finland, France, German D.R., German F.R., Great Britain (England, Scotland), Hungary, Italy, Mongolia, Poland, Romania, Sweden, Switzerland, U.S.S.R. (Estonia, m.Siberia), Yugoslavia.

Genus: *Stenokelisia* RIBAUT, 1934.

Type-species: *Stenokelisia angusta* RIBAUT, 1934, by original designation.

*Stenokelisia angusta* RIBAUT, 1934.

Distribution: France.

Subfamily: *Saccharosydinae* VILBASTE, 1968.

Genus: *Saccharosydne* KIRKALDY, 1907.

Type-species: *Delphax saccharivora* WESTWOOD, 1833, by original designation.

*Saccharosydne procerus* (MATSUMURA, 1931).

*Oxycranus procerus* MATSUMURA, 1910.

Distribution: China (Manchuria), Japan (Honshu, Kyushu, Shikoku), U.S.S.R. (Maritime Territory); [Oriental region].

Subfamily: *Stenocraninae* WAGNER, 1963.

Genus: *Stenocranus* FIEBER, 1866.

Type-species: *Fulgora minuta* FABRICIUS, 1787, by subsequent designation by OSHANIN, 1912.

*Stenocranus akashiensis* MATSUMURA, 1935.

Distribution: Japan (Honshu).

*Stenocranus aurantiacus* DUBOVSKY, 1966.

Distribution: U.S.S.R. (Kazakhstan, Tadzhikistan, Uzbekistan).

*Stenocranus elongatus* MATSUMURA, 1935.

Distribution: Japan (Kyushu).

- Stenocranus fallax* MATSUMURA, 1935.  
Distribution: China (Shansi), Japan (Hokkaido, Honshu, Kyushu); [Oriental region].
- Stenocranus fuscovittatus* (STÅL, 1858).  
*Delphax fuscovittata* STÅL, 1858.  
*Delphax colorata* KIRSCHBAUM, 1868.  
Distribution: Austria, Belgium, Bulgaria, China (Manchuria), Czechoslovakia (Bohemia, Moravia), Denmark, France, German D.R., German F.R., Great Britain (England), Hungary, Italy, Netherlands, Poland, Portugal, Romania, Switzerland, U.S.S.R. (Estonia, Georgia, Kazakhstan, m.Siberia).
- Stenocranus harimensis* MATSUMURA, 1935.  
Distribution: Japan (Honshu, Kyushu, Shikoku).
- Stenocranus hokkaidoensis* METCALF, 1943.  
*Stenocranus vittatus* MATSUMURA, 1935. [Primary homonym].  
Distribution: Japan (Hokkaido, Honshu, Kyushu), U.S.S.R. (Maritime Territory).
- Stenocranus koreanus* MATSUMURA, 1935.  
Distribution: Korean Peninsula.
- Stenocranus longipennis* (CURTIS, 1837).  
*Delphax longipennis* CURTIS, 1837.  
Distribution: Algeria, France, German F.R., Great Britain (England, Wales), Ireland.
- Stenocranus major* (KIRSCHBAUM, 1868).  
*Delphax major* KIRSCHBAUM, 1868.  
Distribution: Austria, Czechoslovakia (Bohemia, Moravia), German D.R., German F.R., Great Britain (England), Netherlands, Poland, Yugoslavia.
- Stenocranus matsumurai* METCALF, 1943.  
*Stenocranus breviceps* MATSUMURA, 1935. [Primary homonym].  
Distribution: China (Shansi), Japan (Hokkaido, Honshu, Kyushu), U.S.S.R. (Maritime Territory).
- Stenocranus minutus* (FABRICIUS, 1787).  
*Fulgora minuta* FABRICIUS, 1787.  
*Delphax lineola* GERMAR, 1818.  
*Delphax longifrons* BOHEMAN, 1847.  
*Liburnia farinosa* BUCKTON, 1890.  
*Stenocranus lineola illabatus* REY, 1894.  
Distribution: Algeria, Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), Hungary, Ireland, Italy, Netherlands, Norway, Poland, Portugal, Romania, Sweden, Switzerland, Tunisia, U.S.S.R. (Armenia, Azerbaijan, Estonia, Kazakhstan, Kirghizia, Latvia, Moldavia, m. and s.Russia, Ukraine, Uzbekistan), n.Yugoslavia.

- Stenocranus niisimai* MATSUMURA, 1935.  
Distribution: Japan (Hokkaido).
- Stenocranus ozenumensis* ISHIHARA, 1952.  
Distribution: Japan (Honshu).
- Stenocranus pallidus* LINNAVUORI, 1962.  
Distribution: Israel.
- Stenocranus parvulus* VILBASTE, 1968.  
Distribution: U.S.S.R. (Maritime Territory).
- Stenocranus sapporensis* MATSUMURA, 1935.  
Distribution: Japan (Hokkaido, Shikoku).
- Stenocranus silvicola* VILBASTE, 1968.  
Distribution: U.S.S.R. (Maritime Territory).
- Stenocranus sukumonus* MATSUMURA, 1935.  
Distribution: Japan (Shikoku).
- Stenocranus takasagonis* MATSUMURA, 1935.  
Distribution: Japan (Honshu).
- Stenocranus tamagawanus* MATSUMURA, 1935.  
Distribution: Japan (Honshu).
- Stenocranus tateyamanus* MATSUMURA, 1935.  
Distribution: Japan (Honshu, Kyushu).
- Stenocranus yasumatsui* ISHIHARA, 1952.  
Distribution: Japan (Shikoku).

Genus: *Terauchiana* MATSUMURA, 1915.

Type-species: *Terauchiana singularis* MATSUMURA, 1915, by monotypy.

- Terauchiana nigripennis* KATO, 1933.  
Distribution: Japan (Honshu, Kyushu).
- Terauchiana sagitta* (KUSNEZOV, 1929).  
*Tropidocephala sagitta* KUSNEZOV, 1929.  
Distribution: U.S.S.R. (Kazakhstan, Uzbekistan).
- Terauchiana singularis* MATSUMURA, 1915.  
Distribution: Japan (Honshu, Kyushu), Korean Peninsula, U.S.S.R. (Maritime Territory).
- Terauchiana yasumatsui* ESAKI et ISHIHARA, 1950.  
Distribution: China (Shansi).

Subfamily: *Jassidaeinae* WAGNER, 1963.

Genus: *Jassidaeus* FIEBER, 1866.

Type-species: *Jassidaeus morio* FIEBER, 1866, by monotypy.

*Jassidaeus lugubris* (SIGNORET, 1865).

*Delphax lugubris* SIGNORET, 1865.

*Jassidaeus morio* FIEBER, 1866.

*Ditropis atratula* KIRSCHBAUM, 1868.

Distribution: Austria, Czechoslovakia (Bohemia, Moravia, Slovakia), France, German D.R., German F.R., Hungary, Poland, Romania, U.S.S.R. (Moldavia).

Subfamily: *Stirominae* WAGNER, 1963.

Genus: *Delphacinus* FIEBER, 1866.

Type-species: *Delphax mesomela* [sic] BOHEMAN, 1850, by monotypy.

*Delphacinus lukjanovitshi* KUSNEZOV, 1929.

Distribution: U.S.S.R. (m.Siberia).

*Delphacinus mesomelas* (BOHEMAN, 1850).

*Delphax mesomela* [sic] BOHEMAN, 1850.

*Delphax mesomelas*: SCHAUM, 1851.

*Delphacinus alpinus* STROBL, 1900.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland), Hungary, Ireland, Netherlands, Poland, Sweden, Switzerland, U.S.S.R. (Estonia, Kazakhstan, Latvia, Moldavia, Ukraine).

Genus: *Delphacinoides* VILBASTE, 1965.

Type-species: *Delphacinoides altaicus* VILBASTE, 1965, by original designation.

*Delphacinoides altaicus* VILBASTE, 1965.

Distribution: U.S.S.R. (Altai Mts.).

Genus: *Delphacellus* HAUPT, 1929.

Type-species: *Liburnia putoni* SCOTT, 1874, by original designation.

*Delphacellus putoni* (SCOTT, 1874).

*Liburnia putoni* SCOTT, 1874.

Distribution: Algeria, Italy (Sardinia), Tunisia.

Genus: *Stiromella* WAGNER, 1963.

Type-species: *Criomorphus obliquus* WAGNER, 1948, by original designation.

*Stiromella albeola* MITJAEV, 1967.

Distribution: U.S.S.R. (Kazakhstan).

*Stiromella allata* DUBOVSKY, 1970.

Distribution: U.S.S.R. (Kirghizia).

*Stiromella euagropyri* (EMELJANOV, 1964).

*Criomorphus euagropyri* EMELJANOV, 1964.

Distribution: U.S.S.R. (Kazakhstan).

*Stiromella fusca* (LINNAVUORI, 1953).

*Criomorphus fuscus* LINNAVUORI, 1953.

Distribution: U.S.S.R. (Siberia).

*Stiromella inaequalis* (EMELJANOV, 1964).

*Criomorphus inaequalis* [sic] EMELJANOV, 1964.

*Criomorphus inaequalis*: EMELJANOV, 1969.

Distribution: U.S.S.R. (Kazakhstan).

*Stiromella karatavica* MITJAEV, 1968.

Distribution: U.S.S.R. (Kazakhstan).

*Stiromella luctuosa* NAST, 1966.

Distribution: Mongolia.

*Stiromella notata* LOGVINENKO, 1970.

Distribution: U.S.S.R. (Ukraine <Crimea>).

*Stiromella obliqua* (WAGNER, 1948).

*Criomorphus obliquus* WAGNER, 1948.

Distribution: Austria, U.S.S.R. (Estonia, Kazakhstan).

*Stiromella subinaequalis* MITJAEV, 1968.

*Stiromella subinaequalis* [sic] MITJAEV, 1968.

*Stiromella subinaequalis*: MITJAEV, 1971.

Distribution: U.S.S.R. (Kazakhstan).

*Stiromella vitticeps* NAST, 1966.

Distribution: Mongolia.

Genus: *Ditropis* KIRSCHBAUM, 1868.

Type-species: *Delphax pteridis* SPINOLA, 1839, by subsequent designation by OSHANIN, 1912.

*Ditropis pteridis* (SPINOLA, 1839).

*Delphax pteridis* SPINOLA, 1839.

Distribution: Albania, Austria, Belgium, Cyprus, Czechoslovakia (Bohemia,

Slovakia), Denmark, France, German D.R., German F.R., Great Britain (England), Ireland, Italy (also Sardinia), Netherlands, Norway, Poland, Portugal, Spain, Sweden, Switzerland, U.S.S.R. (Georgia), Yugoslavia.

Genus: *Pastiroma* DLABOLA, 1967.

Type-species: *Calligypona odessana* DLABOLA, 1958, by original designation.

*Pastiroma odessana* (DLABOLA, 1958).

*Calligypona odessana* DLABOLA, 1958.

Distribution: Hungary, Mongolia, Romania, U.S.S.R. (Ukraine).

Genus: *Kormus* FIEBER, 1866.

Type-species: *Kormus artemisiae* FIEBER, 1866, by monotypy.

*Kormus artemisiae* FIEBER, 1866.

Distribution: U.S.S.R. (Kazakhstan, s.Russia).

Genus: *Eurysa* FIEBER, 1866.

Type-species: *Delphax lineata* PERRIS, 1857, by subsequent designation by OSHANIN, 1912.

*Epeurysa* MATSUMURA, 1900. Type-species: *Epeurysa nawaii* MATSUMURA, 1900, by monotypy.

*Eurysa brunnea* MELICHAR, 1896.

*Eurysa singeri* KUPKA, 1941.

Distribution: Austria, Czechoslovakia (Moravia), German D.R., German F.R., Hungary, Tunisia.

*Eurysa carinata* MELICHAR, 1899.

Distribution: Tunisia.

*Eurysa clypeata* HORVÁTH, 1897.

Distribution: Austria, Hungary.

*Eurysa douglasi* (SCOTT, 1870).

*Liburnia douglasi* SCOTT, 1870.

Distribution: Great Britain (England), U.S.S.R. (m.Siberia).

*Eurysa immunda* HORVÁTH, 1916.

Distribution: Albania, Italy, Yugoslavia.

*Eurysa laeticiae* DLABOLA, 1967.

Distribution: Italy.

*Eurysa lineata* (PERRIS, 1857).

*Delphax lineata* PERRIS, 1857.

*Eurysa vittata* FIEBER, 1866.

*Delphax quadrivittata* KIRSCHBAUM, 1868.

*Eurysa lineata livens* REY, 1894.

*Eurysa lineata syriaca* DLABOLA, 1965.

Distribution: Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), France, German D.R., German F.R., Great Britain (England), Hungary, Ireland, Italy, Jordan, Mongolia, Netherlands, Poland, Portugal, Romania, Spain, Sweden, Switzerland, Syria, U.S.S.R. (Georgia, Moldavia, Ukraine, Uzbekistan), Yugoslavia.

*Eurysa nawaii* (MATSUMURA, 1900).

*Epeurysa nawaii* MATSUMURA, 1900.

Distribution: Japan (Honshu, Kyushu, Shikoku); [Oriental region].

*Eurysa pyrenaee* FIEBER, 1866.

*Eurysa pyrenaee* [sic] FIEBER, 1866.

*Eurysa pyrenaee dimidiata* REY, 1891.

Distribution: France, Spain.

*Eurysa rabatana* LINDBERG, 1963.

Distribution: Morocco.

*Eurysa ribauti* LINDBERG, 1936.

Distribution: Canary Is., Madeira Archipelago.

*Eurysa sublineata* EMELJANOV, 1964.

Distribution: U.S.S.R. (Kazakhstan).

*Eurysa transbaicalica* KUSNEZOV, 1929.

Distribution: U.S.S.R. (m.Siberia).

Genus: *Stiromeurysa* DLABOLA, 1965.

Type-species: *Stiromeurysa vitoshaensis* DLABOLA, 1965, by original designation.

*Stiromeurysa vitoshaensis* DLABOLA, 1965.

Distribution: Bulgaria.

Genus: *Eurysula* VILBASTE, 1968.

Type-species: *Eurysula lurida* FIEBER, 1866, by original designation.

*Eurysula lurida* (FIEBER, 1866).

*Eurysula lurida* FIEBER, 1866.

*Atropis laevifrons* J. SAHLBERG, 1871.

*Stiroma ruficeps* MELICHAR, 1896.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Finland, France, German D.R., German F.R., Great Britain (England), Hungary, Italy, Netherlands, Norway, Poland, Sweden, U.S.S.R. (Estonia, Kazakhstan, Latvia, n.Russia, Ukraine).

Genus: *Eurybregma* SCOTT, 1875.

Type-species: *Eurybregma nigrolineata* SCOTT, 1875, by monotypy.

*Eurybregma nigrolineata* SCOTT, 1875.

Distribution: Austria, China (Sinkiang), Czechoslovakia (Bohemia, Moravia, Slovakia), France, German D.R., German F.R., Great Britain (England), Hungary, Italy, Mongolia, Poland, U.S.S.R. (Kazakhstan, Moldavia, m.Russia, m.Siberia, Ukraine, Uzbekistan).

*Eurybregma porcus* (EMELJANOV, 1964).

*Criomorphus porcus* EMELJANOV, 1964.

Distribution: Mongolia, U.S.S.R. (Kazakhstan, n.Russia).

*Eurybregma pseudagropyri* EMELJANOV, 1966.

Distribution: U.S.S.R. (m.Siberia).

*Eurybregma sparsa* LOGVINENKO, 1970.

Distribution: U.S.S.R. (Azerbaijan).

Genus: *Niphisa* EMELJANOV, 1966.

Type-species: *Niphisa candens* EMELJANOV, 1966, by original designation.

*Niphisa candens* EMELJANOV, 1966.

Distribution: Mongolia, U.S.S.R. (m.Siberia).

Genus: *Stiroma* FIEBER, 1866.

Type-species: *Stiroma affinis* FIEBER, 1866, by subsequent designation by OSHANIN, 1912.

*Stiroma affinis* FIEBER, 1866.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), n.Italy, Mongolia, Netherlands, Norway, Poland, Romania, Sweden, Switzerland, U.S.S.R. (Altai Mts., Estonia, Kazakhstan, Latvia, n.Russia, Ukraine).

*Stiroma bicarinata* (HERRICH-SCHÄFFER, 1835).

*Delphax bicarinata* HERRICH-SCHÄFFER, 1835.

*Delphax mutabilis* BOHEMAN, 1847.

*Delphax nasalis* BOHEMAN, 1847.

*Stiroma bicarinata nigra* METCALF, 1943.

*Stiroma bicarinata alpha* METCALF, 1943.

*Stiroma bicarinata vittata* METCALF, 1943.

Distribution: Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland), n.Italy, Mongolia, Norway, Poland, Romania,

Sweden, Switzerland, Tunisia, U.S.S.R. (Estonia, Georgia, Kazakhstan, n. and m.Russia, w.Siberia, Ukraine).

*Stiroma montana* HORVÁTH, 1897.

Distribution: Austria, Romania.

*Stiroma nigrifrons* (KUSNEZOV, 1929).

*Metrops* [sic] *nigrifrons* KUSNEZOV, 1929.

Distribution: Mongolia, U.S.S.R. (m.Siberia).

Genus: *Stiromoides* VILBASTE, 1971.

Type-species: *Eurysa maculiceps* HORVÁTH, 1903, by original designation.

*Stiromoides maculiceps* (HORVÁTH, 1903).

*Eurysa maculiceps* HORVÁTH, 1903.

Distribution: Finland, Hungary, Mongolia, U.S.S.R. (Estonia, Kazakhstan, Siberia).

Genus: *Metropis* FIEBER, 1866.

Type-species: *Metropis mayri* FIEBER, 1866, by subsequent designation by OSHANIN, 1912.

*Atropis* KIRSCHBAUM, 1868. Type-species: *Atropis latifrons* KIRSCHBAUM, 1868, by monotypy.

*Metropis achnatheri* EMELJANOV, 1964.

Distribution: China (Inner Mongolia), Mongolia, U.S.S.R. (Kazakhstan, m.Siberia).

*Metropis albipennis* KUSNEZOV, 1929.

Distribution: U.S.S.R. (m.Siberia).

*Metropis festucicola* DLABOLA, 1963.

Distribution: U.S.S.R. (m.Siberia).

*Metropis forficula* HORVÁTH, 1908.

Distribution: France.

*Metropis incisus* LOGVINENKO, 1970.

Distribution: U.S.S.R. (Georgia).

*Metropis inermis* WAGNER, 1939.

Distribution: Austria, Czechoslovakia (Bohemia, Moravia, Slovakia), Italy, Poland, Romania, U.S.S.R. (Kazakhstan, Ukraine <Crimea>).

*Metropis latifrons* (KIRSCHBAUM, 1868).

*Atropis latifrons* KIRSCHBAUM, 1868.

Distribution: Austria, Czechoslovakia (Slovakia), France, German F.R., Hungary, Italy, Poland, Romania.

*Metropis latinus* LINNAVUORI, 1959.

Distribution: Italy.

*Metropis maurus* FIEBER, 1866.

Distribution: Austria, France (Pyrenees), Italy.

*Metropis mayri* FIEBER, 1866.

Distribution: Austria, Belgium, Czechoslovakia (Moravia), France, Hungary, Italy, Switzerland, U.S.S.R. (Altai Mts., Kazakhstan, Moldavia, Ukraine <Crimea>).

*Metropis rubripes* MATSUMURA, 1910.

Distribution: Italy (Sicily).

*Metropis songinitus* DLABOLA, 1967.

Distribution: Mongolia.

*Metropis tridentatus* LOGVINENKO, 1970.

Distribution: U.S.S.R. (Azerbaijan).

*Metropis ugamicus* MITJAEV, 1969.

Distribution: U.S.S.R. (Kazakhstan).

Subfamily: *Tropidocephalinae* MUIR, 1915.

*Tropidocephalini* MUIR, 1915.

Genus: *Tropidocephala* STÅL, 1853.

Type-species: *Tropidocephala flaviceps* STÅL, 1855, by monotypy.

*Nephropsia* A. COSTA, 1862. Type-species: *Fulgura elegans* O. COSTA, 1834, by monotypy.  
*Conicoda* MATSUMURA, 1900. Type-species: *Conicoda graminea* MATSUMURA, 1900, by subsequent designation by KIRKALDY, 1907.

*Smara* DISTANT, 1906. Type-species: *Smara festiva* DISTANT, 1906, by original designation.

*Tropidocephala andropogonis* HORVÁTH, 1895.

Distribution: Czechoslovakia (Slovakia), Yugoslavia.

*Tropidocephala brunnipennis* SIGNORET, 1860.

*Conicoda graminea* MATSUMURA, 1900.

Distribution: Egypt, Japan (Honshu, Kyushu, Shikoku); [Australian, Ethiopian and Oriental regions].

*Tropidocephala erianthi* DLABOLA, 1963.

Distribution: U.S.S.R. (Uzbekistan).

*Tropidocephala festiva* (DISTANT, 1906).

*Smara festiva* DISTANT, 1906.

Distribution: Japan (Honshu, Shikoku); [Oriental region].

***Tropidocephala nigra*** (MATSUMURA, 1900).

*Conicoda nigra* MATSUMURA, 1900.

Distribution: Japan (Honshu, Kyushu, Shikoku).

***Tropidocephala prasina*** MELICHAR, 1902.

*Tropidocephala prasina lateralis* MELICHAR, 1902.

Distribution: Iran.

***Tropidocephala pulchella*** DUBOVSKY, 1966.

Distribution: U.S.S.R. (Tadzhikistan, Uzbekistan).

***Tropidocephala tuberipennis*** (MULSANT et REY, 1855).

*Fulgura* [sic] *elegans* O. COSTA, 1834. [Primary homonym].

*Delphax tuberipennis* MULSANT et REY, 1855.

Distribution: Albania, Algeria, Cyprus, Egypt, s.France, Greece, Israel, Italy (also Sardinia and Sicily), Morocco, Spain, Tunisia, Turkey (Anatolia), Yugoslavia.

Subfamily: *Achorotilinae* WAGNER, 1963.

Genus: *Achorotile* FIEBER, 1866.

Type-species: *Delphax albosignata* DAHLBOM, 1850, by monotypy.

***Achorotile albosignata*** (DAHLBOM, 1850).

*Delphax albosignata* DAHLBOM, 1850.

*Delphax fuscineris* BOHEMAN, 1852.

Distribution: Denmark, Finland, France, German D.R., n.Italy, Norway, n.Poland, Sweden, U.S.S.R. (n. and m.Russia, w.Siberia).

***Achorotile longicornis*** (J. SAHLBERG, 1871).

*Ditropis longicornis* J. SAHLBERG, 1871.

Distribution: Finland, Poland (Tatra Mts.), Sweden, U.S.S.R. (n.Russia).

***Achorotile nobilis*** DLABOLA, 1961.

Distribution: Romania.

***Achorotile transbaicalica*** KUSNEZOV, 1929.

Distribution: U.S.S.R. (m. and e.Siberia).

Genus: *Euconomelus* HAUPT, 1929.

Type-species: *Delphax lepida* BOHEMAN, 1847, by original designation.

***Euconomelus lepidus*** (BOHEMAN, 1847).

*Fulgura limbata* FABRICIUS, 1794. [Primary homonym].

*Delphax lepida* BOHEMAN, 1847.

*Delphax tristis* BOHEMAN, 1847.

Distribution: Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain

(England, Scotland), Hungary, Ireland, Italy, Mongolia, Netherlands, Norway, Poland, Romania, Sweden, Switzerland, U.S.S.R. (Altai Mts., Azerbaijan, Estonia, Georgia, Kazakhstan, Kirghizia, Latvia, n. and m.Russia, w. and m.Siberia, Ukraine, Uzbekistan), Yugoslavia.

Subfamily: *Delphacinae* LEACH, 1815.

*Delphacida* LEACH, 1815.

*Araeopinae* METCALF, 1938.

Genus: *Conomelus* FIEBER, 1866.

Type-species: *Delphax anceps* GERMAR, 1821, by designation under the Plenary Powers of the International Commission on Zoological Nomenclature.

*Conomelus anceps* (GERMAR, 1821).

*Delphax anceps* GERMAR, 1821.

*Delphax pictipennis* CURTIS, 1837.

*Delphax signifera* BOHEMAN, 1845.

*Delphax palliata* BOHEMAN, 1849.

Distribution: Algeria, Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Hungary, Ireland, Italy (Sardinia), Netherlands, Norway, Poland, Romania, Spain, Sweden, Turkey (Anatolia), U.S.S.R. (Estonia, Georgia, Latvia, Moldavia, n. and m.Russia, Ukraine), Yugoslavia.

*Conomelus dehnli* NAST, 1966.

Distribution: Poland.

*Conomelus lorifer* RIBAUT, 1948.

Distribution: German D.R., Italy.

*Conomelus odrysius* DLABOLA, 1965.

Distribution: Bulgaria.

Genus: *Delphax* FABRICIUS, 1798.

Type-species: *Cicada crassicornis* PANZER, 1796, by subsequent designation under the Plenary Powers of the International Commission on Zoological Nomenclature.

*Araeopus* SPINOLA, 1839. Type-species: *Cicada crassicornis* PANZER, 1796, by monotypy.

*Delphax alachanicus* ANUFRIEV, 1970.

Distribution: China (Alashan Desert).

*Delphax armeniacus* ANUFRIEV, 1970.

Distribution: U.S.S.R. (Armenia).

***Delphax crassicornis*** (PANZER, 1796).

*Cicada crassicornis* PANZER, 1796.

*Cicada dubia* PANZER, 1796. [Primary homonym].

*Asiraca grisea* LATREILLE, 1804.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia), Denmark, Finland, German D.R., German F.R., Hungary, Italy (also Sicily), Japan (Hokkaido, Honshu, Kyushu), Netherlands, Poland, Romania, Spain, Sweden, Tunisia, U.S.S.R. (Azerbaijan, Estonia, Kazakhstan, Kirghizia, Latvia, n. and m. Russia, Tadzhikistan, Ukraine), Yugoslavia.

***Delphax inermis*** RIBAUT, 1934.

Distribution: Afghanistan, Algeria, Cyprus, France, Israel, Italy, Morocco, Spain, Spanish Sahara, Tunisia.

***Delphax meridionalis*** (HAUPT, 1924).

*Araeopus meridionalis* HAUPT, 1924.

Distribution: Greece, Italy, Yugoslavia.

***Delphax narbonensis*** RIBAUT, 1934.

Distribution: France.

***Delphax orientalis*** (LINNAVUORI, 1955).

*Araeopus orientalis* LINNAVUORI, 1955.

Distribution: U.S.S.R. (Kazakhstan, Kirghizia, Turkmenia).

***Delphax pulchellus*** (CURTIS, 1833).

*Asiraca pulchella* CURTIS, 1833.

*Asiraca vittata* SNELLEN VAN VOLLENHOVEN, 1862.

*Areopus* [sic] *minki* FIEBER, 1866.

Distribution: Austria, Belgium, Bulgaria, Czechoslovakia (Moravia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Wales), Ireland, Italy (also Sardinia), Netherlands, Norway, Poland, Sweden, U.S.S.R. (Estonia), Yugoslavia.

Genus: ***Euides*** FIEBER, 1866.

Type-species: *Delphax basilinea* GERMAR, 1821, by subsequent designation by MUIR, 1915.

*Euidella* PUTON, 1886. Type-species: *Delphax basilinea* GERMAR, by subsequent designation by OSHANIN, 1912.

***Euides alpina*** WAGNER, 1948.

Distribution: Austria, Poland, U.S.S.R. (Kazakhstan, Kirghizia, s. Russia <Dagestan>).

***Euides basilinea*** (GERMAR, 1821).

*Delphax basilinea* GERMAR, 1821.

Distribution: U.S.S.R. (Ukraine <Crimea>).

*Euides caspiana* (DLABOLA, 1961).

*Euidella caspiana* DLABOLA, 1961.

Distribution: U.S.S.R. (s.Russia <Dagestan>).

*Euides speciosa* (BOHEMAN, 1845).

*Delphax speciosa* BOHEMAN, 1845.

*Delphax asiracoides* FOREL, 1858.

*Euidella speciosa obscura* METCALF, 1943.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), Hungary, Italy, Netherlands, Poland, Romania, Sweden, U.S.S.R. (Azerbaijan, Estonia, Kazakhstan, n. and s.Russia, Ukraine), Yugoslavia.

*Euides velitchkovskyi* (MELICHAR, 1913).

*Euidella velitchkovkyi* MELICHAR, 1913.

Distribution: U.S.S.R. (Kazakhstan, s.Russia).

Genus: *Epunka* MATSUMURA, 1935.

Type-species: *Epunka bilineata* MATSUMURA, 1935, by original designation.

*Epunka bilineata* MATSUMURA, 1935.

Distribution: Japan (Hokkaido).

Subfamily: *Chlorioninae* WAGNER, 1963.

Genus: *Chloriona* FIEBER, 1866.

Type-species: *Delphax unicolor* HERRICH-SCHÄFFER, 1835, by subsequent designation by KIRKALDY, 1907.

*Chloriona alaica* DUBOVSKY, 1970.

Distribution: U.S.S.R. (Kirghizia).

*Chloriona arakawai* MATSUMURA, 1935.

Distribution: Japan (Shikoku).

*Chloriona canariensis* LINDBERG, 1954.

Distribution: Canary Is., Turkey (Anatolia), U.S.S.R. (Kazakhstan, Moldavia, s.Russia <Dagestan>), Uzbekistan).

*Chloriona chinai* OSSIANNILSSON, 1946.

Distribution: Finland, Sweden, U.S.S.R. (Estonia, n.Russia, Siberia).

*Chloriona clavata* DLABOLA, 1960.

Distribution: U.S.S.R. (Kazakhstan, Moldavia, s.Russia <Dagestan>), Uzbekistan).

*Chloriona dorsata* EDWARDS, 1898.

*Chloriona danica* JENSEN-HAARUP, 1917.

Distribution: Denmark, France, Great Britain (England), Poland.

*Chloriona flaveola* LINDBERG, 1948.

Distribution: Cyprus, Egypt, Israel, Morocco, Turkey (Anatolia), U.S.S.R. (Kazakhstan).

*Chloriona glaucescens* FIEBER, 1866.

*Chloriona unicolor alba* METCALF, 1943.

*Chloriona unicolor dorsalis* METCALF, 1943.

*Chloriona unicolor flava* METCALF, 1943.

*Chloriona unicolor nigrofasciata* METCALF, 1943.

*Chloriona unicolor nigromaculata* METCALF, 1943.

Distribution: Czechoslovakia (Bohemia, Moravia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland), Hungary, n.Italy, Netherlands, Norway, Poland, Romania, Sweden, U.S.S.R. (Estonia, Georgia, Kazakhstan, Uzbekistan).

*Chloriona indigena* DUBOVSKY, 1966.

Distribution: U.S.S.R. (Kirghizia).

*Chloriona japonica* MATSUMURA, 1935.

Distribution: Japan (Hokkaido, Honshu, Kyushu).

*Chloriona litoralis* MITJAEV, 1971.

Distribution: Kazakhstan.

*Chloriona oranensis* MATSUMURA, 1910.

Distribution: Algeria.

*Chloriona oriens* (DLABOLA, 1961).

*Calligypona oriens* DLABOLA, 1961.

Distribution: Bulgaria, Mongolia, Romania, U.S.S.R. (s.Russia <Dagestan>).

*Chloriona sachalinensis* MATSUMURA, 1935.

Distribution: U.S.S.R. (Sakhalin).

*Chloriona shikokuana* MATSUMURA, 1935.

*Chloriona sukumonis* MATSUMURA, 1935.

Distribution: Japan (Honshu, Shikoku).

*Chloriona sicula* MATSUMURA, 1910.

Distribution: Italy (Sicily).

*Chloriona smaragdula* (STÅL, 1853).

*Delphax smaragdula* STÅL, 1835.

*Chloriona prasinula* FIEBER, 1872.

Distribution: Austria, Belgium, Czechoslovakia (Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), Hungary, n.Italy, Netherlands, Poland, Sweden, Switzerland, [?] Syria, [?] Tunisia, U.S.S.R. (Estonia, Kazakhstan, n. and m.Russia, w.Siberia), Yugoslavia.

*Chloriona stenoptera* (FLOR, 1861).

*Delphax stenoptera* FLOR, 1861.

Distribution: Poland, U.S.S.R. (Latvia, Kazakhstan).

*Chloriona superba* EMELJANOV, 1964.

Distribution: U.S.S.R. (Kazakhstan).

*Chloriona tateyamana* MATSUMURA, 1935.

Distribution: Japan (Honshu), U.S.S.R. (Kazakhstan, Maritime Territory).

*Chloriona tosaensis* MATSUMURA, 1935.

Distribution: Japan (Shikoku).

*Chloriona unicolor* (HERRICH-SCHÄFFER, 1835).

*Delphax unicolor* HERRICH-SCHÄFFER, 1835.

*Chloriona edwardsi* LE QUESNE, 1960.

Distribution: Austria, France, German D.R., German F.R., Great Britain (England), Hungary, [?] Iran, Ireland, Italy (also Sardinia), Tunisia, U.S.S.R. (Georgia, Kazakhstan, Kirghizia, m. and s.Russia, Ukraine, Uzbekistan), Yugoslavia.

*Chloriona vasconica* RIBAUT, 1934.

Distribution: Czechoslovakia (Bohemia, Slovakia), France, German D.R., German F.R., Great Britain (England), Hungary, Italy, Poland, Sweden, U.S.S.R. (s.Russia).

*Chloriona zaisanica* MITJAEV, 1971.

Distribution: U.S.S.R. (Kazakhstan).

Subfamily: *Criomorphinae* KIRKALDY, 1910.

*Criomorphini* KIRKALDY, 1910.

*Megamelinae* HAUPT, 1929.

Genus: *Megamelus* FIEBER, 1866.

Type-species: *Delphax notula* GERMAR, 1830, by monotypy.

*Megamelus discrepans* HAUPT, 1930.

Distribution: Italy.

*Megamelus leptus* FIEBER, 1878.

Distribution: Czechoslovakia (Bohemia), n.Italy.

*Megamelus notula* (GERMAR, 1830).

*Delphax notula* GERMAR, 1830.

*Delphax truncatipennis* BOHEMAN, 1847.

*Megamelus brevifrons* REUTER, 1880.

*Megamelus notula fuscus* METCALF, 1943.

*Megamelus notula pallidus* METCALF, 1943.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia),

Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Hungary, Ireland, Italy, Japan (Honshu), Mongolia, Netherlands, Norway, Poland, Romania, Sweden, U.S.S.R. (Altai Mts., Azerbaijan, Estonia, Kazakhstan, Latvia, n. and m. Russia, w. Siberia, Ukraine), Yugoslavia.

*Megamelus olorinus* DLABOLA, 1961.

Distribution: Romania, U.S.S.R. (s. Russia <Dagestan>, Uzbekistan).

Genus: *Kakuna* MATSUMURA, 1935.

Type-species: *Kakuna kuwayamai* MATSUMURA, 1935, by original designation.

*Kakuna kuwayamai* MATSUMURA, 1935.

Distribution: Japan (Hokkaido, Honshu, Kyushu, Shikoku).

*Kakuna sapporonis* (MATSUMURA, 1935).

*Toya sapporonis* MATSUMURA, 1935.

Distribution: Japan (Hokkaido, Honshu, Kyushu), U.S.S.R. (Maritime Territory).

Genus: *Coracodelphax* VILBASTE, 1968.

Type-species: *Coracodelphax obscurus* VILBASTE, 1968, by original designation.

*Coracodelphax obscurus* VILBASTE, 1968.

Distribution: U.S.S.R. (Maritime Territory).

Genus: *Unkanodes* FENNAH, 1956.

Type-species: *Unkana sapporona* MATSUMURA, 1935, by original designation.

*Elymodelphax* WAGNER, 1963. Type-species: *Liburnia excisa* MELICHAR, 1898, by original designation.

*Chilodelphax* VILBASTE, 1968. [Subgenus of *Unkanodes*]. Type-species: *Unkanodes (Chilodelphax) silvaticus* VILBASTE, 1968, by original designation.

*Unkanodes albifascia* (MATSUMURA, 1900).

*Liburnia albifascia* MATSUMURA, 1900.

Distribution: Japan (Honshu), U.S.S.R. (Maritime Territory).

*Unkanodes excisa* (MELICHAR, 1898).

*Liburnia excisa* MELICHAR, 1898.

*Liburnia elymi* JENSEN-HAARUP, 1915.

Distribution: Denmark, Finland, German D.R., German F.R., Poland, Sweden, U.S.S.R. (Estonia, Kurile Is., n. Russia, Ukraine).

*Unkanodes latespinosa* (DLABOLA, 1957).

*Calligypona latespinosa* DLABOLA, 1957.

Distribution: Afghanistan, Mongolia, Turkey (Anatolia).

***Unkanodes paramarginata* (DLABOLA, 1961).**

*Calligypona paramarginata* DLABOLA, 1961.

Distribution: U.S.S.R. (Tadzhikistan, Uzbekistan).

***Unkanodes sapporona* (MATSUMURA, 1935).**

*Unkana sapporona* MATSUMURA, 1935.

Distribution: China (Shansi), Japan (Hokkaido, Kyushu), U.S.S.R. (Maritime Territory); [Oriental region].

***Unkanodes sibirica* (KUSNEZOV, 1929).**

*Liburnia sibirica* KUSNEZOV, 1929.

*Megadelphax ustkanica* VILBASTE, 1965.

Distribution: Mongolia, U.S.S.R. (Altai Mts., m.Siberia).

***Unkanodes silvatica* VILBASTE, 1968.**

*Unkanodes (Chilodelphax) silvaticus* VILBASTE, 1968.

Distribution: U.S.S.R. (Maritime Territory).

***Unkanodes tanasijevici* (DLABOLA, 1965).**

*Elymodelphax tanasijevici* DLABOLA, 1965.

Distribution: Yugoslavia.

Genus: ***Unkanodella* VILBASTE, 1968.**

Type-species: *Unkanodella ussuriensis* VILBASTE, 1968, by original designation.

***Unkanodella ussuriensis* VILBASTE, 1968.**

Distribution: U.S.S.R. (Maritime Territory).

Genus: ***Megadelphax* WAGNER, 1963.**

Type-species: *Delphax sordidula* STÅL, 1853, by original designation.

***Megadelphax kangauzi* ANUFRIEV, 1970.**

Distribution: U.S.S.R. (Maritime Territory).

***Megadelphax sobrinus* DLABOLA, 1967.**

*Megadelphax sobrina* DLABOLA, 1967.

Distribution: Mongolia.

***Megadelphax sordidulus* (STÅL, 1853).**

*Delphax sordidula* STÅL, 1853.

*Delphacodes sordidula sahlbergi* METCALF, 1943.

Distribution: Algeria, Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Finland, France, German D.R., German F.R., Hungary, Italy, Mongolia, Netherlands, Poland, Romania, Sweden, Switzerland, Tunisia, U.S.S.R. (Estonia, Kazakhstan, Latvia, n. and s.Russia, m.Siberia), Yugoslavia.

Genus: *Laodelphax* FENNAH, 1963.

Type-species: *Delphax striatella* FALLÉN, 1826, by original designation.

*Callidelphax* WAGNER, 1963. Type-species: *Delphax striatella* FALLÉN, 1826, by original designation.

*Laodelphax striatellus* (FALLÉN, 1826).

*Delphax striatella* FALLÉN, 1826.

*Liburnia striatella lateralis* FIEBER, 1879.

*Delphax striatella fimbriata* REY, 1894.

*Delphax striatella dorsalis* REY, 1894. [Primary homonym].

*Liburnia devastans* MATSUMURA, 1900.

*Liburnia nipponica* MATSUMURA, 1900.

*Liburnia minoensis* MATSUMURA, 1900.

*Liburnia giffuensis* MATSUMURA, 1900.

*Liburnia akashiensis* MATSUMURA, 1900.

*Liburnia maikoensis* MATSUMURA, 1900.

*Liburnia niveopicta* HAUPT, 1927.

*Delphacodes striatella reyana* METCALF, 1943.

Distribution: Afghanistan, Albania, Algeria, Austria, Bulgaria, Canary Is., China (Manchuria, Shansi), Czechoslovakia (Bohemia, Moravia, Slovakia), Finland, France, German D.R., German F.R., Great Britain (England), Hungary, Iran, Iraq, Israel, Italy (also Sardinia and Sicily), Japan (Hokkaido, Honshu, Kyushu, Shikoku), Korean Peninsula, Lebanon, Madeira Archipelago, Mongolia, Netherlands, Poland, Portugal, Romania, Spain, Sweden, Tunisia, Turkey (Anatolia), U.S.S.R. (Altai Mts., Armenia, Azerbaijan, Estonia, Georgia, Kazakhstan, Kirghizia, Kurile Is., Latvia, Maritime Territory, Moldavia, n., m. and s.Russia, w.Siberia, Tadzhikistan, Ukraine, Uzbekistan), Yugoslavia.; [Oriental region].

Genus: *Leptodelphax* HAUPT, 1927.

Type-species: *Leptodelphax cyclops* HAUPT, 1927, by original designation.

*Leptodelphax cyclops* HAUPT, 1927.

Distribution: Cyprus, Egypt, Israel; [Ethiopian region].

Genus: *Himeunka* MATSUMURA et ISHIHARA, 1945.

Type-species: *Unkana tateyamaella* MATSUMURA, 1935, by original designation.

*Himeunka tateyamaella* (MATSUMURA, 1935).

*Unkana tateyamaella* MATSUMURA, 1935.

*Unkana kushiana* MATSUMURA, 1935.

Distribution: Japan (Honshu, Kyushu).

Genus: *Hosunka* MATSUMURA, 1935.

Type-species: *Hosunka pallidula* MATSUMURA, 1935, by original designation.

*Unkana* MATSUMURA, 1935. [Preoccupied]. Type-species: *Unkana hakonensis* MATSUMURA, 1935, by original designation.

*Unkanella* ESAKI et ISHIHARA, 1943. Type-species: *Unkana hakonensis* MATSUMURA, 1935, by original designation.

***Hosunka hakonensis* (MATSUMURA, 1935).**

*Unkana hakonensis* MATSUMURA, 1935.

Distribution: Japan (Hokkaido, Honshu, Kyushu).

***Hosunka testacea* (MATSUMURA, 1900).**

*Chlorionidea testacea* MATSUMURA, 1900.

Distribution: Japan (Honshu).

Genus: *Matsumuranoda* METCALF, 1943.

Type-species: *Noda karafutona* MATSUMURA, 1935, by original designation.

*Noda* MATSUMURA, 1935. [Preoccupied]. Type-species: *Noda karafutona* MATSUMURA, 1935, by original designation.

***Matsumuranoda karafutona* (MATSUMURA, 1935).**

*Noda karafutona* MATSUMURA, 1935.

Distribution: U.S.S.R. (Sakhalin).

Genus: *Sogatella* FENNAH, 1956.

Type-species: *Delphax furcifera* HORVÁTH, 1899, by original designation.

*Sogatella* FENNAH, 1956. [Subgenus of *Chloriona*].

***Sogatella catoptron* FENNAH, 1963.**

Distribution: Egypt, Israel, Jordan.

***Sogatella elegantissima* (ISHIHARA, 1952).**

*Delphacodes elegantissima* ISHIHARA, 1952.

Distribution: Japan (Shikoku).

***Sogatella furcifera* (HORVÁTH, 1899).**

*Delphax furcifera* HORVÁTH, 1899.

Distribution: Canary Is., China (Manchuria, Shansi), Egypt, Italy (Sicily), Japan (Hokkaido, Honshu, Kyushu, Shikoku), Korean Peninsula, Madeira Archipelago, Mongolia, U.S.S.R. (Kurile Is., Maritime Territory); [Oriental region].

***Sogatella kyushuensis* (MATSUMURA et ISHIHARA, 1945).**

*Sogata kyushuensis* MATSUMURA et ISHIHARA, 1945.

Distribution: Japan (Kyushu).

***Sogatella longifurcifera* (ESAKI et ISHIHARA, 1947).**

*Delphacodes longifurcifera* ESAKI et ISHIHARA, 1947.

Distribution: Japan (Kyushu), Mongolia, U.S.S.R. (Maritime Territory); [Oriental and Australian regions].

*Sogatella nigeriensis* (MUIR, 1920).

*Megamelus furcifer nigeriensis* MUIR, 1920.

Distribution: Egypt; [Ethiopian region].

*Sogatella panicicola* (ISHIHARA, 1949).

*Delphacodes panicicola* ISHIHARA, 1949.

Distribution: Japan (Honshu, Shikoku), U.S.S.R. (Maritime Territory).

*Sogatella petax* FENNAH, 1963.

Distribution: Egypt, Jordan.

*Sogatella sirokata* (MATSUMURA et ISHIHARA, 1945).

*Sogata sirokata* MATSUMURA et ISHIHARA, 1945.

Distribution: Japan (Honshu, Kyushu); [Oriental region].

*Sogatella suezensis* (MATSUMURA, 1910).

*Delphax suezensis* MATSUMURA, 1910.

*Liburnia vibix* HAUPT, 1927.

Distribution: Azores, Canary Is., Cyprus, Egypt, Iran, Iraq, Israel, Jordan, Lebanon, Madeira Archipelago, Morocco, U.S.S.R. (Ukraine <Crimea>), Yugoslavia; [Ethiopian region].

Genus: *Ditropsis* WAGNER, 1963.

Type-species: *Delphax flavipes* SIGNORET, 1865, by original designation.

*Ditropsis flavipes* (SIGNORET, 1865).

*Delphax flavipes* SIGNORET, 1865.

*Dicranotropis nitida* HAUPT, 1924.

Distribution: Austria, Belgium, Czechoslovakia (Moravia, Slovakia), Denmark, France, German D.R., German F.R., Hungary, Italy, Mongolia, Poland, Spain, Switzerland, Turkey (Anatolia), U.S.S.R. (Kazakhstan, m.Russia, m.Siberia, Ukraine).

Genus: *Nilaparvata* DISTANT, 1906.

Type-species: *Nilaparvata greeni* DISTANT, 1906, by original designation.

*Nilaparvata bakeri* (MUIR, 1917).

*Delphacodes bakeri* MUIR, 1917.

Distribution: Japan (Honshu, Kyushu); [Oriental region].

*Nilaparvata lugens* (STÅL, 1854).

*Delphax lugens* STÅL, 1854.

*Delphax oryzae* MATSUMURA, 1907.

Distribution: China (Manchuria), Japan (Hokkaido, Honshu, Kyushu, Shikoku), Korean Peninsula; [Oriental and Australian regions].

Genus: *Paraliburnia* JENSEN-HAARUP, 1917.

Type-species: *Paraliburnia jacobsoni* JENSEN-HAARUP, 1917, by original designation.

*Paraliburnia adela* (FLOR, 1861).

*Delphax adela* FLOR, 1861.

*Delphax concolor* FIEBER, 1866.

*Delphax sordida* KIRSCHBAUM, 1868.

*Liburnia signoreti* SCOTT, 1870.

*Paraliburnia jacobsoni* JENSEN-HAARUP, 1917.

Distribution: Austria, Czechoslovakia (Bohemia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), Ireland, Netherlands, Poland, Sweden, U.S.S.R. (Estonia, Kazakhstan, Latvia, s.Russia, Siberia).

*Paraliburnia clypealis* (J. SAHLBERG, 1871).

*Liburnia clypealis* J. SAHLBERG, 1871.

Distribution: Czechoslovakia (Bohemia), Finland, German D.R., Great Britain (England), Sweden, U.S.S.R. (n.Russia).

Genus: *Hyledelphax* VILBASTE, 1968.

Type-species: *Delphax elegantula* BOHEMAN, 1847, by original designation.

*Hyledelphax elegantulus* (BOHEMAN, 1847).

*Delphax elegantula* BOHEMAN, 1847.

*Liburnia aemulator* SCOTT, 1873.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland), Hungary, Ireland, Italy, Mongolia, Netherlands, Norway, Poland, Romania, Sweden, Tunisia, U.S.S.R. (Estonia, Kazakhstan, Latvia, Moldavia, n. and m.Russia, Ukraine), Yugoslavia.

Genus: *Megamelodes* LE QUESNE, 1960.

Type-species: *Delphax quadrimaculatus* SIGNORET, 1865, by original designation.

*Megamelodes lequesnei* WAGNER, 1963.

Distribution: [?] Belgium, German D.R., Great Britain (England, Scotland), Ireland, [?] Netherlands, [?] Portugal, [?] Romania, U.S.S.R. ([?] s.Russia <Dagestan>).

*Megamelodes quadrimaculatus* (SIGNORET, 1865).

*Delphax quadrimaculatus* SIGNORET, 1865.

*Liburnia fieberi* SCOTT, 1870.

Distribution: Austria, Belgium, France, German D.R., German F.R., Great Britain (England, Wales), Hungary, Ireland, Italy, Madeira Archipelago, Netherlands.

Genus: *Calligypona* J. SAHLBERG, 1871.

Type-species: *Calligypona albicollis* J. SAHLBERG, 1871, by monotypy.

***Calligypona reyi* (FIEBER, 1866).**

*Delphax reyi* FIEBER, 1866.

*Delphax longipes* KIRSCHBAUM, 1868.

*Calligypona albicollis* J. SAHLBERG, 1871.

*Delphacodes sahlbergiana* ISHIHARA, 1949.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), Hungary, Italy, Mongolia, Netherlands, Poland, Sweden, U.S.S.R. (Estonia, Maritime Territory, Tadzhikistan, Uzbekistan), Yugoslavia.

Genus: *Delphacodes* FIEBER, 1866.

Type-species: *Delphax (Delphacodes) mulsanti* FIEBER, 1866, by subsequent designation by KIRKALDY, 1904.

***Delphacodes albifrons* (FIEBER, 1879).**

*Liburnia albifrons* FIEBER, 1879.

Distribution: Austria, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), France, German D.R., German F.R., Netherlands, Poland, Tunisia, U.S.S.R. (Kazakhstan), Yugoslavia.

***Delphacodes audrasi* RIBAUT, 1954.**

*Delphacodes linnavuorii* LE QUESNE, 1960.

Distribution: Albania, Bulgaria, Czechoslovakia (Slovakia), France, Israel, Romania, Turkey (Anatolia), Yugoslavia.

***Delphacodes capnodes* (SCOTT, 1870).**

*Liburnia capnodes* SCOTT, 1870.

*Megamelus brevifrons pilosa* HAUPT, 1935.

*Megamelus paludicola* LINDBERG, 1937.

Distribution: Czechoslovakia (Bohemia), German D.R., German F.R., Great Britain (England), Hungary, Netherlands, Poland, Sweden, U.S.S.R. (Estonia), Yugoslavia.

***Delphacodes mulsanti* (FIEBER, 1866).**

*Delphax (Delphacodes) mulsanti* FIEBER, 1866.

Distribution: France.

***Delphacodes venosus* (GERMAR, 1830).**

*Delphax venosa* GERMAR, 1830.

*Delphax rhypara* FLOR, 1861.

*Delphax piceola* KIRSCHBAUM, 1868.

*Liburnia melanopachys* SCOTT, 1870.

*Liburnia curtula* J. SAHLBERG, 1871.

*Liburnia thoulessi* EDWARDS, 1894.

Distribution: Albania, Austria, Belgium, Czechoslovakia (Bohemia), Den-

mark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland), Hungary, Italy, Netherlands, Norway, Poland, Sweden, Switzerland, U.S.S.R. (Estonia, Latvia, Moldavia, n.Russia, Ukraine), Yugoslavia.

Genus: *Gravesteiniella* WAGNER, 1963.

Type-species: *Liburnia boldi* SCOTT, 1870, by original designation.

*Gravesteiniella boldi* (SCOTT, 1870).

*Liburnia boldi* SCOTT, 1870.

*Liburnia consanguinea* SCOTT, 1873.

*Liburnia tschikoica* KUSNEZOV, 1929.

*Calligypona bicolor* LINDBERG, 1948.

*Gravesteiniella boldi rubricarina* MITJAEV, 1967.

Distribution: Cyprus, Czechoslovakia (Slovakia), Denmark, German D.R., German F.R., Great Britain (England, Scotland), Hungary, Ireland, Mongolia, Netherlands, Poland, Sweden, U.S.S.R. (Altai Mts., Estonia, Kazakhstan, Latvia, Maritime Territory, s.Russia, m.Siberia, Ukraine).

Genus: *Elachodelphax* VILBASTE, 1965.

Type-species: *Liburnia metcalfi* KUSNEZOV, 1929, by original designation.

*Elachodelphax metcalfi* (KUSNEZOV, 1929).

*Liburnia metcalfi* KUSNEZOV, 1929.

Distribution: U.S.S.R. (Altai Mts., Kazakhstan, m.Siberia).

Genus: *Muellerianella* WAGNER, 1963.

Type-species: *Delphax fairmairei* PERRIS, 1857, by original designation.

*Muellerianella brevipennis* (BOHEMAN, 1847).

*Fulgora flavescens* FABRICIUS, 1794. [Primary homonym].

*Delphax brevipennis* BOHEMAN, 1847.

*Delphax bivittata* BOHEMAN, 1850.

*Delphax hyalinipennis* STÅL, 1854.

*Delphax neglecta* FLOR, 1861, partim.

*Delphacodes brevipennis sordidella* METCALF, 1943.

Distribution: Austria, Azores, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Hungary, Italy (also Sardinia), Netherlands, Norway, Poland, Romania, Sweden, Switzerland, U.S.S.R. (Estonia, Latvia, n.Russia), Yugoslavia.

*Muellerianella fairmairei* (PERRIS, 1857).

*Delphax fairmairii* [sic] PERRIS, 1857.

*Delphax neglecta* FLOR, 1861, partim.

*Liburnia extrusa* SCOTT, 1871.

*Delphax fairmairei signicollis* REY, 1894.

Distribution: Austria, Azores, Belgium, China (Kansu), Czechoslovakia (Bohemia, Moravia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Ireland, Italy, Madeira Archipelago, Netherlands, Poland, Romania, Sweden, Switzerland, U.S.S.R. (Estonia, Latvia, Maritime Territory, n. and m. Russia, Ukraine).

Genus: *Trichodelphax* VILBASTE, 1968.

Type-species: *Trichodelphax splendidulus* VILBASTE, 1968, by original designation.

*Trichodelphax splendidulus* VILBASTE, 1968.

Distribution: U.S.S.R. (Maritime Territory).

Genus: *Chlorionidea* Löw, 1885.

Type-species: *Chlorionidea flava* Löw, 1885, by monotypy.

*Chlorionidea bromi* EMELJANOV, 1964.

Distribution: U.S.S.R. (Kazakhstan, Kirghizia).

*Chlorionidea flava* Löw, 1885.

*Stenocranus styriacus* STROBL, 1900.

Distribution: Austria, Czechoslovakia (Moravia, Slovakia), German D.R., Italy, s. Poland, n. Yugoslavia.

Genus: *Muirodelphax* WAGNER, 1963.

Type-species: *Delphax aubei* PERRIS, 1857, by original designation.

*Muirodelphax atratulus* VILBASTE, 1968.

Distribution: U.S.S.R. (Maritime Territory).

*Muirodelphax aubei* (PERRIS, 1857).

*Delphax aubei* PERRIS, 1857.

*Delphax cognata* FIEBER, 1866.

*Delphax protrusa* FIEBER, 1866.

*Delphax obsoleta* KIRSCHBAUM, 1868.

*Delphax obscurinervis* KIRSCHBAUM, 1868.

Distribution: Afghanistan, Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Ireland, Italy, Hungary, Mongolia, Netherlands, Poland, Portugal, Sweden, Tunisia, Turkey (Anatolia), U.S.S.R. (Estonia, Georgia, Kazakhstan, Kirghizia, Moldavia, Tadzhikistan, Ukraine, Uzbekistan), Yugoslavia.

*Muirodelphax litoralis* VILBASTE, 1968.

Distribution: U.S.S.R. (Maritime Territory).

Genus: *Acanthodelphax* LE QUESNE, 1964.

Type-species: *Delphax denticauda* BOHEMAN, 1847, by original designation.  
*Acanthodelphax* LE QUESNE, 1964. [Subgenus of *Muirodelphax*].

*Acanthodelphax denticauda* (BOHEMAN, 1847).

*Delphax denticauda* BOHEMAN, 1847.

*Liburnia insignis* SCOTT, 1882.

*Delphax conwentzi* MATSUMURA, 1906.

*Liburnia oxyura* HAUPT, 1935.

*Delphacodes denticauda palliditestacea* METCALF, 1943.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland), Norway, Poland, Switzerland, Sweden, U.S.S.R. (Estonia, Latvia, n.Russia).

*Acanthodelphax spinosus* (FIEBER, 1866).

*Delphax spinosa* FIEBER, 1866.

*Delphax ferruginea* KIRSCHBAUM, 1868.

Distribution: Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, France, German D.R., German F.R., Hungary, Italy, Netherlands, Poland, Romania, U.S.S.R. (Ukraine).

Genus: *Dichromina* MITJAEV, 1971.

Type-species: *Calligypona venustula* EMELJANOV, 1964, by monotypy.

*Dichromina venustula* (EMELJANOV, 1964).

*Calligypona venustula* EMELJANOV, 1964.

Distribution: U.S.S.R. (Kazakhstan).

Genus: *Halmyra* MITJAEV, 1971.

Type-species: *Calligypona aeluropodis* EMELJANOV, 1964, by monotypy.

*Halmyra aeluropodis* (EMELJANOV, 1964).

*Calligypona aeluropodis* EMELJANOV, 1964.

Distribution: U.S.S.R. (Kazakhstan).

Genus: *Tyrphodelphax* VILBASTE, 1968.

Type-species: *Delphax distincta* FLOR, 1861, by original designation.

*Tyrphodelphax albocarinatus* (STÅL, 1858).

*Delphax albocarinata* STÅL, 1858.

*Liburnia albocarinata intermedia* J. SAHLBERG, 1871.

*Delphacodes albocarinata palliata* METCALF, 1943.

Distribution: Austria, Czechoslovakia (Bohemia), Denmark, Finland, German D.R., German F.R., Norway, Poland, Sweden, U.S.S.R. (Estonia, n.Russia).

*Tyrphodelphax distinctus* (FLOR, 1861).

*Delphax distincta* FLOR, 1861.

Distribution: Austria, Czechoslovakia (Bohemia), Denmark, Finland, German D.R., German F.R., Great Britain (England, Scotland), Ireland, Poland, Sweden, Switzerland, U.S.S.R. (Estonia, Latvia).

*Tyrphodelphax haglundi* (J. SAHLBERG, 1871).

*Liburnia haglundi* J. SAHLBERG, 1871.

Distribution: Czechoslovakia (Bohemia, Moravia), Sweden.

Genus: *Dicranotropis* FIEBER, 1866.

Type-species: *Delphax hamata* BOHEMAN, 1847, by subsequent designation by DISTANT, 1906.

*Leimonodite* KIRKALDY, 1907. Type-species: *Dicranotropis beckeri* FIEBER, 1866, by original designation.

*Dicranotropis beckeri* FIEBER, 1866.

Distribution: U.S.S.R. (Armenia, Kazakhstan, s.Russia, Tadzhikistan, Ukraine, Uzbekistan).

*Dicranotropis dimorpha* MATSUMURA, 1910.

Distribution: Italy (Sicily).

*Dicranotropis divergens* KIRSCHBAUM, 1868.

*Dicranotropis carpathica* HORVÁTH, 1884.

Distribution: Albania, Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), France, German D.R., German F.R., Great Britain (Scotland), Italy, Poland, Romania, Switzerland, U.S.S.R. (Caucasus, Kazakhstan, Ukraine).

*Dicranotropis esakii* ISHIHARA, 1949.

Distribution: Japan (Kyushu).

*Dicranotropis hamata* (BOHEMAN, 1847).

*Delphax hamata* BOHEMAN, 1847.

Distribution: Algeria, Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Hungary, Italy, Ireland, Netherlands, Norway, Poland, Romania, Spain, Sweden, Tunisia, U.S.S.R. (Altai Mts., Estonia, Georgia, Kazakhstan, Latvia, Moldavia, n., m. and s.Russia, m.Siberia, Ukraine), Yugoslavia.

*Dicranotropis montana* VILBASTE, 1965.

Distribution: U.S.S.R. (Altai Mts.).

*Dicranotropis tenellula* DLABOLA, 1965.

Distribution: Mongolia.

*Dicranotropis tikuzenensis* MATSUMURA et ISHIHARA, 1949.

Distribution: Japan (Kyushu, Shikoku).

*Dicranotropis zenata* LOGVINENKO, 1969.

Distribution: U.S.S.R. (Georgia).

Genus: *Zuleika* DISTANT, 1912.

Type-species: *Zuleika bengalensis* DISTANT, 1912, by monotypy.

*Zuleika nipponica* MATSUMURA et ISHIHARA, 1945.

*Zuleica* [sic] *nipponica* MATSUMURA et ISHIHARA, 1945.

Distribution: Japan (Honshu, Kyushu).

Genus: *Yanunka* ISHIHARA, 1952.

Type-species: *Yanunka miscanthi* ISHIHARA, 1952, by original designation.

*Yanunka miscanthi* ISHIHARA, 1952.

Distribution: Japan (Shikoku).

Genus: *Florodelphax* VILBASTE, 1968.

Type-species: *Delphax paryphasma* FLOR, 1861, by original designation.

*Florodelphax leptosoma* (FLOR, 1861).

*Delphax leptosoma* FLOR, 1861.

*Delphax albofimbriata* FIEBER, 1866.

*Delphax albolimbata* KIRSCHBAUM, 1868.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Hungary, Ireland, Italy, Madeira Archipelago, Netherlands, Poland, Portugal, Romania, Spain, Sweden, Switzerland, Turkey (Anatolia), U.S.S.R. (Estonia, Latvia, n.Russia, Ukraine), Yugoslavia.

*Florodelphax paryphasma* (FLOR, 1861).

*Delphax paryphasma* FLOR, 1861.

*Liburnia niveimarginata* SCOTT, 1870.

*Delphax lucticolor* J. SAHLBERG, 1868.

Distribution: Finland, German D.R., German F.R., Great Britain (England), U.S.S.R. (Estonia, Kazakhstan, Latvia, n.Russia).

Genus: *Kosswigianella* WAGNER, 1963.

Type-species: *Delphax exigua* BOHEMAN, 1847, by original designation.

*Kosswigianella exigua* (Boheman, 1847).

*Delphax exigua* BOHEMAN, 1847.

*Delphax limitata* FIEBER, 1866.

*Liburnia scutellata* SCOTT, 1873.

*Delphacodes exigua arida* METCALF, 1943.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Finland, France, German D.R., German F.R., Great Britain (England, Wales), Hungary, Italy, Japan (Honshu), Netherlands, Poland, Sweden, Switzerland, Tunisia, U.S.S.R. (Estonia, Latvia, Ukraine), Yugoslavia.

***Koswigianella guentheri*** DLABOLA, 1966.

*Koswigianella* [sic] *guentheri* DLABOLA, 1966.

Distribution: Mongolia.

Genus: ***Struebingianella*** WAGNER, 1963.

Type-species: *Delphax lugubrina* BOHEMAN, 1847, by original designation.

***Struebingianella dalei*** (SCOTT, 1870).

*Liburnia dalei* SCOTT, 1870.

*Liburnia dalei bohemani* [sic] FIEBER, 1879.

*Delphacodes dalei brunnea* METCALF, 1943.

Distribution: France, Great Britain (England).

***Struebingianella detecta*** (LINNAVUORI, 1953).

*Calligypona detecta* LINNAVUORI, 1953.

Distribution: U.S.S.R. (Maritime Territory, e.Siberia).

***Struebingianella litoralis*** (REUTER, 1880).

*Liburnia litoralis* REUTER, 1880.

Distribution: Finland, Great Britain (Scotland).

***Struebingianella lugubrina*** (BOHEMAN, 1847).

*Delphax lugubrina* BOHEMAN, 1847.

*Liburnia lugubrina melanoptera* KUNTZE, 1937.

*Liburnia lugubrina hyalipennis* KUNTZE, 1937.

*Delphacodes lugubrina albidoflavis* METCALF, 1943.

*Delphacodes lugubrina bimacula* METCALF, 1943.

*Delphacodes lugubrina piceonigra* METCALF, 1943.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), Hungary, Ireland, n.Italy, Netherlands, Poland, Sweden, Switzerland, U.S.S.R. (Estonia, Latvia, n.Russia, Ukraine), n.Yugoslavia.

***Struebingianella palliceps*** (HORVÁTH, 1897).

*Delphax palliceps* HORVÁTH, 1897.

Distribution: Bulgaria, Hungary, Italy, Romania, Yugoslavia.

Genus: ***Ferganodelphax*** DUBOVSKY, 1970.

Type-species: *Ferganodelphax crassiuscula* DUBOVSKY, 1970, by monotypy.

***Ferganodelphax crassiusculus*** DUBOVSKY, 1970.

*Ferganodelphax crassiuscula* DUBOVSKY, 1970.

Distribution: U.S.S.R. (Kirghizia).

Genus: *Kusnezoviella* VILBASTE, 1965.

Type-species: *Liburnia dimidiatifrons* KUSNEZOV, 1929, by original designation.

*Kusnezoviella dimidiatifrons* (KUSNEZOV, 1929).

*Liburnia dimidiatifrons* KUSNEZOV, 1929.

Distribution: Mongolia, U.S.S.R. (Altai Mts., m.Siberia).

*Kusnezoviella euagropyri* (EMELJANOV, 1964).

*Calligypona euagropyri* EMELJANOV, 1964.

Distribution: U.S.S.R. (Kazakhstan).

Genus: *Xanthodelphax* WAGNER, 1963.

Type-species: *Delphax flaveola* FLOR, 1861, by original designation.

*Xanthodelphax flaveolus* (FLOR, 1861).

*Delphax flaveola* FLOR, 1861.

*Delphacodes flaveola dorsoinfusca* METCALF, 1943.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England), Hungary, n.Italy, Norway, Poland, Romania, Sweden, U.S.S.R. (Estonia, Latvia, n.Russia, w.Siberia, Ukraine).

*Xanthodelphax stramineus* (STÅL, 1858).

*Delphax straminea* STÅL, 1858.

*Delphax fulveola* KIRSCHBAUM, 1868.

*Liburnia v-flava* SCOTT, 1881.

Distribution: Austria, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland), Hungary, n.Italy, Netherlands, Poland, Romania, Sweden, U.S.S.R. (Estonia, Kazakhstan, Latvia, Moldavia, n.Russia, Ukraine), Yugoslavia.

*Xanthodelphax xanthus* VILBASTE, 1965.

*Xanthodelphax xantha* VILBASTE, 1965.

*Xanthodelphax soosi* NAST, 1966.

Distribution: Poland, U.S.S.R. (Altai Mts., Kazakhstan).

Genus: *Paradelphacodes* WAGNER, 1963.

Type-species: *Delphax paludosa* FLOR, 1861, by original designation.

*Paradelphacodes paludosa* (FLOR, 1861).

*Delphax paludosa* FLOR, 1861.

*Delphacodes kuvaharai* ISHIHARA, 1949.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia), Finland, France, German D.R., German F.R., Great Britain (England), Italy, Japan (Honshu), Mongolia, Netherlands, Poland, Sweden, U.S.S.R. (Estonia, Latvia, Maritime Territory, n.Russia).

*Paradelphacodes tengaica* VILBASTE, 1965

Distribution: U.S.S.R. (Altai Mts.).

Genus: *Oncodelphax* WAGNER, 1963

Type-species: *Delphax pullula* BOHEMAN, 1852, by original designation.

*Oncodelphax pullulus* (BOHEMAN, 1852).

*Delphax pullula* BOHEMAN, 1852.

*Delphacodes pullula basiflavis* METCALF, 1943.

*Delphacodes pullula infuscata* METCALF, 1943.

Distribution: Austria, Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland), Netherlands, Poland, Sweden, Switzerland, U.S.S.R. (Estonia, n.Russia), Yugoslavia.

Genus: *Criomorphus* CURTIS, 1833.

Type-species: *Criomorphus albomarginatus* CURTIS, 1833, by original designation.

*Criomorphus albomarginatus* CURTIS, 1833.

*Delphax collaris* STÅL, 1853.

*Delphax adelpha* FLOR, 1861.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Ireland, Israel, Netherlands, Norway, Poland, Romania, Sweden, Switzerland, Tunisia, U.S.S.R. (Estonia, Latvia, n. and m.Russia, Ukraine).

*Criomorphus borealis* (J. SAHLBERG, 1871).

*Ditropis borealis* J. SAHLBERG, 1871.

*Stiroma borealis carinata* METCALF, 1943.

Distribution: Czechoslovakia ([?] Bohemia), Finland, German D.R., Mongolia, Norway, Poland, Sweden, U.S.S.R. (Estonia, Latvia, n.Russia, n.w.Siberia).

*Criomorphus moestus* (BOHEMAN, 1847).

*Delphax moesta* BOHEMAN, 1847.

*Delphax thoracica* STÅL, 1858.

*Stiroma germanica* MATSUMURA, 1910.

Distribution: Czechoslovakia (Bohemia), Finland, France, German D.R., Great Britain (England), Sweden, U.S.S.R. (Estonia, Latvia, n.Russia).

*Criomorphus nigerrimus* DLABOLA, 1965.

Distribution: Mongolia.

*Criomorphus williamsi* CHINA, 1939.

Distribution: Great Britain (England), U.S.S.R. (Kazakhstan).

Genus: *Pseudodelphacodes* WAGNER, 1963.

Type-species: *Delphax flaviceps* FIEBER, 1866, by original designation.

***Pseudodelphacodes flaviceps* (FIEBER, 1866).**

*Delphax flaviceps* FIEBER, 1866.

Distribution: Austria, German F.R., Switzerland, Yugoslavia.

Genus: *Falcotoya* FENNAH, 1969.

Type-species: *Falcotoya aurinia* FENNAH, 1969, by original designation.

***Falcotoya lyraeformis* (MATSUMURA, 1900).**

*Liburnia lyraeformis* MATSUMURA, 1900.

Distribution: Japan (Honshu, Kyushu, Shikoku).

Genus: *Toya* DISTANT, 1906.

Type-species: *Toya attenuata* DISTANT, 1906, by original designation.

*Metadelphax* WAGNER, 1963. Type-species: *Delphax propinqua* FIEBER, 1866, by original designation.

***Toya minuscula* (HORVÁTH, 1897).**

*Delphax minuscula* HORVÁTH, 1897.

Distribution: Afghanistan, Czechoslovakia (Slovakia), Egypt, Hungary, Iran, Romania, Turkey (Anatolia), U.S.S.R. (Armenia, Kazakhstan, Tadzhikistan), Yugoslavia.

***Toya propinqua* (FIEBER, 1866).**

*Delphax propinqua* FIEBER, 1866.

*Delphax hamulata* KIRSCHBAUM, 1868.

Distribution: Afghanistan, Albania, Algeria, Austria, Bulgaria, Canary Is., Cyprus, Czechoslovakia (Bohemia, Moravia, Slovakia), Egypt, France, German D.R., German F.R., Hungary, Iran, Iraq, Israel, Italy (also Sicily), Japan (Honshu), Jordan, Libya, Madeira Archipelago, Morocco, Poland, Portugal, Romania, Spain, Spanish Sahara, Tunisia, Turkey (Anatolia), U.S.S.R. (Armenia, Georgia, Kazakhstan, Kirghizia, Moldavia, s.Russia, Tadzhikistan, Uzbekistan), Yugoslavia; [Ethiopian, [?] Nearctic and [?] Oriental regions].

***Toya simulans* (DLABOLA, 1958).**

*Calligypona simulans* DLABOLA, 1958.

Distribution: U.S.S.R. (Georgia, Kazakhstan, Tadzhikistan, Ukraine).

***Toya tricolorata* (DLABOLA, 1961).**

*Calligypona tricolorata* DLABOLA, 1961.

Distribution: U.S.S.R. (Kazakhstan, Tadzhikistan).

***Toya yanoi* (ISHIHARA, 1952).**

*Sogata yanoi* ISHIHARA, 1952.

Distribution: Japan (Shikoku).

Genus: *Terthron* FENNAH, 1965.

Type-species: *Delphax anemonias* KIRKALDY, 1907, by original designation.

*Terthron albovittatum* (MATSUMURA, 1900).

*Dicranotropis albovittata* MATSUMURA, 1900.

Distribution: Japan (Honshu, Kyushu, Shikoku), Korean Peninsula; [Oriental region].

Genus: *Terthronella* VILBASTE, 1968.

Type-species: *Liburnia basalis* MATSUMURA, 1915, by original designation.

*Terthronella basalis* (MATSUMURA, 1915).

*Liburnia (Delphax) basalis* MATSUMURA, 1915.

Distribution: Korean Peninsula, U.S.S.R. (Maritime Territory).

Genus: *Harmalia* FENNAH, 1969.

Type-species: *Sogata thoracica* DISTANT, 1916, by original designation.

*Harmalia albicollis* (MOTSCHULSKY, 1863).

*Delphax albicollis* MOTSCHULSKY, 1863.

*Delphacodes shirozui* ISHIHARA, 1949.

Distribution: Japan (Kyushu, Shikoku), U.S.S.R. (Maritime Territory); [Oriental region].

*Harmalia sameshimai* (MATSUMURA et ISHIHARA, 1945).

*Kakuna sameshimai* MATSUMURA et ISHIHARA, 1945.

Distribution: Japan (Kyushu).

Genus: *Javesella* FENNAH, 1963.

Type-species: *Fulgora pellucida* FABRICIUS, 1794, by original designation.

*Weidnerianella* WAGNER, 1963. Type-species: *Fulgora pellucida* FABRICIUS, 1794, by original designation.

*Haffnerianella* WAGNER, 1966. [Subgenus of *Javesella*]. Type-species: *Delphacodes stali* METCALF, 1943, by original designation.

*Javesella discolor* (BOHEMAN, 1847).

*Delphax discolor* BOHEMAN, 1847.

*Delphax similis* KIRSCHBAUM, 1868.

*Delphax nitidula* KIRSCHBAUM, 1868.

*Delphax patens* KIRSCHBAUM, 1868.

Distribution: Algeria, Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Ireland, Italy (also Sicily), Mongolia, Netherlands, Norway, Poland, Romania, Sweden, Switzerland, U.S.S.R. (Estonia, Latvia, n.Russia, n. and m.Siberia, Ukraine).

***Javesella dubia* (KIRSCHBAUM, 1868).**

*Delphax dubia* KIRSCHBAUM, 1868.  
*Delphax herrichi* KIRSCHBAUM, 1868, partim.  
*Delphax nitidipennis* KIRSCHBAUM, 1868.  
*Liburnia pargasensis* REUTER, 1880.  
*Liburnia difficilis* EDWARDS, 1888.

Distribution: Austria, Azores, Belgium, Bulgaria, Cyprus, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland), Ireland, Italy, Madeira Archipelago, Morocco, Netherlands, Norway, Poland, Romania, Sweden, U.S.S.R. (Altai Mts., Estonia, Kazakhstan, Latvia, Moldavia, n. and m.Russia, Ukraine, Uzbekistan).

***Javesella eforiae* (DLABOLA, 1967).**

*Calligypona eforiae* DLABOLA, 1967.

Distribution: Mongolia, Romania, U.S.S.R. (Kazakhstan).

***Javesella forcipata* (BOHEMAN, 1847).**

*Delphax forcipata* BOHEMAN, 1847.  
*Liburnia alpina* J. SAHLBERG, 1871.  
*Delphacodes alpina dorsoflava* METCALF, 1943.  
*Delphacodes alpina griseoflavescens* METCALF, 1943.  
*Delphacodes alpina grisescens* METCALF, 1943.  
*Delphacodes alpina nigrovariegata* METCALF, 1943.  
*Delphacodes alpina tota* METCALF, 1943.  
*Delphacodes forcipata angustata* METCALF, 1943.  
*Delphacodes forcipata dorsigera* METCALF, 1943.  
*Delphacodes forcipata rufescens* METCALF, 1943.

Distribution: Austria, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Ireland, Italy, Netherlands, Norway, Poland, Romania, Sweden, U.S.S.R. (Estonia, Latvia, n. and m.Russia, n.Siberia, Ukraine).

***Javesella nuchtica* DLABOLA, 1967.**

Distribution: Mongolia.

***Javesella obscurella* (BOHEMAN, 1847).**

*Delphax obscurella* BOHEMAN, 1847.  
*Liburnia discreta* EDWARDS, 1888.  
*Delphacodes obscurella fuscotestacea* METCALF, 1943.

Distribution: Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Hungary, Italy (also Sardinia), Ireland, Mongolia, Netherlands, Norway, Poland, Portugal, Romania, Sweden, Switzerland, Turkey (Anatolia), U.S.S.R. (Estonia, Georgia, Latvia, Moldavia, n. and s.Russia, n.Siberia, Ukraine), Yugoslavia; [Nearctic region].

***Javesella pellucida*** (FABRICIUS, 1794).

- Fulgora pellucida* FABRICIUS, 1794.  
*Fulgora striata* FABRICIUS, 1794.  
*Fulgora marginata* FABRICIUS, 1794.  
*Delphax dispar* FALLÉN, 1806.  
*Delphax suturalis* CURTIS, 1837.  
*Delphax herrichii* KIRSCHBAUM, 1868, partim.  
*Delphax ochroleuca* KIRSCHBAUM, 1868.  
*Delphax frontalis* KIRSCHBAUM, 1868.  
*Liburnia flavipennis* J. SAHLBERG, 1871.  
*Liburnia pellucida obscura* FIEBER, 1879.  
*Delphax pellucida fuscicollis* REY, 1891.  
*Delphax pellucida varicollis* REY, 1894.  
*Delphacodes pellucida bilineata* METCALF, 1943.  
*Delphacodes pellucida flavotestacea* METCALF, 1943.  
*Delphacodes pellucida griseoflava* METCALF, 1943.  
*Delphacodes pellucida testacea* METCALF, 1943.

Distribution: Algeria, Austria, Azores, Belgium, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Great Britain (England, Scotland, Wales), Hungary, Ireland, Iceland, Italy, Libya, Mongolia, Morocco, Netherlands, Norway, Poland, Romania, Spain, Sweden, Switzerland, Turkey (Anatolia), U.S.S.R. (Altai Mts., Estonia, Georgia, Kamchatka, Kazakhstan, Kirghizia, Kurile Is., Latvia, Maritime Territory, Moldavia, n., m. and s.Russia, Sakhalin, Siberia, Ukraine, Uzbekistan), Yugoslavia.

***Javesella salina*** (HAUPT, 1924).

- Delphax salina* HAUPT, 1924.  
*Liburnia juncea* HAUPT, 1935.

Distribution: Czechoslovakia (Slovakia), German D.R., German F.R., Mongolia, Poland, Sweden, Turkey (Anatolia), U.S.S.R. (Altai Mts., Estonia, Maritime Territory).

***Javesella selengica*** DLABOLA, 1970.

Distribution: Mongolia.

***Javesella simillima*** (LINNAVUORI, 1948).

- Calligypona simillima* LINNAVUORI, 1948.

Distribution: Finland, U.S.S.R. (Estonia, n.Siberia <Taymyr>).

***Javesella stali*** (METCALF, 1943).

- Delphax bohemani* STÅL, 1858. [Primary homonym].  
*Delphacodes stali* METCALF, 1943.  
*Delphacodes stali flavalis* METCALF, 1943.

Distribution: Austria, France, German D.R., [?] Italy, Poland, Sweden, U.S.S.R. (Estonia, Latvia, n.Russia, Ukraine).

Genus: *Ribautodelphax* WAGNER, 1963.

Type-species: *Delphax collina* BOHEMAN, 1847, by original designation.

***Ribautodelphax affinis* LOGVINENKO, 1970.**

Distribution: U.S.S.R. (Georgia, Ukraine <Crimea>).

***Ribautodelphax albostriatus* (FIEBER, 1866).**

*Delphax albostriata* FIEBER, 1866.

*Delphax distinguenda* KIRSCHBAUM, 1868.

Distribution: Austria, Belgium, Cyprus, Czechoslovakia (Bohemia, Moravia, Slovakia), Denmark, Finland, France, German D.R., German F.R., Hungary, Italy, Mongolia, Netherlands, Norway, Poland, Portugal, Romania, Spain, Sweden, Switzerland, Tunisia, U.S.S.R. (Caucasus, Estonia, Kazakhstan, Kirghizia, Moldavia, n. and m.Russia, m.Siberia, Ukraine), Yugoslavia.

***Ribautodelphax altaicus* VILBASTE, 1965.**

*Ribautodelphax altaica* VILBASTE, 1965.

Distribution: U.S.S.R. (Altai Mts., Kazakhstan).

***Ribautodelphax angulosus* (RIBAUT, 1953).**

*Calligypona angulosa* RIBAUT, 1953.

Distribution: Austria, Finland, France, German D.R., German F.R., Great Britain (England), Mongolia, Netherlands, Poland, Romania, Sweden, Switzerland, U.S.S.R. (Kazakhstan, Latvia, Moldavia, Ukraine).

***Ribautodelphax bicolor* LOGVINENKO, 1970.**

Distribution: U.S.S.R. (Ukraine <Crimea>).

***Ribautodelphax bidentatus* ANUFRIEV, 1970.**

Distribution: U.S.S.R. (Maritime Territory).

***Ribautodelphax bogdul* DLABOLA, 1968.**

Distribution: Mongolia.

***Ribautodelphax collinus* (BOHEMAN, 1847).**

*Delphax collina* BOHEMAN, 1847.

*Delphax concinna* FIEBER, 1866. [Primary homonym].

*Liburnia biarmica* J. SAHLBERG, 1871.

*Delphacodes annae* KIRKALDY, 1906.

Distribution: Austria, Belgium, Bulgaria, Czechoslovakia (Bohemia, Moravia, Slovakia), Finland, France, German D.R., German F.R., Italy, Netherlands, Poland, Sweden, U.S.S.R. (Estonia, Kazakhstan, Latvia, n.Russia, Ukraine), Yugoslavia.

***Ribautodelphax exquisitus* ANUFRIEV, 1970.**

*Ribautodelphax exquisita* ANUFRIEV, 1970.

Distribution: U.S.S.R. (m.Siberia <Chita distr.>).

***Ribautodelphax flavicans* VILBASTE, 1965.**

Distribution: Mongolia, U.S.S.R. (Altai Mts.).

***Ribautodelphax imitans* (RIBAUT, 1953).**

*Calligypona imitans* RIBAUT, 1953.

Distribution: Austria, Czechoslovakia (Slovakia), France, German F.R., Great Britain (England), Italy, Netherlands, Poland, Romania.

***Ribautodelphax notabilis* LOGVINENKO, 1970.**

Distribution: U.S.S.R. (Ukraine <Crimea>).

***Ribautodelphax ochreateus* VILBASTE, 1965.**

*Ribautodelphax ochreatea* VILBASTE, 1965.

Distribution: Mongolia, U.S.S.R. (Altai Mts., Kirghizia).

***Ribautodelphax pallens* (STÅL, 1854).**

*Delphax pallens* STÅL, 1854.

Distribution: Austria, Finland, German F.R., Great Britain (England), Norway, Sweden, U.S.S.R. (Estonia, Kaliningrad distr., Latvia, n.Russia).

***Ribautodelphax pungens* (RIBAUT, 1953).**

*Calligypona pungens* RIBAUT, 1953.

Distribution: Austria, Czechoslovakia (Bohemia, Moravia), France, German D.R., German F.R., Great Britain (England), Italy, Netherlands, Poland.

***Ribautodelphax vicinus* (LINNAVUORI, 1953).**

*Calligypona vicina* LINNAVUORI, 1953.

Distribution: Mongolia, U.S.S.R. (w.Siberia).

Genus: *Phyllodinus* VAN DUZEE, 1897.

Typ-species: *Phyllodinus nervata* VAN DUZEE, 1897, by monotypy.

*Jamiphax* MATSUMURA, 1940. [Preoccupied]. Type-species: *Jamiphax kotoshonis* MATSUMURA, 1940, by original designation.

***Phyllodinus nigropunctatus* (MOTSCHULSKY, 1863).**

*Mestus nigropunctatus* MOTSCHULSKY, 1863.

Distribution: Japan (Honshu, Kyushu, Shikoku); [Oriental region].

Genus: *Pseudaraeopus* KIRKALDY, 1904.

Type-species: *Araeopus lethierryi* MULSANT et REY, 1879, by original designation.

*Delphacodes* MELICHAR, 1901. [Preoccupied]. Type-species: *Araeopus lethierryi* MULSANT et REY, 1879, by subsequent designation by OSHANIN, 1912.

*Tricranoneura* HORVÁTH, 1922. [Subgenus of *Pseudaraeopus*]. Type-species: *Delphacodes bolivari* MELICHAR, 1901, by monotypy.

***Pseudaraeopus bolivari* (MELICHAR, 1901).**

*Delphacodes bolivari* MELICHAR, 1901.

Distribution: Canary Is., Egypt, Iran, Israel, Portugal, Spain.

***Pseudaraeopus curtulus* DLABOLA, 1960.**

Distribution: Iran.

*Pseudaraeopus dalmatinus* HORVÁTH, 1922.

Distribution: Yugoslavia.

*Pseudaraeopus frontalis* (LINDBERG, 1958).

*Bostaera frontalis* LINDBERG, 1958.

Distribution: Palestine; [Ethiopian region].

*Pseudaraeopus iranicus* DLABOLA, 1960.

Distribution: Iran.

*Pseudaraeopus lethierryi* (MULSANT et REY, 1879).

*Araeopus lethierryi* MULSANT et REY, 1879.

*Stiroma ribauti* LALLEMAND, 1921.

Distribution: Canary Is., Cyprus, France, Israel, Madeira Archipelago, Yugoslavia.

Genus: *Nagara* VILBASTE, 1968.

Type-species: *Liburnia nagaragawana* MATSUMURA, 1900, by original designation.

*Nagara nagaragawana* (MATSUMURA, 1900).

*Liburnia nagaragawana* MATSUMURA, 1900.

Distribution: Japan (Honshu, Kyushu, Shikoku), U.S.S.R. (Maritime Territory); [Oriental region].

Genus: *Cemus* FENNAH, 1964.

Type-species: *Cemus leviculus* FENNAH, 1964, by original designation.

*Cemus nigromaculosus* (MUIR, 1917).

*Phyllostinus nigromaculosus* MUIR, 1917.

Distribution: Japan (Honshu, Kyushu); [Oriental region].

Genus: *Peregrinus* KIRKALDY, 1904.

Type-species: *Delphax maidis* ASHMEAD, 1890, by original designation.

*Peregrinus maidis* (ASHMEAD, 1890).

*Delphax maidis* ASHMEAD, 1890.

Distribution: Canary Is.; [Oriental, Ethiopian, Nearctic and Neotropic regions].

Genus: *Euidopsis* RIBAUT, 1948.

Type-species: *Euidopsis truncata* RIBAUT, 1948, by original designation.

*Euidopsis truncata* RIBAUT, 1948.

Distribution: Cyprus.

Genus: *Alatades* DLABOLA, 1957.

Type-species: *Alatades trilineatus* DLABOLA, 1957, by original designation.

*Alatades trilineatus* DLABOLA, 1957.

Distribution: Israel, Turkey (Anatolia).

Genus: *Perkinsiella* KIRKALDY, 1903.

Type-species: *Perkinsiella saccharicida* KIRKALDY, 1903, by original designation.

*Araeopides* RIBAUT, 1948. Type-species: *Araeopides picta* RIBAUT, 1948, by original designation.

*Perkinsiella insignis* (DISTANT, 1912).

*Pundaluoya insignis* DISTANT, 1912.

*Araeopides picta* RIBAUT, 1948.

Distribution: Afghanistan, Cyprus, Iraq, Israel, Turkey (Anatolia); [Oriental and Ethiopian regions].

*Perkinsiella rivularis* LINNAVUORI, 1964.

Distribution: Egypt, Israel.

*Perkinsiella sinensis* KIRKALDY, 1907.

Distribution: Japan (Honshu, Kyushu); [Oriental region].

*Perkinsiella yakushimensis* ISHIHARA, 1954.

Distribution: Japan (Yakushima).

Genus: *Hirozunka* MATSUMURA et ISHIHARA, 1945.

Type-species: *Hirozunka japonica* MATSUMURA et ISHIHARA, 1945, by original designation.

*Hirozunka japonica* MATSUMURA et ISHIHARA, 1945.

Distribution: Japan (Honshu, Kyushu).

#### **DELPHACIDAE OF UNCERTAIN GENERIC POSITION.**

The species are arranged in alphabetic order of their specific names (**bold-faced**).

*Delphacodes agropyri* ISHIHARA, 1949.

Distribution: Japan (Honshu).

*Liburnia angara* KUSNEZOV, 1929.

Distribution: U.S.S.R. (m.Siberia).

*Liburnia anthracina* HORVÁTH, 1909.

Distribution: Canary Is., Madeira Archipelago; [Ethiopian region].

- Calligypona appropinqua* DLABOLA, 1961.  
Distribution: U.S.S.R. (Tadzhikistan, Uzbekistan).
- Liburnia axillaris* J. SAHLBERG, 1876.  
Distribution: Finland.
- Calligypona bella* LINNAVUORI, 1964.  
Distribution: Egypt.
- Calligypona bifurcata* LINDBERG, 1961.  
Distribution: Madeira Archipelago, Portugal.
- Calligypona brunneola* DUBOVSKY, 1966.  
Distribution: U.S.S.R. (Kirghizia).
- Liburnia burjata* KUSNEZOV, 1929.  
Distribution: U.S.S.R. (m.Siberia).
- Delphax cataniae* MATSUMURA, 1910.  
Distribution: Italy (Sicily).
- Calligypona cochlearia* EMELJANOV, 1964.  
Distribution: U.S.S.R. (Kazakhstan).
- Delphax conspicua* HORVÁTH, 1904.  
Distribution: U.S.S.R. (Kazakhstan).
- Liburnia coracina* CSIKI, 1940.  
Distribution: Albania.
- Liburnia cornigera* KUSNEZOV, 1929.  
Distribution: U.S.S.R. (m.Siberia).
- Delphacodes dogensis* ISHIHARA, 1952.  
Distribution: Japan (Shikoku).
- Calligypona fascia* LINDBERG, 1960.  
Distribution: Portugal.
- Delphacodes flava* METCALF, 1943.  
*Delphax flavescens* IVANOFF, 1885. [Primary homonym].  
Distribution: U.S.S.R. (Ukraine).
- Calligypona flavobrunnea* DLABOLA, 1956.  
Distribution: Italy.
- Delphacodes fukuokae* ISHIHARA, 1949.  
Distribution: Japan (Kyushu).
- Calligypona fumata* LINDBERG, 1948.  
Distribution: Bulgaria, Cyprus.

- Delphax fumipennis* FIEBER, 1866.  
Distribution: France, Greece.
- Liburnia fuscifrons* FIEBER, 1879.  
Distribution: France, Belgium.
- Delphax gilveola* KIRSCHBAUM, 1868.  
Distribution: German F.R.
- Liburnia (Delphax) gracilis* MATSUMURA, 1915.  
Distribution: Korean Peninsula.
- Liburnia guaramensis* SCOTT, 1873.  
Distribution: Spain.
- Calligypona gudruna* GYLLENSVÄRD, 1967.  
Distribution: Canary Is.
- Delphax hemiptera* GERMAR, 1818.  
Distribution: no locality given.
- Calligypona hispidula* LINDBERG, 1954.  
Distribution: Canary Is., Egypt, Madeira Archipelago, Morocco.
- Calligypona isis* LINNAVUORI, 1958.  
Distribution: Egypt.
- Calligypona kirghisica* DUBOVSKY, 1966.  
Distribution: U.S.S.R. (Kirghizia).
- Liburnia latifrons* FIEBER, 1879.  
Distribution: France, Germany.
- Liburnia lethierryi* SCOTT, 1873.  
Distribution: Italy (Sardinia), France (Corsica).
- Delphax macroptera* O. COSTA, 1834.  
Distribution: Italy.
- Calligypona maculipennis* LINNAVUORI, 1962.  
Distribution: Israel.
- Liburnia marshalli* SCOTT, 1873.  
Distribution: France (Corsica).
- Liburnia matsumurana* METCALF, 1943.  
*Delphax furcata* MATSUMURA, 1910. [Primary homonym].  
Distribution: Italy (Sicily).
- Delphacodes matsuyamensis* ISHIHARA, 1952.  
Distribution: Japan (Shikoku).

- Liburnia melanocephala* FIEBER, 1879.  
Distribution: German F.R., U.S.S.R. (Ukraine).
- Calligypona minutissima* DUBOVSKY, 1966.  
Distribution: U.S.S.R. (Fergana Valley).
- Delphax mocsaryi* HORVÁTH, 1897.  
Distribution: Hungary.
- Delphax modesta* FIEBER, 1866.  
Distribution: France.
- Liburnia mongolica* KUSNEZOV, 1929.  
Distribution: U.S.S.R. (m.Siberia).
- Calligypona negrita* DLABOLA, 1961.  
Distribution: U.S.S.R. (Tadzhikistan).
- Delphacodes nigerrima* ISHIHARA, 1949.  
Distribution: Japan (Honshu).
- Delphacodes nigrella* ISHIHARA, 1954.  
Distribution: Japan (Kyushu).
- Delphax nigricans* MATSUMURA, 1910.  
Distribution: Italy (Sicily), Morocco.
- Delphax nigrifrons* MATSUMURA, 1910.  
Distribution: Italy (Sicily).
- Delphacodes nigrigena* MATSUMURA et ISHIHARA, 1945.  
Distribution: Japan (Kyushu).
- Liburnia nigrostriata* KUSNEZOV, 1929.  
Distribution: U.S.S.R. (m.Siberia).
- Calligypona obtusangula* LINNAVUORI, 1957.  
Distribution: Cyprus, Israel, Italy.
- Megamelus ornatipennis* HAUPT, 1927.  
Distribution: Afghanistan, Iran, Israel, Jordan.
- Delphax radiata* O. COSTA, 1834.  
Distribution: Italy.
- Liburnia segetum* HAUPT, 1927.  
Distribution: Palestine.
- Delphacodes sporoboli* LINDBERG, 1958.  
Distribution: Egypt; [Ethiopian region].
- Delphax stejnegeri* ASHMEAD, 1899.  
Distribution: U.S.S.R. (Commander Is.); [Nearctic region].

- Liburnia stigmatica* MELICHAR, 1897.  
Distribution: Italy, Yugoslavia.
- Delphax stigmatalis* CURTIS, 1837.  
Distribution: Great Britain.
- Delphax strigosa* MATSUMURA, 1910.  
Distribution: Italy (Sicily).
- Delphax tangira* MATSUMURA, 1910.  
Distribution: Morocco.
- Delphax tapina* FIEBER, 1866.  
Distribution: France, Denmark, Italy (Sardinia).
- Delphacodes terryi* MUIR, 1917.  
Distribution: Japan (Kyushu); [Oriental region].
- Calligypona typhae* LINDBERG, 1960.  
Distribution: Canary Is., Egypt, Israel, Morocco.
- Delphax uncinata* FIEBER, 1866.  
Distribution: Great Britain, Greece, Hungary, Portugal.
- Calligypona urceolata* DUBOVSKY, 1966.  
Distribution: U.S.S.R. (Kirghizia).
- Calligypona uzbekistanica* DUBOVSKY, 1966.  
Distribution: U.S.S.R. (Uzbekistan).
- Delphacodes walkeri* METCLAF, 1943.  
*Delphax dorsalis* WALKER, 1870. [Primary homonym].  
Distribution: Egypt.
- Liburnia yezoana* MATSUMURA, 1900.  
Distribution: Japan (Hokkaido, Honshu, Shikoku).
- Calligypona zeravshanica* DUBOVSKY, 1967.  
Distribution: U.S.S.R. (Uzbekistan).

Family: **MEENOPLIDAE** FIEBER, 1872.

Genus: *Nisia* MELICHAR, 1903.

Type-species: *Meenoplus atrovexus* LETHIERRY, 1888, by monotypy.

*Nisia atrovexa* (LETHIERRY, 1888).

*Meenoplus atrovexus* LETHIERRY, 1888.

Distribution: China (Chekiang), Egypt, Israel, Japan (Honshu, Kyushu, Shikoku), Korean Peninsula, Morocco, Palestine, Spanish Sahara; [Oriental, Ethiopian and Australian regions].