



Figs 1–3. *Thionia simplex* (Germar, 1830), male, Italy. **1**, dorsal view; **2**, lateral view; **3**, frontal view. Total length of specimen is 5.2 mm.

First record of Nearctic issid planthopper *Thionia simplex* (Hemiptera: Fulgoroidea: Issidae) from Europe

Первое указание неарктической иссиды *Thionia simplex* (Hemiptera: Fulgoroidea: Issidae) из Европы

V.M. GNEZDILOV* & F. POGGI

В.М. ГНЕЗДИЛОВ & Ф. ПОДЖИ

V.M. Gnezdilov, Zoological Institute of the Russian Academy of Sciences, 1 Universitetskaya Emb., St Petersburg 199034, Russia. E-mails: vmgnezdilov@mail.ru, vgnezdilov@zin.ru

F. Poggi, Via Madonnina 6, I-23873 Missaglia (LC), Italia. E-mail: erythria@gmail.com

The Nearctic issid species *Thionia simplex* (Germar, 1830) is recorded for the first time from Europe. Other alien Auchenorrhyncha species in Europe are listed and discussed.

Неарктическая иссида *Thionia simplex* (Germar, 1830) впервые указана из Европы. Перечислены и обсуждены другие случаи завозов в Европу цикадовых.

Key words: alien species, U.S.A., Europe, Italy, Auchenorrhyncha, Fulgoroidea, Issidae, Issini, planthopper, new record

Ключевые слова: инвазивный вид, США, Европа, Италия, Auchenorrhyncha, Fulgoroidea, Issidae, Issini, фулгориодные цикадовые, новое указание

INTRODUCTION

Italy has become the “gateway for New World planthoppers in Europe” since the last century as several species which are adventive for Europe were first recorded in this country, e.g. *Metcalfa pruinosa* (Say, 1830) from Flatidae and *Acanalonia conica* (Say, 1830) from Acanaloniidae (Zangheri & Donadini, 1980; D’Urso & Uliana, 2006). Also a ricaniid species *Ricania speculum* (Walker, 1851) from the Orient was first recorded for Europe in Italy (Mazza et al., 2014).

Below we report the first representative of the Nearctic Issidae, *Thionia simplex* (Germar, 1830), found in Europe. This species was first collected in Lombardia in 2012 and identified using illustrations and

description (including of the male genitalia) given by Doering (1938).

Thionia simplex (Germar, 1830) was described from Kentucky in U.S.A. (Germar, 1830). Currently this species is recorded from 19 states in eastern U.S.A. (Bartlett et al., 2014). The genus *Thionia* Stål, 1859 belongs to the tribe Issini Spinola, 1839 (= Thioniini Melichar, 1906) sensu Gnezdilov (2009) and currently comprises 72 extant species, most of them distributed in the Neotropic Realm, with just eight species known from the Nearctic Realm (U.S.A.) (Gnezdilov, 2013; Bartlett et al., 2014).

Thionia simplex is known as a polyphagous species in the U.S.A., where it has been collected from 12 host plants in ten families (Wheeler & Wilson, 1988). It may become a serious pest in Europe as has already happened with *Metcalfa pruinosa* which spread from Italy east to southern Russia and Republic of Korea and west to Spain and has

* Corresponding author.

became a serious pest on many plants (Lucchi, 1997; Gnezdilov & Sugonyaev, 2009; Kim et al., 2011; Preda & Skolka, 2011).

MATERIAL AND METHODS

The material examined is deposited in the private collection of Mr Francesco Poggi, Missaglia, Italy (CFP) and in the Zoological Institute of the Russian Academy of Sciences, Saint Petersburg, Russia (ZIN).

The photos were taken by Leica MZ 95 with camera Leica DFC 290 and then assembled with Helicon Focus 5.3 and Adobe Photoshop CS6.

Superfamily **FULGOROIDEA**

Family **ISSIDAE**

Subfamily **ISSINAE**

Tribe **ISSINI**

Genus *Thionia* Stål, 1859

Thionia simplex (Germar, 1830)
(Figs 1–3)

Material examined. **Italy, Lombardia**, 20 km N of Milan, Marezzo (LC), 350 m: 3 females, 2.IX.2012, F. Poggi leg. (CFP); 2 males, 1 female, 15.IX.2012, F. Poggi leg. (CFP); 5 males, 6.IX.2014, F. Poggi leg. (ZIN); 5 males, 1 female, 6.IX.2014, F. Poggi leg. (CFP).

Remarks. Nine males and one female from the material listed above were collected 6 September 2014 on a yellow outside house wall, one male was collected 6 September 2014 at light, and two males and four females were collected since 2 to 15 September 2012 on *Ulmus laevis* Pall. No nymphs were collected.

DISCUSSION

Transcontinental invasions are very exceptional for the Issidae as a whole. Up to now there is just one issid species which has crossed the Atlantic Ocean. This South European species, *Agalmatium bilobum* (Fieber, 1877), was found in California, U.S.A. Probably it has been introduced with grapevines

(Gnezdilov & O'Brien, 2006). One Western Palaearctic species, *Issus coleoptratus* (Fabricius, 1781), has been found on Saint Helena Island in the Atlantic Ocean (Synave, 1976), and the Indo-Malayan *Thabena brunifrons* (Bonfils, Attie et Reynaud, 2001) has been found on Reunion Island in the Indian Ocean (Bonfils et al., 2001; Gnezdilov, 2009, 2013). Two more Indo-Malayan species formally described from Europe (Fieber, 1876; Gnezdilov et al., 2011), *Thabena fisisala* (Fieber, 1876) and *Thabenula sulcata* (Fieber, 1876), either resulted from labeling errors or reflected actual cases of movement of Indo-Malayan species to Europe by sea (Gnezdilov, 2013). Other planthopper, the species of the family Caliscelidae Amyot et Serville, *Caliscelis bonellii* (Latreille, 1807), described from Italy, also was introduced from Southern Europe to California, U.S.A. (O'Brien, 1967).

In Italy *Thionia simplex* (Germar) was collected on *Ulmus laevis* Pall. (Ulmaceae). In the U.S.A. this species was also recorded from an elm, *Ulmus alata* Michx. (Wheeler & Wilson, 1988). However during two years of an observation (September 2012 – September 2014) no nymphs were found in Italy at the place of invasion. This can indicate several invasions by imago from the U.S.A. to Italy with any cargos, probably some ornamental plants, without full development and hibernation in Europe. However it can also be caused by a lack of effective methods for finding nymphs. It is noteworthy that the collecting locality of *T. simplex* is the same as for another Nearctic Auchenorrhyncha, the leafhopper from Cicadellidae: Coelidiinae, *Jikradia olitoria* (Say, 1830), also recently found in Europe (Nielson et al., 2014). Further study is needed to investigate the life cycle of *T. simplex* and its possible pest importance in Italy and Europe.

ACKNOWLEDGEMENTS

We are glad to thank Dr Lois O'Brien (Tucson, AZ, U.S.A.) and Dr Charles Bartlett (Newark, DE, U.S.A.) for their valuable comments on the manuscript.

REFERENCES

- Bartlett C.R., O'Brien L.B. & Wilson S.W.** 2014. A review of the planthoppers (Hemiptera: Fulgoroidea) of the United States. *Memoires of the American Entomological Society*, **50**: 1–187.
- Bonfils J., Attié M. & Reynaud B.** 2001. Un nouveau genre d'Issidae de l'île de la Réunion: *Borbonissus* n. gen. (Hemiptera, Fulgoromorpha). *Bulletin de la Société entomologique de France*, **106**(3): 217–224.
- Doering K.C.** 1938. A contribution to the taxonomy of the subfamily Issinae in America North of Mexico (Fulgoroidea, Homoptera). Part II. *The University of Kansas Science Bulletin*, **25**(20): 447–575.
- D'Urso V. & Uliana M.** 2006. *Acanalonia conica* (Hemiptera, Fulgoromorpha, Acanaloniidae), a Nearctic species recently introduced in Europe. *Deutsche Entomologische Zeitschrift*, **53**(1): 103–107.
- Fieber F.-X.** 1876. Les Cicadines d'Europe d'après les originaux et les publications les plus récentes. Deuxième partie: Descriptions des espèces. Traduit de l'allemand par Ferd. Rieber *Revue et Magasin de Zoologie*, Ser. 3, **4**: 11–268.
- Germar, E.F.** 1830. Species Cicadarium enumeratae et sub genera distributae. *Thon's Entomologisches Archiv*, **2**(2): 1–57.
- Gnezdilov V.M.** 2009. Revisionary notes on some tropical Issidae and Nogodinidae (Hemiptera: Fulgoroidea). *Acta Entomologica Musei Nationalis Pragae*, **49**(1): 75–92.
- Gnezdilov V.M.** 2013. Modern classification and the distribution of the family Issidae Spinola (Homoptera, Auchenorrhyncha, Fulgoroidea). *Entomologicheskoe obozrenie*, **92**(4): 724–738. (In Russian; English translation: *Entomological Review*, 2014, **94**(5): 687–697).
- Gnezdilov V.M. & O'Brien L.B.** 2006. *Hysteropterum severini* Caldwell & DeLong, 1948, a synonym of *Agalmatium bilobum* (Fieber, 1877) (Hemiptera: Fulgoroidea: Issidae). *The Pan-Pacific Entomologist*, **82**(1): 50–53.
- Gnezdilov V.M. & Sugonyaev E.S.** 2009. First record of *Metcalfa pruinosa* (Homoptera, Fulgoroidea, Flatidae) from Russia. *Zoosystematica Rossica*, **18**(2): 260–261.
- Gnezdilov V.M., Soulier-Perkins A. & Bourgoin T.** 2011. Fieber's original drawings and their corresponding types for the family Issidae (Hemiptera, Fulgoromorpha) in the Muséum national d'Histoire naturelle of Paris, France. *Zootaxa*, 2806: 24–34.
- Kim Y.Y., Kim M.Y., Hong K. & Lee S.H.** 2011. Outbreak of an exotic flatid, *Metcalfa pruinosa* (Say) (Hemiptera: Flatidae), in the capital region of Korea. *Journal of Asia-Pacific Entomology*, **14**(4): 473–478.
- Lucchi A.** 1997. *Metcalfa pruinosa* and honey production in Italy. *American Bee Journal*, **137**(7): 532–535.
- Mazza G., Pennacchio F., Gargani E., Franceschini I., Roversi P.F. & Cianferoni F.** 2014. First report of *Ricania speculum* (Walker, 1851) in Europe (Hemiptera: Fulgoromorpha: Ricaniidae). *Zootaxa*, 3861: 297–300.
- Nielson M.W., Zack R.S., Poggi F. & Nickel H.** 2014. New leafhopper species of *Jikradia* from Mesoamerica with new records, revised key to species, distribution, origin, and checklist (Hemiptera: Cicadellidae: Coelidiinae: Teruliini). *Revista de Biologia Tropical*, **62**: 1375–1383.
- O'Brien L.B.** 1967. *Caliscelis bonellii* (Latreille), a European genus of Issidae new to the United States (Homoptera: Fulgoroidea). *Pan-Pacific Entomologist*, **43**: 130–133.
- Preda C. & Skolka M.** 2011. Range expansion of *Metcalfa pruinosa* (Homoptera: Fulgoroidea) in Southeastern Europe. *Ecologia Balkanica*, **3**(1): 79–87.
- Synave H.** 1976. Fam. Issidae. La faune terrestre de l'île de Sainte-Hélène. Pt. 3. *Annales du Musée royal de l'Afrique centrale. Ser. IN-8. Sciences Zoologiques*, **215**: 260–262.
- Wheeler A.G. Jr. & Wilson S.W.** 1988. Notes on the biology and immatures of the issid planthoppers *Thionia bullata* and *T. simplex* (Homoptera: Fulgoroidea). *Journal of the New York Entomological Society*, **96**(3): 266–273.
- Zangheri S. & Donadini P.** 1980. Comparsa nel Veneto di un Omottero nearctico: *Metcalfa pruinosa* Say (Homoptera: Flatidae). *Redia*, **63**: 301–304.

Received October 8, 2014 / Accepted November 7, 2014

Editorial responsibility: D.A. Gapon