

# *Kakuna taibaiensis* sp. n. and a newly recorded species of *Dicranotropis* (Hemiptera, Fulgoroidea, Delphacidae) from China

Feng-Juan Ren<sup>1</sup>, Qi Xie<sup>2</sup>, Li Qiao<sup>1,3</sup>, Dao-zheng Qin<sup>1</sup>

**1** Key Laboratory of Plant Protection Resources and Pest Management of the Ministry of Education, Entomological Museum, Northwest A&F University, Yangling, Shaanxi 712100, China **2** Baoji University of Art and Science, Baoji, Shaanxi 721013, China **3** Xinyang City Academy of Agricultural Sciences, Xinyang, Henan 464000, China

Corresponding author: Dao-zheng Qin ([qindaozh@nwsuaf.edu.cn](mailto:qindaozh@nwsuaf.edu.cn))

---

Academic editor: Mike Wilson | Received 29 April 2014 | Accepted 1 September 2014 | Published 8 October 2014

---

<http://zoobank.org/B4288100-0C42-4A6E-804F-4C54582D7825>

---

**Citation:** Ren F-J, Xie Q, Qiao L, Qin D-z (2014) *Kakuna taibaiensis* sp. n. and a newly recorded species of *Dicranotropis* (Hemiptera, Fulgoroidea, Delphacidae) from China. ZooKeys 444: 119–130. doi: 10.3897/zookeys.444.7810

---

## Abstract

One new species of the delphacid genus *Kakuna* Matsumura, *K. taibaiensis* Ren & Qin, **sp. n.** is described from Mt. Taibai in Shaanxi Province, China. *Dicranotropis montana* (Horvath, 1897) is reported for the first time from China. Habitus photos and illustrations of male genitalia of the two species are given.

## Keywords

Auchenorrhyncha, planthoppers, Fulgoromorpha, taxonomy, distribution

## Introduction

The genus *Kakuna* was established by Matsumura (1935) for the type species *K. kuwayamai* Matsumura, 1935 from Japan (Hokkaido, Sapporo). Ding (2006) regarded *Parametopina* Yang as a junior synonym of *Kakuna* Matsumura and transferred *Parametopina yushaniae* Yang to *Kakuna*. Recently, Chen and Yang (2010) redefined the generic

characteristics and added three more species, *K. lii*, *K. nonspina* and *K. zhongtuana* to the genus from southwestern China (Guizhou). To date, five *Kakuna* species are known all distributed in China and Japan. In this paper, we add a new species, *K. taibaiensis* from Mt. Taibai (China: Shaanxi) and provide a key to all species in this genus.

Another delphacid species, *Dicranotropis montana*, was described by Horvath (1897), which was originally arranged as a member of the genus *Stiroma* Fieber. Ashe (1982) studied the type material of this species and transferred it into *Dicranotropis* Fieber. This species is currently distributed in the Palaearctic Region. After checking the specimens collected in 2010 (now deposited in the Entomological Museum in NWFU), we found this species in Hebei (northern China) and record it for the first time in the Chinese fauna.

## Material and methods

All specimens examined in this study are deposited in the Entomological Museum, Northwest A & F University, Yangling, Shaanxi, China (NWFU). The genital segments of the examined specimens were macerated in 10% KOH and drawn from preparations in glycerin jelly with the aid of a light microscope. Illustrations of the specimens were made using a Leica MZ 12.5 stereomicroscope. Habitus photos were taken using a Scientific Digital micrography system equipped with an Auto-montage imaging system and a highly sensitive QIMAGING Retiga 4000R digital camera (CCD). Multiple photographs were compiled into final images. The terminology in this paper follows that of Ding (2006). Measurements of the body length were from the apex of the vertex to the posterior tip of the abdomen (macropterous) or to the tip of abdomen (brachypterous). All measurements are in millimeters (mm).

## Taxonomy

### *Kakuna* Matsumura, 1935

*Kakuna* Matsumura 1935: 76; Ding 2006: 404; Chen and Yang 2010: 30. Type species: *Kakuna kuwayamai* Matsumura, 1935, by original designation.

*Parametopina* Yang, 1989: 308. Type species: *Parametopina yushaniae* Yang 1989: 308, synonymized by Ding 2006: 404.

**Diagnosis.** Relatively large, brown delphacids. Head including eyes narrower than pronotum. Submedian carinae uniting at apex of vertex. Fastigium in lateral view rounded. Dorsum of body with a milky longitudinal stripe from middle of vertex to middle of posterior margin of forewing. Median carina of frons forked at extreme base. Antennae cylindrical. Forewing with large, longitudinal, brown marking. Metabasitarsus longer than tarsomere 2 + 3 combined, calcar thin, tectiform, with many black-tipped teeth on lateral margin. Male pygofer in lateral view with laterodorsal angle

produced, medioventral process absent. Diaphragm of pygofer narrow, dorsal margin produced dorsad. Suspensorium ventrally ring-like. Aedeagus tubular, long. Parameres fairly developed, apically convergent. Anal segment deeply sunk into dorsal emargination of pygofer, ring-like, caudoventral processes present or absent.

**Distribution.** China (Guizhou, Taiwan, Zhejiang, Fujian, Shaanxi), Japan.

**Key to species in the genus *Kakuna* (males)**

- 1 Male anal segment produced caudoventrally..... 2
- Male anal segment not produced caudoventrally..... 3
- 2 Male anal segment with two spine-like processes along caudoventral margin; mediodorsal processes of diaphragm separated at bases; aedeagus not bearing spinous processes at apex. China (Guizhou).....  
..... ***K. nonspinata* Chen & Yang**
- Male anal segment with an elongate process at caudoventral margin; mediodorsal processes of diaphragm fused at their bases; aedeagus bearing spinous processes at apex. China (Taiwan) ..... ***K. yushaniae* (Yang)**
- 3 Aedeagus in profile distinctly expanded at apex, dorsal margin with spinous processes ..... 4
- Aedeagus in profile not expanded at apex, dorsal margin without spinous processes ..... 5
- 4 Mediodorsal processes of diaphragm curved laterad apically; aedeagus without spinous processes ventrally near apex. China (Guizhou) .....  
..... ***K. lii* Chen & Yang**
- Mediodorsal processes of diaphragm straight apically; aedeagus with spinous processes ventrally near apex. China (Zhejiang, Fujian, Guizhou), Japan (Hokkaido, Honshu, Kyushu) ..... ***K. kuwayamai* Matsumura**
- 5 Mediodorsal processes of diaphragm long, reaching to the level of anal segment; aedeagus in profile curved ventrad medially; inner margins of parameres with denticles medially. China (Shaanxi) .....  
..... ***K. taibaiensis* Ren & Qin, sp. n.**
- Mediodorsal processes of diaphragm short, not reaching to the level of anal segment; aedeagus in profile curved dorsad medially; inner margins of parameres without denticles but with a nipple-like process medially. China (Guizhou)..... ***K. zhongtuana* Chen & Yang**

***Kakuna taibaiensis* Ren & Qin, sp. n.**

<http://zoobank.org/0DE83AE8-F62C-4649-8C6E-459D960AA940>

Figs 1–16

**Description.** Macropterous male: Body length: male 5.82–5.91 mm; forewing length: male 5.06–5.13 mm (n=2).




**Figures 1–5.** *Kakuna taibaiensis* Ren & Qin, sp. n. **1** male adult, dorsal view **2** male adult, left lateral view **3** head and thorax, dorsal view **4** frons and clypeus **5** metatarsus and post-tibial spur. Scale bars = 1.0 mm (Figs **1**, **2**); 0.5 mm (Figs **3–5**).

**Color.** General color brown. Ocelli reddish brown, eyes black. Dorsum of body with a milky longitudinal stripe from the junction of Y-shaped carina to the middle of posterior margin of forewing. Forewing yellowish brown, membrane has a large, longitudinal, fuscous marking from base of costal area to apex, veins fuscous, longitudinal veins ornamented with blackish brown granules. Abdomen fuscous. Fore and middle legs brown, hind legs yellowish brown, apices of spines on tibiae and tarsi black.

**Structure.** Head including eyes narrower than pronotum (about 0.81:1) (Figs 1, 3). Vertex shorter in midline than wide at base (about 0.82:1), narrower at apex than at base (about 0.78:1), lateral margins slightly concave in dorsal view, submedian carinae convex, originating from near 1/3 base of lateral carinae and uniting at apex of vertex (Figs 1, 3). Y-shaped carina distinct, basal compartment shallowly concave (Fig. 3). Fastigium rounded (Fig. 2). Frons longer in midline than maximum width about 2.05:1, widest above the level of ocelli, median carina conspicuous, forked at extreme base (Fig. 4). Postclypeus wider at base than frons at apex, post- and anteclypeus together approximately 0.86 × of the length of frons (Fig. 4). Rostrum almost reaching mesotrochanters. Antennae terete, nearly attaining middle level of postclypeus, scape longer than wide at apex (about 1.83:1), shorter than pedicel (about 0.52:1) (Fig. 4).

Pronotum in midline slightly shorter than length of vertex (about 0.79:1), lateral carinae developed, slightly curved, not reaching posterior margin (Fig. 3). Mesonotum medially ca. 1.64 times longer than vertex and pronotum together, lateral carina almost straight, reaching posterior margin, median carina obscure apically (Fig. 3). Forewings long and narrow, longer than widest part about 3.35:1, widest in middle (Figs 1, 2, 16). Spination of apex of hind leg 5 (3+2) (tibia), 8 (6+2) (basitarsus) and 4 (2nd



**Figures 6–16.** *Kakuna taibaiensis* Ren & Qin, sp. n. **6** male genitalia, posterior view **7** male genitalia, left lateral view **8** male pygofer, posterior view **9** male pygofer, left lateral view **10** anal segment, posterior view **11** anal segment, aedeagal complex, connective and parameres, left lateral view **12** aedeagus, dorsocaudal view **13** aedeagus, left lateral view **14** suspensorium, posterior view **15** parameres, posterior view **16** forewing. Scale bars = 0.2 mm (Figs 6–9, 11, 15); 0.1 mm (Figs 10, 12–14); 0.5 mm (Fig. 16).

tarsomere) (Fig. 5). Metabasisarsus distinctly longer than tarsomere 2+3 combined (about 1.79:1), calcar shorter than metabasisarsus (about 0.77:1), thin, bearing 29 black-tipped teeth on lateral margin (Fig. 5).

**Male genitalia.** Male pygofer slightly wider ventrally than dorsally, laterodorsal angles roundly produced caudad; in posterior view with opening longer than wide, ventral margin shallowly excavated, medioventral process absent (Figs 6–9). Suspensorium ventrally ring-like, dorsally with a process at each side leading to the anal segment ventrolaterally (Fig. 14). Diaphragm narrow, mediodorsal processes fairly developed, pillar-like, basally with a broad common stalk, thence contiguous apicad, apical part separated and curved laterad, tips truncated (Figs 6, 8). Parameres fairly long, reaching to the level of anal segment, in posterior view contiguous basally, apical 2/5 convergent mesad, apices rounded, inner margins expanded and ornamented with denticles medially (Figs 6, 11, 15). Aedeagus tubular, arch-shaped in profile, moderately dilated near the base, near apex on the dorsal side to the ventral apex provided with small teeth, gonopore apical on the slightly membranous dorsal side (Figs 12, 13). Male anal segment deeply sunk into dorsal emargination of pygofer, ring-like, caudoventral processes absent (Figs 6, 7, 10, 11).

**Type materials. Holotype.** ♂ (macropterous, NWAUFU), China, Shaanxi Province, Mt. Taibai, 13-VIII-2010, by light trap, coll. A. P. Dong. **Paratype.** 1♂ (macropterous, NWAUFU), same data as holotype.

**Female.** Unknown.

**Host plant.** Unknown.

**Etymology.** The species epithet is named after the type locality, Mt. Taibai in Shaanxi, China.

**Distribution.** Known currently from the type locality in northwest China (Shaanxi Province).

**Remarks.** *Kakuna taibaiensis* is similar to *K. zhongtuana* Chen & Yang (2010) in the male anal segment not produced caudoventrally, aedeagus not bearing spinous processes and mediodorsal processes of diaphragm having a common stalk basally. However, the new species differs from the latter in the mediodorsal processes fairly long, reaching to the level of anal segment (mediodorsal processes short, not reaching to the level of anal segment in *zhongtuana*), aedeagus curved ventrad medially in profile (aedeagus curved dorsad medially in profile in *zhongtuana*), parameres rounded at apex in posterior view, inner margins ornamented with denticles medially (parameres acute at apex and adorned with a nipple-like process medially along each inner margin in *zhongtuana*).

**Discussion.** The Himalaya-Qinling-Huai River line is the most distinctive barrier and may serve as the division of the Palaearctic and Oriental Regions since the Pleistocene. However, the north-south transitional affects have been much more pronounced in species and a broad transitional zone has resulted (Zhang 2002). The new finding in this paper based on the specimens from Mt. Taibai (the main peak of Mts. Qinling in Shaanxi, China) confirms the suggestion of Chen and Yang (2010) that

the members of the genus *Kakuna* have extended into the southern area of the Palearctic Region. During our investigations of Delphacidae on Mt. Taibai, we found many species in this family have extended into the border of the two Regions which were traditionally thought to be confined in the Palearctic or Oriental Region only, including some new species described in recent years (Qin 2007, Qin et al. 2012). We suspect that the delphacid fauna in this border area will be more extensive if more investigations are conducted.

***Dicranotropis montana* (Horvath, 1897)**

new record to China

Figs 17–34

*Stiroma montana* Horvath 1897: 625.

*Dicranotropis tenellula* Dlabola 1965: 84; Emeljanov 1977: 109, synonymized by Asche 1982: 30.

*Dicranotropis gratiosa* Dlabola 1997: 315, synonymized by Holzinger 1999: 259.


*Dicranotropis montanus* Vilbaste 1965: 15, synonymized by Asche 1982: 30.

*Dicranotropis montana* (Horvath), Asche 1982: 30; Holzinger 1999: 259; Holzinger et al. 2003: 267; Nickel 2003: 55; Trivellone 2010: 100.

**Description.** Macropterous male: Body length (from apex of vertex to the tip of forewing): male 3.40–3.64 mm, female 3.44–3.90 mm; forewing length: male 2.72–2.96 mm, female 3.04–3.24 mm. Brachypterous male: Body length (from apex of vertex to the tip of abdomen): male 2.24–2.56 mm, female 2.64–2.96 mm; forewing length: male 0.99–1.08 mm, female 1.04–1.24 mm.

**Color.** General color of male (macropterous) blackish brown. Ocelli reddish black, eyes grayish black. Vertex anteriorly, frons, clypeus, lateral area of pronotum behind eyes black. Antennae yellowish brown except apex of scape and base of pedicel fuscous. Pronotum between lateral carinae and laterobasal angles sordid whitish. All carinae and margins of vertex, frons and clypeus whitish. Rostrum fuscous at apex. Mesonotum mostly dark brown, scutellum whitish apically. Abdomen dark. Legs brown to yellowish brown. Tegmina subhyaline, veins yellowish brown. Female with ovipositor brown. Male (brachypterous) with the same color as macropterous except pronotum, mesonotum and tegmina yellowish brown, abdomen of female mostly yellowish white, abdomen with small brown spots dorsally and ventrally on each segment.

**Structure.** Head including eyes slightly narrower than pronotum (about 0.92:1). Vertex shorter in midline than wide at base (about 0.62:1), narrower at apex than at base (about 0.89:1). Submedian carinae originating from near 1/4 base of lateral carinae, not uniting at apex of vertex. Y-shaped carina distinct (Figs 17, 19, 21). Frons longer in midline than maximum width about 1.64:1, widest above the level of ocelli, carinae conspicuous, median carina forked at basal fourth (Fig. 22). Postclypeus broader




**Figures 17–22.** *Dicranotropis montana* (Horvath, 1897) **17** macropterous male, dorsal view **18** macropterous male, left lateral view **19** brachypterous male, dorsal view **20** brachypterous male, left lateral view **21** head and thorax (macropterous male), dorsal view **22** frons and clypeus (brachypterous male). Scale bars = 1.0 mm (Figs 17, 18); 0.5 mm (Figs 19–22).

at base than frons at apex, postclypeus and anteclypeus together approximately  $0.80 \times$  the length of the frons (Fig. 22). Rostrum almost reaching mesotrochanters. Antennae terete, reaching frontoclypeal suture, scape longer than apical width (about 1.59:1), shorter than pedicel (about 0.64:1) (Figs 18, 20, 22).

Pronotum shorter than vertex in midline (about 0.91:1), lateral carinae straight, not reaching to posterior margin (Figs 17, 19, 21). Mesonotum medially ca. 1.34





**Figures 23–34.** *Dicranotropis montana* (Horvath, 1897). **23** male genitalia, posterior view **24** male genitalia, left lateral view **25** male pygofer, posterior view **26** anal segment, posterior view **27** anal segment, aedeagal complex, connective and parameres, left lateral view **28** aedeagus, left lateral view **29** aedeagus, right lateral view **30** aedeagus, dorsal view **31** suspensorium, posterior view **32** parameres, posterior view **33** macropterous forewing **34** brachypterous forewing. Scale bars = 0.2 mm (Figs 23–25, 27); 0.1 mm (Figs 26, 28–30, 32); 0.04 mm (Fig. 31); 0.5 mm (Fig. 33); 0.25 mm (Fig. 34).

times longer than vertex and pronotum together, lateral carina reaching posterior margin, median carina obscure apically (Figs 17, 21). Macropterous forewings surpassing tip of abdomen approximately 2/5 of its total length (Figs 17, 18), longer than widest part (about 2.86:1). Spinal formula of hind leg 5–7–4, post-tibial spur shorter than metabasitarsus, sparsely with about 8 tiny teeth along hind margin.

**Male genitalia.** Male pygofer in profile wider ventrally than dorsally, anterior margin distinctly convex submedially (Fig. 24); in posterior view opening of pygofer obcordate, medioventral process absent (Figs 23, 25), below laterodorsal angle interiorly with a spine-like process on each side, transverse-directed (Figs 23, 25). Suspensorium n-shaped, dorsally arched medially with two small triangular processes on both ends (Fig. 31). Diaphragm broad, mediadorsal processes strongly sclerotized and laterally beset with many granulations, incised medially (Figs 23, 25). Opening for parameres large, dorsal margin nearly straight, lateral margins slightly sinuate (Fig. 25). Parameres long, contiguous at bases, narrowed and divergent apically, inner margins expanded subapically, in profile apical margin emarginated in two triangular processes (Fig. 27). Aedeagus tubular, short and broad, with five rows of teeth on surface, including four longitudinal rows and one transverse row basad of gonopore (Figs 28, 29, 30). Male anal segment ring-like, laterodistal angles produced into a short, stout spinose process on each side (Figs 23, 24, 26, 27).

**Species examined.** 23 ♂♂ 22 ♀♀ (macropterous) and 35 ♂♂ 46 ♀♀ (brachypterous), China: Hebei Province, Mt. Xiaowutai, 24-VI-2009, coll. D. Z. Qin.

**Distribution.** China (Hebei), Russia, Austria, Switzerland, Germany, France, Italy, Hungary, Romania, Mongolia.

**Host plant.** Unknown.

**Discussion.** Dlabola (1965) established *D. tenellula* Dlabola based on the specimens from Mongolia; Vilbaste (1965) described *D. montana* Vilbaste from Russia but it was regarded as a junior synonymy of *D. tenellula* Dlabola by Emeljanov (1977). Asche (1982) treated both *D. tenellula* Dlabola and *D. montana* Vilbaste as junior synonyms of *D. montana* (Horvath, 1897). However, the treatment of *D. tenellula* Dlabola was not accepted by Anufriev and Emeljanov (1988). Holzinger et al. (2003) studied the *Dicranotropis* species in central Europe, in *D. montana* (Horvath) part, they redrew the male genitalia of this species and noted: “according to Emeljanov and Gnezdilov (pers. common.), the central Asian *Dicranotropis tenellula* Dlabola, 1965 is not conspecific with *D. montana* (Horvath, 1897)”. After checking the specimens deposited in NWFU, and also these illustrations of male genitalia provided by Dlabola (1965, 1997), Vilbaste (1965), Anufriev and Emeljanov (1988), Holzinger (1999) and Holzinger et al. (2003), we found *D. tenellula* Dlabola to be very similar to *D. montana* (Horvath) and very difficult to distinguish. We hope the status of *D. tenellula* can be reconsidered and firmly established using more advanced methods in the future. Here we consider *D. tenellula* Dlabola as a junior synonym of *D. montana* (Horvath).

## Acknowledgements

We are grateful to Prof. Richard John Schrock (Emporia State University, Emporia, KS, USA) for reviewing the manuscript. This work was supported by the National Natural Science Foundation of China (Nos. 30970387 and 31172126).

## References

- Anufriev GA, Emeljanov AF (1988) Suborder Cicadinea (Auchenorrhyncha)-Cicadas. In: Ler PA (Ed.) Keys to the identification of insects of the Soviet Far East. Vol. 2: Homoptera and Heteroptera. Nauka, Leningrad, 12–495.
- Asche M (1982) Zur Kenntnis einiger von Geza Horvath beschriebener Delphaciden (Homoptera Cicadina Fulgoromorpha). Marburger Entomologische Publikationen 1(7): 25–36.
- Chen XS, Yang L (2010) Oriental bamboo delphacid planthoppers: three new species of genus *Kakuna* Matsumura (Hemiptera: Fulgoromorpha: Delphacidae) from Guizhou Province, China. *Zootaxa* 2344: 29–38.
- Ding JH (2006) Fauna Sinica. Insecta Vol. 45. Homoptera Delphacidae. Editorial Committee of Fauna Sinica, Chinese Academy of Science. Science Press, Beijing, China, 776 pp.
- Dlabola J (1965) Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. 54; Homoptera, Auchenorrhyncha. *Acta Faunistica Entomologica Musei Nationalis Pragae* 11(100): 79–136.
- Dlabola J (1997) *Mycterodus* verwandte Taxone und sieben neue Zikadenarten (Homoptera, Auchenorrhyncha). *Acta Entomologica Musei Nationalis Pragae* 44 (1995): 301–319.
- Emeljanov AF (1977) Leaf-hoppers (Homoptera, Auchenorrhyncha) from the Mongolian People's Republic based mainly on materials of the Soviet-Mongolian zoological expeditions (1967–1969). *Nasekomye Mongolii* 5: 96–195.
- Holzinger WE (1999) Taxonomie und Verbreitung ausgewählter Zikadenarten Österreichs (Insecta: Hemiptera: Auchenorrhyncha). *Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden* 21(17): 259–264.
- Holzinger WE, Kammerlander I, Nickel H (2003) The Auchenorrhyncha of Central Europe—Die Zikaden Mitteleuropas Fulgoromorpha, Cicadomorpha excl. Cicadellidae. Brill Publishers, Leiden, the Netherlands, 673 pp.
- Horvath G (1897) Homoptera nova ex Hungaria. *Termeszetráji Füzetek*, Budapest 20: 620–643.
- Matsumura S (1935) Supplementary note to the revision of *Stenocranus* and its allied species in Japan-Empire. *Insecta Matsumurana* 10: 71–78.
- Nickel H (2003) The Leafhoppers and Planthoppers of Germany (Hemiptera, Auchenorrhyncha): patterns and strategies in a highly diverse group of phytophagous insects. *Pensoft Series Faunistica* No. 28, Sofia-Moscow, Keltern, 1–460.
- Qin DZ (2007) Two new species of the Chinese endemic delphacid genus *Neuterthron* Ding (Hemiptera: Fulgoromorpha) from Yunnan and Shaanxi Provinces. *Zootaxa* 1547: 59–64.

- Qin DZ, Liu TT, Lin YF (2012) A new species in the *Bambusiphaga fascia* group (Hemiptera, Fulgoroidea, Delphacidae) from China, with a key to all species in the group. *Acta Zootaxonomica Sinica* 37(4): 777–780.
- Trivellone V (2010) Contribution to the knowledge of the Auchenorrhyncha fauna of bogs and fens of Ticino and Grisons, with some new records for Switzerland (Hemiptera: Fulgoro-morpha et Cicadomorpha). *Cicadina* 11: 97–106.
- Vilbaste J (1965) Uber die Zikadenfauna des Altai (in Russian). *Akademia Nauk. Tartu* 1–144.
- Yang CT (1989) Delphacidae of Taiwan (II) (Homoptera: Fulgoroidea). *National Science Council Special Publication* 6: 334 pp.
- Zhang RZ (2002) Geological events and mammalian distribution in China. *Acta Zoologica Sinica* 48(2): 141–153.