

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/313657892>

New distributional records of Delphacidae (Hemiptera: Fulgoroidea) from Chile

Article in *Archivos Entomológicos* · February 2017

CITATIONS

0

READS

243

1 author:



Juan Francisco Campodonico

University of Chile

8 PUBLICATIONS 1 CITATION

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Contributions to the knowledge of Fulgoroidea (Hemiptera) fauna in Chile [View project](#)



Contributions to the knowledge of Membracoidea (Hemiptera) fauna in Chile [View project](#)

All content following this page was uploaded by [Juan Francisco Campodonico](#) on 13 February 2017.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.

ARTIGO / ARTÍCULO / ARTICLE

New distributional records of Delphacidae (Hemiptera: Fulgoroidea) from Chile.

Juan F. Campodonico

J.M. Perceval 10259, Vitacura, Santiago, CHILE. e-mail: juanfranciscocampodonico@gmail.com

Abstract: *Chionomus haywardi* (Muir, 1929) (previously known for Argentina and Uruguay), *Syndelphax dissipatus* (Muir, 1926) (previously known for Ecuador, Brazil, Panamá and Costa Rica) and *Dicranotropis bipectinata* Muir, 1926 (previously known for Ecuador) are recorded for the first time from Chile. Additionally, the known distribution range of *Delphacodes darwini* Muir, 1929 (previously known only from Los Lagos region) is expanded up to Coquimbo region and *Delphacodes kuscheli* Fennah, 1955 (previously known from Juan Fernández Islands, Argentina and Uruguay) is recorded for first time from continental Chile. Photographs and diagnostic characters of the males are provided.

Key words: Hemiptera, Fulgoroidea, Delphacidae, Delphacini, *Chionomus*, *Syndelphax*, *Dicranotropis*, *Delphacodes*, *Delphacodes kuscheli*, Faunistics, Chile.

Resumen: Nuevos registros de distribución de Delphacidae (Hemiptera: Fulgoroidea) de Chile. Se registran por primera vez para Chile las especies *Chionomus haywardi* (Muir, 1929) (previamente conocida de Argentina y Uruguay), *Syndelphax dissipatus* (Muir, 1926) (previamente conocida de Ecuador, Brasil, Panamá y Costa Rica) y *Dicranotropis bipectinata* Muir, 1926 (previamente conocida de Ecuador). Adicionalmente, el rango de distribución conocido de *Delphacodes darwini* Muir, 1929 (previamente conocida sólo de la región de Los Lagos) se amplía hasta la región de Coquimbo y *Delphacodes kuscheli* Fennah, 1955 (previamente conocida del archipiélago de Juan Fernández, Argentina y Uruguay) se registra por primera vez para Chile continental. Se proporcionan fotografías y caracteres diagnósticos de los machos.

Palabras clave: Hemiptera, Fulgoroidea, Delphacidae, Delphacini, *Chionomus*, *Syndelphax*, *Dicranotropis*, *Delphacodes*, *Delphacodes kuscheli*, Faunística, Chile.

Recibido: 18 de enero de 2017

Publicado on-line: 13 de febrero de 2017

Aceptado: 2 de febrero de 2017

Introduction

Delphacidae (Hemiptera: Fulgoroidea), including 2152 described species and 411 genera (Bourgoin 2017), is the most economically important family within the Fulgoroidea (Urban *et al.* 2010). It is characterized by the presence of a movable spur (the calcar) at the apex of the hind tibia (Bartlett 2014). Up to this date, 25 species classified in 11 genera have been recorded from Chile (Spinola 1852, Bergroth 1924, Muir 1927, 1929, 1934, Fennah 1955, 1957, 1965, 1969, Aguilera 1972, Remes-Lenicov 1996, Rioja *et al.* 2006, 2010, Remes-Lenicov & Rioja 2007, Gonzon & Bartlett 2007, Rossi-Batiz 2014, Campodonico 2015). Among these, 10 species have been recorded in Chile only from the Juan Fernández Islands, and one is shared between insular and continental Chilean territories (Fennah 1955, 1957, Campodonico 2015).

Materials and methods

For identification, Muir (1926, 1929), Fennah (1955, 1957), Remes-Lenicov & Tesón (1978) and

Weglarz (2012) were followed. For observation of the male genitalia, the abdomen was removed and placed in a saturated KOH solution at room temperature for about 24 hours. When cleared, it was neutralized. Posterior to dissection and examination in glycerine under both stereoscopic and optical microscope, pieces were stored in a micro vial pinned below the respective specimen. Photographs of specimens (Figs. 1-41) were taken with a conventional digital camera adapted to stereoscopic and optical microscopes. The maps (Figs. 42-44) were developed with ArcGIS®.

The material examined was deposited on the following collections: Museo Entomológico, Facultad de Ciencias Agronómicas, Universidad de Chile, Santiago, Chile (MEUC); Museo Nacional de Historia Natural, Santiago, Chile (MNNC); Juan F. Campodonico Particular Collection, Santiago, Chile (JFCW); University of Delaware Insect Research Collection, Newark, DE, USA (UDCC).

Results

Chionomus haywardi (Muir, 1929) (Figs. 1-8, 43)

Delphacodes haywardi Muir, 1929 - Muir (1929:83); Remes-Lenicov & Virla (1996:165); Tesón & Remes-Lenicov (1983:322); Remes-Lenicov & Virla (1999:11); Remes-Lenicov *et al.* (2000:93); Velázquez *et al.* (2003:669).

Chionomus haywardi (Muir, 1929) - Fennah (1971:324) (Combination); Weglarz (2012:44).

Delphacodes collaris Remes-Lenicov & Tesón, 1978 - Remes-Lenicov & Tesón (1978:17); Tesón & Remes-Lenicov (1983:322) (Synonymy).

Material examined: CHILE: VALPARAÍSO REGION: **Marga-Marga province:** Quilpué, Fundo El Carmen, 33°02'S, 71°27'W, 90-120 m, 14.III.2015, J.F. Campodonico leg., 2♂♂ (macropterous), 1♀ (macropterous) (JFCW); Las Cruces, 33°29'S, 71°38'W, 10 m, 18.VII.2016, J.F. Campodonico leg., 1♂ (macropterous) (JFCW); METROPOLITANA REGION: **Santiago province:** Vitacura, Plaza Los Tribunales, 33°22'S, 70°32'W, 765 m, 19.V.2015, J.F. Campodonico leg., sweeping, 4♂♂ (macropterous), 1♀ (macropterous) (JFCW), 2♂♂ (macropterous), 1♀ (macropterous) (MNNC), 2♂♂ (macropterous), 1♀ (macropterous) (UDCC); Maipú, Quebrada de la Plata, 33°29'S, 70°54'W, 650 m, 25.V.2016, J.F. Campodonico leg., sweeping, 1♂ (macropterous) (JFCW), 2♂♂ (macropterous), 1♀ (macropterous) (MEUC); MAULE REGION: **Talca province:** Putú, 35°40'S, 72°11'W, 11 m, 16.VII.2016, J.F. Campodonico leg., sweeping, 3♂♂ (macropterous) (JFCW).

Male genitalia: Pygofer (Figs. 4-5) with opening as an inverted triangle; armature of diaphragm distinctly projecting caudad; anal tube (Fig. 6) with two converging spines directed ventrad; parameres (Fig. 7) wide and short, narrower at the middle, inner angle acute and outer angles rounded; aedeagus (Fig. 8) tubular, wider near base, slightly curved dorsad (Muir 1929, Remes-Lenicov & Tesón 1978, Weglarz 2012).

Previously recorded from Argentina and Uruguay (Muir 1929, Remes-Lenicov *et al.* 2000). Description of female in Remes-Lenicov & Tesón (1978) and Tesón & Remes-Lenicov (1983). Immature stages described by Remes-Lenicov & Virla (1996). *Chionomus haywardi* has been syndicated as a vector of Mal de Río Cuarto maize virus in Argentina (Velázquez *et al.* 2003).

Syndelphax dissipatus (Muir, 1926) (Figs. 9-16, 42)

Delphacodes dissipata Muir, 1929 - Muir (1926:33).

Syndelphax dissipatus (Muir, 1929) - Fennah (1967:76) (Combination); Bartlett & Kunz (2015:600).

Material examined: CHILE: ARICA AND PARINACOTA REGION: **Arica province:** Km. 23, Lluta, 12.VI.1968, N. Hichins leg., 1♀ (macropterous) (MEUC); Km. 23, Lluta, 13.VI.1968, N. Hichins leg., 1♂ (macropterous) (MEUC); ATACAMA REGION: **Huasco province:** Freirina, 10.X.2015, J.F. Campodonico leg., sweeping, 2♂♂ (brachypterous), 1♂ (macropterous), 2♀♀ (brachypterous), 2♀♀ (macropterous) (JFCW), 1♂ (brachypterous), 1♀ (brachypterous) (MNNC), 1♂ (brachypterous) (MEUC), 2♂♂ (brachypterous) (UDCC).

Male genitalia: Opening of pygofer (Fig. 12) a little wider than long; armature of diaphragm (Fig. 13) as a large, subconical process shagreened along the sides; anal tube (Fig. 14) with two large flat spines arising from the middle of the ventral side and the ventro-apical margin curved ventrad; parameres (Fig. 15) broad at basal half, at apical half narrow with outer and inner margins concave, apex truncate, slightly sinuous, angles slightly produced; aedeagus (Fig. 16) subtubular, slightly wider at base, orifice in a dorso-apical position (Muir 1926).

Previously recorded for Ecuador, Brazil, Panamá and Costa Rica (Muir 1926, Fennah 1967, Bartlett & Kunz 2015). Specimens from Freirina were collected by sweeping on wetland vegetation.

***Dicranotropis bipectinata* Muir, 1926** (Figs 17-24, 42)

Dicranotropis bipectinata Muir, 1926 - Muir (1926:25).

Material examined: CHILE: TARAPACÁ REGION: Tamarugal province: Mamiña, 20°04'S, 69°12'W, 2750 m, 29.I.2016, J.F. Campodonico leg., sweeping, 1♂ (brachypterous) (JFCW).

Male genitalia: Pygofer (Figs. 20-21) with latero-ventral and medio-ventral processes; anal tube (Fig. 22) with two large, curved spines on the ventral side; parameres (Figs. 23-24) furcate, main projection slightly sinuous, gradually tapering to apex, second projection arises about middle of the antero-lateral surface of the main projection and curved upwards with apex curved laterad; aedeagus (Fig. 22) with two conspicuous comb-like processes, in lateral view curved, wide at base, dorsal margin convex, ventral margin concave, orifice long (Muir 1926).

Previously recorded for Ecuador (Muir 1926). Specimen from Mamiña collected by sweeping riparian vegetation.

***Delphacodes darwini* Muir, 1929** (Figs. 25-32, 44)

Delphacodes darwini Muir, 1929 - Muir (1929:78); Bourgoin (2017).

Material examined: CHILE: COQUIMBO REGION: Elqui province: Vicuña, 4.X.1966, R. Charlin leg., 1♂ (macropterous) (MEUC); VALPARAÍSO REGION: Los Andes province: Los Andes, 6.XI.1979, G. Barría leg., 2♂♂ (macropterous) (MEUC); Valparaíso province: Humedal de Mantagua, 32°52'S, 71°30'W, 11 m, 28.XII.2014, J.F. Campodonico leg., sweeping, 1♂ (brachypterous) (JFCW); METROPOLITANA REGION: Santiago province: Vitacura, Plaza Los Tribunales, 33°22'S, 70°32'W, 765 m, 19.V.2015, J.F. Campodonico leg., sweeping, 1♂ (macropterous) (JFCW); Cerro Renca, 26.XII.1983, G. Arriagada leg., 1♂ (macropterous) (MNNC); Antumapu, 6.XI.1978, G. Barría leg., 3♂♂ (macropterous), 6♀♀ (brachypterous) (MEUC); Antumapu, 19.XII.1978, G. Barría leg., 3♂♂ (macropterous), 8♀♀ (brachypterous) (MEUC); Antumapu, 25.VI.1979, G. Barría leg., 1♂ (brachypterous) (MEUC); Antumapu, 4.IX.1979, G. Barría leg., 1♂ (brachypterous) (MEUC); Cordillera province: Pirque, 13.VII.1979, G. Barría leg., 2♂♂ (brachypterous) (MEUC); MAULE REGION: Talca province: Putú, 35°40'S, 72°11'W, 11 m, 16.VII.2016, J.F. Campodonico leg., sweeping, 3♂♂ (brachypterous) (JFCW); LOS RÍOS REGION: Valdivia province: Panguipulli, Puyumen, 39°40'S, 72°17'W, 150 m, 19.IX.2015, J.F. Campodonico leg., sweeping, 2♂♂ (brachypterous), 2♀♀ (brachypterous) (JFCW); Chaihuín, 39°57'S, 73°34'W, 5 m, 11.II.2015, J.F. Campodonico leg., sweeping, 1♂ (brachypterous) (JFCW); LOS LAGOS REGION: Chiloé province: Caulín, I.2015, J.M. Contreras leg., sweeping, 2♂♂ (brachypterous), 4♀♀ (brachypterous) (JFCW).

Male genitalia: Pygofer (Figs. 28-29) with a large anal emargination, anal angles widely produced, rounded and curved; armature of diaphragm as a small plate, truncated at apex; anal tube (Fig. 30) with the medio-ventral margin of apex produced ventrad and with two broad, spines arising from the ventral surface and curved basad; parameres (Fig. 31) long, flattened, widest on the apical half, subtruncate at apex; aedeagus (Fig. 32) with the orifice on the upper surface and some minute teeth just basad of the orifice (Muir 1929).

Previously only recorded for Chiloé, Los Lagos Region (Muir 1929). Photograph of the type in Bourgoin (2017).

***Delphacodes kuscheli* Fennah, 1955 (Figs 33-41, 43)**

Delphacodes kuscheli Fennah, 1955 - Fennah (1955: 137); Fennah (1957:383); Remes-Lenicov & Tesón (1978:22); Tesón & Remes-Lenicov (1983:320); Remes-Lenicov *et al.* (1985:251); Grilli & Gorla (1997:45); Remes-Lenicov & Virla (1999:9); Remes-Lenicov *et al.* (2000:93); Truol *et al.* (2001:39); Grilli & Gorla (2002:187); Remes-Lenicov *et al.* (2008:25).

Material examined: CHILE: VALPARAÍSO REGION: Valparaíso province: Humedal de Mantagua, 32°52'S, 71°30'W, 11 m, 28.XII.2014, J.F. Campodonico leg., sweeping, 5♂♂ (brachypterous), 1♂ (macropterous), 4♀♀ (brachypterous), 1♀ (macropterous) (JFCW), 3♂♂ (brachypterous) (MNNC), 3♂♂ (brachypterous) (MEUC), 3♂♂ (brachypterous) (UDCC).

Male genitalia: Pygofer (Figs. 36-37) in posterior view almost circular; dorsal margin of diaphragm with two triangular lobes; anal tube (Figs. 38-39) with each latero-apical angles produced ventrad in a slender spine; parameres (Fig. 40) broad, "S" shaped with the latero-apical angle strongly produced; aedeagus (Fig. 41) abruptly bent near base, distal portion tubular, porrect, with a ridge thickening traversing each side almost through the whole length (Fennah 1955, 1957).

Previously recorded for Robinson Crusoe Island, Juan Fernández (Chile), Argentina and Uruguay (Fennah 1955, Remes-Lenicov & Tesón 1978, Remes-Lenicov *et al.* 2000). This record fills the gap between Juan Fernández Islands and the oriental side of the southern cone. Description of the female in Remes-Lenicov & Tesón (1978) and Tesón & Remes-Lenicov (1983). Immature stages described by Remes-Lenicov *et al.* (2008). This species has being extensively studied for being the main vector of Mal de Río Cuarto maize virus in Argentina (e.g. Remes-Lenicov *et al.* 1985, Grilli & Gorla 1997, 2002, Truol *et al.* 2001).

Discussion

Syndelphax dissipatus and *Dicranotropis bipectinata* are tropical elements that reach their southern distribution limits in wet habitats between the arid areas of northern Chile. On the other side, *Chionomus haywardi*, *Delphacodes darwini* and *Delphacodes kuscheli* are elements distributed on the southern cone of America, to which is added the colonization of Juan Fernández Islands by the last species. With these records the number of species and genera recorded from Chile ascend up to 28 and 12 respectively and the number of Chilean species shared between the continent and Juan Fernández Islands rises up to two. It is worthy to mention that the South American species assigned to *Delphacodes* Fieber, 1866 and *Dicranotropis* Fieber, 1866 do not actually belong to these genera and their classification need to be revised.

Two species with agricultural importance in Argentina are recorded from the neighbouring country. However, unlike Argentina, the studies related with the phytosanitary role of Delphacidae are scarce in Chile and limited to the far north (Rioja *et al.* 2006, 2010). Faunistic and taxonomical investigations on Chilean Delphacidae should be an input to facilitate research for future studies on their phytosanitary importance.

Acknowledgements

I thank Eduardo I. Faúndez for his help on the redaction of a previous version of the manuscript; Aira Faúndez and Rodrigo Flores for their help on the elaboration of the maps and Danilo Cepeda (MEUC) and Mario Elgueta (MNNC) for allowing me to study the material deposited in their respective collections.

References

- Aguilera, A. 1972. Colección entomológica del Centro de Investigación y Capacitación Agrícola (CICA), Arica. Nuevas determinaciones. II contribución. *Idesia* **2**: 99-116.
- Bartlett, C.R. 2014. *Delphacid planthoppers of North America*. University of Delaware, College of Agriculture & Natural Resources, Newark, Delaware, USA. Accessed at: <http://ag.udel.edu/enwc/research/delphacid/index.html> [4 November 2016].
- Bartlett, C.R. & Kunz, G. 2015. Erratum. A new genus and species of delphacid planthopper (Hemiptera: Fulgoroidea: Delphacidae) from Central America with a preliminary regional species list. *Zootaxa* **3963**(4): 598-600.
- Bergroth, E. 1924. *Hemiptera from Juan Fernandez and Easter Island*. In: Skottsberg, C. (ed.). *The Natural History of Juan Fernandez and Easter Island*. Vol. III Zoology: 395-399. Almqvist & Wiksells Boktryckeri, Uppsala, Sweden.
- Bourgoin, T. 2017. *FLOW (Fulgoromorpha Lists on The Web): a world knowledge base dedicated to Fulgoromorpha*. Version 8. Accessed at: <http://hemiptera-databases.org/flow> [18 January 2017].
- Campononico, J.F. 2015. Sobre la distribución de *Dicranotropis acheron* Fennah, 1957 (Hemiptera: Fulgoroidea: Delphacidae) en Chile. *Archivos Entomológicos* **14**: 265-268.
- Fennah, R.G. 1955. Delphacidae from Juan Fernandez (Homoptera: Fulgoroidea). *Proceedings of the Royal Entomological Society of London (B)* **24**(7-8): 129-138.
- Fennah, R.G. 1957. Los insectos de las Islas Juan Fernández 29. Fulgoroidea. *Revista Chilena de Entomología* **5**: 375-384.
- Fennah, R.G. 1965. Fulgoroidea from Southern Chile (Hemiptera). *Bulletin of the British Museum (Natural History), Entomology* **17**: 233-272.
- Fennah, R.G. 1967. Fulgoroidea from the Galápagos Archipelago. *Proceedings of the California Academy of Sciences* **35**(4): 53-102.
- Fennah, R.G. 1969. A revision of *Idiosystatus* Berg (Homoptera: Fulgoroidea, Delphacidae). *Proceedings of the Royal Entomological Society of London* **38**(3-4): 47-52.
- Fennah, R.G. 1971. Fulgoroidea from the Cayman Islands and adjacent areas. *Journal of Natural History* **5**: 299-342.
- Gonzon, A.T. & Bartlett, C.R. 2007. Systematics of *Hadropygos* n.g., *Metadelphax* Wagner and New World *Toya* Distant (Hemiptera: Delphacidae). *Transactions of the American Entomological Society* **133**(3): 205-277.
- Grilli, M.P. & Gorla, D.E. 1997. The spatio-temporal pattern of *Delphacodes kuscheli* (Homoptera: Delphacidae) abundance in central Argentina. *Bulletin of Entomological Research* **87**(1): 45-53.
- Grilli, M.P. & Gorla, D.E. 2002. Variación geográfica de la abundancia poblacional de *Delphacodes kuscheli* (Fennah) en la región central de Argentina. *Ecología austral* **12**(2): 187-195.

Muir, F. 1926. Contribution to our knowledge of South American Fulgoroidea (Homoptera) Part I. The Family Delphacidae. *Bulletin of the Hawaiian Sugar Planters Association (Entomological Series)* **18**: 1-51.

Muir, F. 1927. A new species of *Sparnia* Stål, from South Chile (Delphacidae, Homoptera). *Annals and Magazine of Natural History (Ser. 9)* **20**: 296-297.

Muir, F. 1929. New and little-known South-American Delphacidae (Homoptera, Fulgoroidea) in the collection of the British Museum. *Journal of Natural History* **3**(13): 75-85.

Muir, F. 1934. New and little-known Fulgoroidea (Homoptera). *Journal of Natural History* **14**(84): 561-586.

Remes-Lenicov, A.M.M. 1996. El género *Dicranotropis* Fieber, 1866, en la República Argentina y Chile (Insecta: Homoptera: Delphacidae). *Acta Entomológica Chilena* **20**: 123-128.

Remes-Lenicov, A.M.M.; Brentassi, M.E. & Toledo, A.V. 2008. Description of the immature stages of *Delphacodes kuscheli* Fennah (Hemiptera: Delphacidae), vector of "Mal de Río Cuarto virus" on maize in Argentina. *Studies on Neotropical Fauna and Environment* **43**(1): 25-33.

Remes-Lenicov, A.M.M. & Rioja, T.C. 2007. Presencia de *Toya argentinensis* (Muir, 1929) (Hemiptera: Auchenorrhyncha: Delphacidae) en el Valle de Azapa, Región de Arica y Parinacota, Chile. *Acta Entomológica Chilena* **31**: 11-14.

Remes-Lenicov, A.M.M. & Tesón, A. 1978. Contribución al estudio de los Fulgóridos Argentinos I (Homoptera, Fulgoroidea, Delphacidae). *Revista de la Sociedad Entomológica Argentina* **37**(1-4): 17-22.

Remes-Lenicov, A.M.M.; Tesón, A.; Dagoberto, E. & Huguet, N. 1985. Hallazgo de uno de los vectores del "Mal de Río Cuarto" del maíz. *Gaceta Agronómica* **5**(25): 251-258.

Remes-Lenicov, A.M.M. & Virla, E.G. 1996. Descripción de los estados inmaduros de *Delphacodes haywardi* y notas acerca de su comportamiento en condiciones de laboratorio. *Revista de la Sociedad Entomológica Argentina* **55**(1-4): 165-174.

Remes-Lenicov, A.M.M. & Virla, E.G. 1999. Delfácidos asociados al cultivo de maíz en la República Argentina (Insecta-Homoptera-Delphacidae). *Revista de la Facultad de Agronomía, La Plata* **104**(1): 1-15.

Remes-Lenicov, A.M.M.; Zerbino, S. & Demaría, M. 2000. Especies de delfácidos (Homoptera, Delphacidae) presentes en el cultivo de maíz, en Uruguay. *Agrociencia* **4**(1): 93-95.

Rioja, T.C.; Vargas, H.E. & Bobadilla, D.E. 2006. Biología y enemigos naturales de *Peregrinus maidis* (Ashmead) (Hemiptera: Delphacidae) en el valle de Azapa. *Idesia* **24**(1): 41-48.

Rioja, T.C.; Vargas, H.E. & Bobadilla, D.E. 2010. Observaciones sobre la fertilidad diferencial de dos morfotipos alares en *Peregrinus maidis* (Ashmead) (Hemiptera: Delphacidae) en condiciones de laboratorio. *Idesia* **28**(2): 89-95.

Rossi-Batiz, M.F. 2014. *Taxonomía, distribución y biología de la tribu Saccharosydmini (Insecta-Hemiptera-Fulgoromorpha)*. Ph.D. Thesis. Universidad Nacional de La Plata, Facultad de Ciencias Naturales y Museo, La Plata, Argentina, 183 pp.

Spinola, M. 1852. Tribu IV. *Hipocephalocera*. In: Gay, C. *Historia Física y Política de Chile*. Zoología. Tomo 7: 238-305. Maulde et Renou, Paris, France.


Tesón, A. & Remes-Lenicov, A.M.M. 1983. Contribución al estudio de los Fulgóridos Argentinos III. (Homoptera, Fulgoroidea, Delphacidae). *Revista de la Sociedad Entomológica Argentina* 42(1-4): 313-323.

Truol, G.A.; Usugi, T.; Hirao, J.; Arneodo, J.D.; Giménez-Pecci, M.P. & Laguna, I.G. 2001. Transmisión experimental del virus del Mal de Río Cuarto por *Delphacodes kuscheli*. *Fitopatología Brasileira* 26: 39-44.

Urban, J.M.; Bartlett, C.R. & Cryan, J.R. 2010. Evolution of Delphacidae (Hemiptera: Fulgoroidea): combined-evidence phylogenetics reveals importance of grass host shifts. *Systematic Entomology* 35(4): 678-691.

Velázquez, P.D.; Arneodo, J.D.; Guzmán, F.A.; Conci, L.R. & Truol, G.A. 2003. *Delphacodes haywardi* Muir, a new natural vector of Mal de Río Cuarto virus in Argentina. *Journal of Phytopathology* 151: 669-672.


Weglarz, K.M. 2012. *A revision of the genus Chionomus Fennah (Hemiptera: Fulgoroidea: Delphacidae), cleaning out Delphacodes Fieber*. M.Sc. Thesis. University of Delaware, Newark, DE, USA, 154 pp.




Figs. 1-8.- *Chionomus haywardi*, macropterous male. 1.- Habitus, dorsal view. 2.- Habitus, lateral view. 3.- Head, frontal view. 4.- Pygofer, caudal view. 5.- Pygofer, lateral view. 6.- Anal tube, lateral view. 7.- Right paramere, caudal view. 8.- Aedeagus, lateral view.




Figs. 9-16.- *Syndelphax dissipatus*, brachypterous male. 9.- Habitus, dorsal view. 10.- Habitus, lateral view. 11.- Head, frontal view. 12.- Pygofer, caudal view. 13.- Pygofer, lateral view. 14.- Anal tube, lateral view. 15.- Right paramere, caudal view. 16.- Aedeagus, lateral view.



Figs 17-24.- *Dicranotropis bipectinata*, brachypterous male. 17.- Habitus, dorsal view. 18.- Habitus, lateral view. 19.- Head, frontal view. 20.- Pygofer, caudal view. 21.- Pygofer, lateral view. 22.- Anal tube and aedeagus, lateral view. 23.- Parameres, caudal view. 24.- Right paramere, latero-caudal view.



Figs. 25-32.- *Delphacodes darwini*, brachypterous male. 25.- Habitus, dorsal view. 26.- Habitus, lateral view. 27.- Head, frontal view. 28.- Pygofer, caudal view. 29.- Pygofer, lateral view. 30.- Anal tube, lateral view. 31.- Right paramere, caudal view. 32.- Aedeagus, lateral view.



Figs. 33-41.- *Delphacodes kuscheli*, brachypterous male. 33.- Habitus, dorsal view. 34.- Habitus, lateral view. 35.- Head, frontal view. 36.- Pygofer, caudal view. 37.- Pygofer, lateral view. 38.- Anal tube, lateral view. 39.- Anal tube, caudal view. 40.- Right paramere, caudo-lateral view. 41.- Aedeagus, lateral view.

42





Fig. 42.- Distribution map of *Dicranotropis bipectinata* and *Syndelphax dissipatus*.

Fig. 43.- Distribution map of *Delphacodes kuscheli* and *Chionomus haywardi*.

Fig. 44.- Distribution map of *Delphacodes darwini*.

43



44

