



7th European Hemiptera Congress

and

9th International Workshop on Leafhoppers
and Planthoppers of Economic Importance



Programme

Abstracts of Talks and Posters

Seggau Castle / Graz (Austria), July 19th - 24th, 2015



Contributions to the Auchenorrhyncha fauna of the Turkish part of Thrace (Hemiptera: Fulgoromorpha and Cicadomorpha)

Ahmet DURSUN¹, Meral FENT²

1 Amasya University Faculty of Arts and Sciences Department of Biology, İpekköy, Amasya-Turkey

2 Trakya University Faculty of Sciences Department of Biology, Edirne-Turkey

Corresponding author: ahmetdursun55@hotmail.com

Key words: Turkey fauna, Thrace, Fulgoromorpha, Cicadomorpha

Fulgoromorpha and Cicadomorpha specimens were collected in different localities in Turkish Thrace in the spring and summer period of 2014. The collected material was evaluated and *Metcalfa pruinosa* (Say, 1830) (Flatidae), *Tettigometra laeta* Herrich-Schäffer, 1835 (Tettigometridae), *Scorlupella discolor* (Germar, 1821) (Issidae) and *Euides basilinea* (Germar, 1821) (Delphacidae) from Fulgoromorpha and *Eupelix cuspidata* (Fabricius, 1775) (Cicadellidae) from Cicadomorpha were recorded for the first time for region of Thrace. *E. cuspidata* was so far only known from the Asian part (Anatolia) of Turkey. *S. discolor* was found in several localities, and *T. laeta* and the exotic species *M. pruinosa* were found only in one locality, respectively. *E. basilinea* was already published from Turkey but without any data on the locality. Taxonomical characteristics of the species and their distributions in Turkey and within the Palaearctic Region were given.