

**2nd INTERNATIONAL CONGRESS
CONCERNING THE RHYNCHOTA FAUNA
OF BALKAN AND ADJACENT REGIONS**

PROCEEDINGS

(Edited by SAKIS DROSOPPOULOS)

Sponsors

Ministry of Culture and Science

Prefecture of Florina County

Hellenic Zoological Society

18-22 August 1986

Mikrolimni - Prespa

GREECE

The International Coordinating Committee

Dr. H. Abdul-Nour

Dr. M. Asche

Dr. S. Drosopoulos

Dr. V. D' Urso

Dr. M. Gogala

Dr. H. Hoch

Dr. M. Josifov

Dr. M. Klein

Dr. N. Lodos

The Organizing Committee

Dr. S. Drosopoulos

Dr. M. Loukas

Mr. P. Petrakis

Dr. L. Argyriou

Miss V. Kokkinos

Contributions

The zoogeographical character of Balkan Heteroptera.	M. JOSIEOV	6
A preliminary list and some notes on the Cicadomorpha (Homoptera-Auchenorrhyncha) collected in Greece.	S. DROSOPoulos, M. ASCHE and H. HOCH	8
Heteropteran fauna of Sicilia: present state and characteristics.		
Cicadellidae of Lebanon: new facts and ideas. (Homoptera - Auchenorrhyncha).	H. ABDUL-NOUR	15
The Psylloidea of Lebanon: preliminary records and prospects.	N. ZEIDAN-GEZE	17
Cixiidae and Delphacidae (Homoptera-Fulgoroidea) from Lebanon – a preliminary synopsis.	M. ASCHE and H. HOCH	18
Heteroptera of the island of Salina (Sicily, Italy).	S. IPPOLITO	20
On the Auchenorrhyncha (Homoptera) from Aeolian island (Sicily, Italy).	V. D'URSO	23
On endemic and little known Heteroptera from Crete.	E. HEISS	28
The <i>Arocephalus longiceps</i> (Kbm.) – (Homoptera Cicadelloidea Paralimnini): problems in intraspecific variability, geographical distribution and speciation.	M. ASCHE	30
Patterns of geographic distribution in the planthopper genus <i>Hyalesthes</i> Sign. (Homoptera Fulgoroidea Cixiidae): a phylogenetic approach.	H. HOCH	31
Species-discrimination and geographic distribution in the cydnid genus <i>Tritomegas</i> (Heteroptera, Cydnidae).	D. KAMMERSCHEN	33
Some remarks on the phylogeny of the Lygaeidae based on the male genital system.	I. HOPP	35
Electrophoretic studies on <i>Alebra albostriella</i> complex (Homoptera, Cicadellidae, Typhlocybinae).	M. LOUKAS and S. DROSOPoulos	42
Biosystematic studies on the genus <i>Alebra</i> Fieber.	M. GILLHAM	44
A parthenogenetic planthopper found in Greece.	P. W.F. de VRIJER	46
Structures producing vibrational signals in Heteroptera.	M. GOGALA	47
The vibratory membranes in the genus <i>Euscelis</i> .	H. STRUBING and G. SCHWARZ-MITTELSTAEDT	49
Anti-predatory defence of some Rhynchota especially as it applies to avian insectivores.	D.L. EVANS	53
Structure organization and dynamics of Hemiptera-Plant communities of a mediterranean area: preliminaries and methodology.	P. PETRAKIS, V. TSELEPATIOTI-PETRAKI and S. DROSOPoulos	54
DAPROPHECO: A specialized database system for integrated studies in Plant-Hemiptera communities.	P. PETRAKIS	62
Seasonal aspects and most important biotopes of Greece with reference to Hemiptera (Auchenorrhyncha-Homoptera and Heteroptera).	S. DROSOPoulos	64
Round table on management of nature reserves in Prespa.		69

A preliminary list and some notes on the Cicadomorpha (Homoptera-Auchenorrhyncha) collected in Greece

By S. DROSOPoulos¹, M. ASCHE², and H. HOCH²

1. Benaki Phytopathological Institute Kiphissia - Athens, Greece

2. Laboratory of Zoology, University of Marburg, P.O. Box 1929, W. Germany

As it has been reported for the planthoppers (Drosopoulos, Asche and Hoch, 1985) several localities in Greece have been investigated in the last 10 years as far as Auchenorrhyncha (Homoptera) and Heteroptera are concerned. The material collected counts more than 100.000 specimens. Recently we started a qualitative analysis of the material belonging to the infraorder Cicadomorpha. This analysis resulted in about 400 species (Table 1). More than half of these species (232) are reported for the first time in Greece, while 12 of these species are new and will be described elsewhere. Since this analysis is not complete 10-20 additional species are expected to be found in our material. In addition, there are 37 species which have been reported from Greece but not yet found by us (Table 2). Although it is already known, that some of the species included in Table 2 are erroneously reported from Greece and others are expected to be synonyms, the total number of Cicadomorpha found, so far, in Greece is approximately 450. Considering that many and various types of biotopes have been studied in Greece and the Cicadomorpha fauna of adjacent countries (especially that of Yugoslavia which is rather well investigated), it is estimated that this material covers 85-90% of what is expected to be found in Greece.

It would have been possible to split this material in decades of papers reporting either species from certain geographic areas or biotopes of Greece, or groups of this infraorder from the total area of Greece. However, since the infraorder Fulgoromorpha Evans, 1946 studied by us includes about 300 species and a key to

the species of both infraorders of Auchenorrhyncha is lacking from the eastern mediterranean region, we are planning to publish in future two volumes which will be more useful than separated papers. Such an ambitious job, however, will require some years of work. Therefore we take the chance to announce here part of our data regarding the infraorder of Cicadomorpha.

By looking at this list, it is easy to notice that there are several questions to be solved especially of some well known genera which require careful revision (e.g. *Macropsis*, *Idiocerus*, *Opius*, *Balclutha*, *Fieberiella*, *Platymetopius*, *Anoplotettix*, *Selenocephalus*, *Psammotettix*, *Chlorita*, *Tamaricella* etc.). Consequently, we do not report here species of *Macropsis*, *Tamaricella* and *Chlorita*, although our material contains about 10, 5 and 5 species of these genera respectively. Particularly, species of these three genera, which do not possess good morphological characters for identification, need to be studied biosystematically. Other genera (e.g. *Anoplotettix*) which exhibit an intraspecific variation even within the same population need also to be studied biosystematically. Already, biosystematics appeared to be very useful for the genus *Alebra* especially the electrophoretic studies (Drosopoulos and Loukas, in prep.). However, biosystematic studies have the disadvantage that they consume much more time than pure morphological analysis of some characters (genitalia of the males). Facing this problem we prefer to leave species of some difficult genera without names. In addition, several species are listed in Table 1 with a question mark

since genitalia structures of the material examined are different from those already reported, or identification characters of some species are not sufficient. Care has been taken also, since populations of some species of the Cicadomorpha may have differentiated in the southernmost area of their distribution as that was found in Delphacidae (Drosopoulos, Asche and

Hoch, 1985).

Many of these species are first records for the Balkan peninsula and some even for Europe as for example *Psammotettix pelikani*, which according to Nast (1972) is reported only from Mongolia. Endemism appeared to be relatively low.

The economic importance of these studies is not assessed in this paper.

Table 1. List of species of Cicadomorpha collected in Greece. The asterisk indicates new records for Greece.

Infraorder: CICADOMORPHA Evans, 1946	<i>Macropsis</i> (10 spp. about)
Family: CICADIDAE Leach, 1815	*cf. <i>Hephatus freyi</i> (Fieber, 1868)
<i>Tibicen plebejus</i> (Scopoli, 1763)	*cf. <i>H. nanus</i> (Herrich-Schäffer, 1835)
<i>Cicadatra alhageos</i> (Kolenati, 1857)	<i>Agallia consobrina</i> Curtis, 1833
<i>C. atra</i> (Olivier, 1790)	<i>A. minuta</i> Melichar, 1896
<i>C. hyalina</i> (Fabricius, 1798)	*?A. sp. n.
<i>Cicada ornata</i> Linnaeus, 1758	<i>Anaceratagallia laevis</i> (Ribaut, 1935)
<i>Tibicina haematoxides</i> (Scopoli, 1763)	<i>A. ribauti</i> (Ossiannilsson, 1938)
<i>Pagiphora annulata</i> (Brulle, 1832)	<i>A. venosa</i> (Fourcroy, 1785)
<i>P. ashei</i> Kartal, 1978	<i>Astroagallia sinuata</i> (Mulsant et Rey, 1855)
<i>Cicadetta dimissa</i> (Hagen, 1856)	* <i>Dryodurgades anatolicus</i> Dlabola, 1957
<i>C. gastrica</i> (Stål, 1854)	* <i>D. dlabolai</i> Wagner, 1963
<i>C. hageni</i> Fieber, 1872	<i>D. reticulatus</i> (Herrich-Schäffer, 1834)
<i>C. montana</i> (Scopoli, 1772)	* <i>Rhytidodus decimusquartus</i> (Schrank, 1776)
*?C. sp.n.	* <i>Populicerus albicans</i> (Kirschbaum, 1868)
<i>C. tibialis</i> (Panzer, 1788)	* <i>P. confusus</i> (Flor, 1861)
Family: CERCOPIDAE Leach, 1815	* <i>P. nitidissimus</i> (Herrich-Schäffer, 1835)
<i>Cercopsis arcuata</i> Fieber, 1844	<i>P. populi</i> (Linnaeus, 1761)
<i>C. intermedia</i> Kirschbaum, 1868	* <i>Tremulicerus</i> (?) <i>dimidiatus</i> (Ribaut, 1952)
<i>C. sanguinolenta</i> (Scopoli, 1763)	<i>T. mesopyrrhus</i> (Kirschbaum, 1868)
<i>C. vulnerata</i> Rossi, 1807	* <i>Idiocerus herrichii</i> Kirschbaum, 1868
<i>Haematoloma dorsatum</i> (Ahrens, 1812)	* <i>I. lituratus</i> (Fallen, 1806)
<i>Lepyronia coleoptrata</i> (Linnaeus, 1758)	<i>I. ocellaris</i> (Mulsant et Rey, 1855)
* <i>Neophilaenus albipennis</i> (Fabricius, 1798)	* <i>I. stigmatical</i> Lewis, 1834
<i>N. campestris</i> (Fallen, 1805)	<i>I. vicinus</i> Melichar, 1898
* <i>N. exclamationis</i> (Thunberg, 1784)	<i>Acericerus rotundifrons</i> (Kirschbaum, 1868)
* <i>N. lineatus</i> (Linnaeus, 1758)	*A. sp. n.
* <i>N. minor</i> (Kirschbaum, 1868)	<i>A. vittifrons</i> (Kirschbaum, 1868)
<i>Aphrophora alni</i> (Fallen, 1805)	<i>Metidiocerus rutilans</i> (Kirschbaum, 1868)
* <i>A. corticea</i> Germar, 1821	<i>Viridicerus ustulatus</i> (Mulsant et Rey, 1855)
* <i>A. costalis</i> Matsumura, 1903	<i>Sulamicerus stali</i> (Fieber, 1868)
* <i>A. salicina</i> (Goeze, 1778)	<i>Balcanicerus larvatus</i> (Herrich-Schäffer, 1835)
<i>Philaenus signatus</i> Melichar, 1896	* <i>B. pruni</i> (Ribaut, 1952)
<i>P. spumarius</i> (Linnaeus, 1758)	cf. <i>Batracomorphus irroratus</i> Lewis, 1834
* <i>P. sp. n.</i>	*cf. <i>Iassus lanio</i> (Linnaeus, 1761)
Family: MEMBRACIDAE Rafinesque, 1815	cf. <i>I. scutellaris</i> (Fieber, 1868)
* <i>Gargara genistae</i> (Fabricius, 1775)	cf. <i>Penthimia nigra</i> (Goeze, 1778)
<i>Gentrotus cornutus</i> (Linnaeus, 1758)	*cf. <i>Paradyromidium paradoxum</i> (Herrich-Schäffer, 1837)
<i>Stictocephala bisonia</i> Kopr & Yonke, 1977	<i>Eupelix cuspidata</i> (Fabricius, 1775)
Family: CICADELLIDAE Latreille, 1825	* <i>Hecalus glaucescens</i> (Fieber, 1866)
<i>Ulopa trivialis</i> Germar, 1821	* <i>H. sp.</i>
<i>Megophthalmus scabripennis</i> Edwards, 1915	<i>Aphrodes bicincta</i> (Schrank, 1776)
* <i>M. scanicus</i> (Fallen, 1806)	* <i>A. makarovi</i> Zachvatkin, 1948
* <i>Lebra aurita</i> (Linnaeus, 1758)	* <i>A. (?) carinatus</i> (Stål, 1864)
*cf. <i>Oncopsis flavigollis</i> (Linnaeus, 1761)	<i>Anoscopus albifrons</i> (Linnaeus, 1758)
* <i>O. sp.</i>	*cf. <i>A. albiger</i> (Germar, 1821)

- Planaphrodes bifasciata* (Linnaeus, 1758)
 **P. nigrita* (Kirschbaum, 1868)
P. trifasciata (Fourcroy, 1785)
 **Stroggylocephalus agrestis* (Fallen, 1806)
 **Evacanthus acuminatus* (Fabricius, 1794)
 **E. interruptus* (Linnaeus, 1758)
Cicadella viridis (Linnaeus, 1758)
 **Grypotes puncticollis* (H. S., 1834)
G. staurus Ivanoff, 1885
Goniognathus bolivari (Melichar, 1907)
G. brevis (Herrich-Schaffer, 1835)
G. guttulinervis (Kirschbaum, 1868)
 **Opsius cypriacus* Lindberg, 1958
 *?*O. euxinus* Dlabola, 1965
 *?*O. jucundus* (Lethierry, 1871)
 *?*O. lethierryi* Wagner, 1942
 ?*O. pallasi* (Lethierry, 1874)
O. stactogalus (Fieber, 1866)
 *cf. *Orosius orientalis* (Matsumura, 1914)
Neoaliturus fenestratus (Herrich-Schaffer, 1834)
 **N. gattulatus* (Kirschbaum, 1868)
N. haematoceps (Mulsant et Rey, 1855)
N. tenellus (Baker, 1896)
 **Concavifer bokarensis* Kartal, 1982
Jubrinia distincta Linnauvori, 1962
Nesoclutha erythrocephala (Ferrari, 1882)
Cicadulina bipunctata (Melichar, 1904)
 **Balchutha boica* Wagner, 1950
 **B. nicolasi* (Lethierry, 1876)
B. punctata (Fabricius, 1775)
 ?*B. rhenana* Wagner, 1939
B. rosea (Scott, 1876)
B. saltuella (Kirschbaum, 1868)
 *?*Macrostelus chobauti* Ribaut, 1952
 *?*M. sieberi* (Edwards, 1889)
 *?*M. forficula* (Ribaut, 1927)
 *?*M. frontalis* (Scott, 1875)
 *?*M. halophilus* (Horvath, 1903)
 *?*M. horvathi* (Wagner, 1935)
M. laevis (Ribaut, 1927)
M. ossianilsoni (Lindberg, 1954)
M. quadripunctulatus (Kirschbaum, 1868)
M. ramosus Ribaut, 1952
 **M. salsolae* (Puton, 1872)
 **M. septemnotatus* (Fallen, 1806)
 **M. sexnotatus* (Fallen, 1806)
 **M. viridigriseus* (Edwards, 1922)
 **Sagatus punctifrons* (Fallen, 1826)
 **Deltoccephalus pulicaris* (Fallen, 1806)
 **Recilia coronifera* (Marshall, 1866)
R. schmidtgeni (Wagner, 1939)
Stymphalus rubrolineatus (Stål, 1855)
 **Chiasmus conspurcatus* (Perris, 1857)
 *cf. *Doratura concors* Horvath, 1903
 *?*D. exilis* Horvath, 1903
D. heterophyla Horvath, 1903
D. homophyla (Flor, 1861)
 **D. impudica* Horvath, 1897
D. medvedevi Logvinenko, 1961
D. stylata (Bohemian, 1847)
 **Aconura oculata* (Lindberg, 1948)
Aconurella prolixa (Lethierry, 1885)
Fieberiella duffelsi Wagner, 1963
F. florii (Stål, 1864)
F. ida Dlabola, 1965
 **F. macchiai* Linnauvori, 1962
 **F. sp. n.*
Synophropsis lauri (Horvath, 1897)
- **Placotettix taeniatifrons* (Kirschbaum, 1868)
 **Docotettix cornutus* Ribaut, 1948
 **Doratulina acuticeps* (Ribaut, 1948)
D. instabilis (Ribaut, 1948)
 **D. pallifrons* (Horvath, 1897)
Exitianus capicola (Stål, 1855)
 **Tetartostylus illyricus* (Kirschbaum, 1868)
 **Osbornellus macchiai* (Lindberg, 1948)
Platymetopius complicatus Nast, 1972
 *?*P. dorsofrenestratus* Dlabola, 1958
 **P. filigranus* (Scot, 1876)
 cf. *P. guttatus* Fieber, 1869
 **P. henribaudi* Dlabola, 1961
P. major (Kirschbaum, 1868)
 **P. rostratus* (Herrich-Schaffer, 1934)
 **P. undatus* (De Geer, 1773)
P. wilhemwagneri Dlabola, 1980
Anoplotettix beieri Wagner, 1959
A. fuscovenosus (Ferrari, 1882)
A. graecus Remane, 1966
 **A. putoni* Ribaut, 1952
 *cf. *A. sahyancii* Dlabola, 1970
 **Idiodomus cruentatus* (Panzer, 1799)
Lamprotettix nitidulus (Fabricius, 1787)
 **Paramesodes lucaniae* (Dlabola, 1959)
Allygus atomarius (Fabricius, 1794)
 **A. commutatus* Fieber, 1872
 **A. communis* (Ferrari, 1882)
 *cf. *A. mayri* (Kirschbaum, 1868)
 **A. mixtus* (Fabricius, 1794)
A. modestus Scott, 1876
Phlepsius intricatus (Herrich-Schaffer, 1838)
 ?*P. ornatus* (Perris, 1857)
Selenocephalus abbreviatus Melichar, 1896
 **S. anatolicus* Dlabola, 1957
 **S. obsoletus* (Germar, 1817)
S. pallidus Kirschbaum, 1868
 **S. pallens* Lindberg, 1948
 **Graphocraerus ventralis* (Fallen, 1806)
Rhytidostylus kaloneroi Asche, 1980
 **Hardya anatolica* Zachvatkin, 1946
 **Eohardya fraudulenta* (Horvath, 1903)
 **Sardius argus* (Marshall, 1866)
 **Paluda adumbrata* (C. Sahlberg, 1842)
 **P. preysleri* (Herrich-Schaffer, 1838)
 **P. sp. n.*
 **P. vitripennis* (Flor, 1861)
 **Elymana sulphurella* (Zetterstedt, 1828)
 **Cicadula lineatopunctata*, (Matsumura, 1908)
C. placida (Horvath, 1897)
 **C. quadrinotata* (Fabricius, 1794)
 **Mocydia crocea* (Herrich-Schaffer, 1837)
 **M. sp. n.*
Mocydiopsis longicauda Remane, 1961
 **M. monticola* Remane, 1961
 **Spseudotettix subfuscatus* (Fallen, 1806)
 **Hesium domino* (Reuter, 1880)
 **Thamnotettix confinis* Zetterstedt, 1828
T. creticus Dlabola, 1974
 **T. dilutior* (Kirschbaum, 1868)
 **T. exemptus* Melichar, 1896
T. minoidis Dlabola, 1974
T. thrax Dlabola, 1965
 **T. zelleri* (Kirschbaum, 1868)
 **Macustus grisescens* (Zetterstedt, 1828)
 **Athysanus argentarius* Metcalf, 1955
 **Handianus ignoscus* (Melichar, 1896)
 **H. procerus* (Herrich-Schaffer, 1835)

- **H.* sp.
 - **Stictocoris picturatus* (C. Sahlberg, 1842)
 - **Scleroracus* sp.
 - **Limotettix striola* (Fallen, 1806)
 - **Laburus impictifrons* (Boheman, 1852)
 - Euscelidius schenckii* (Kirschbaum, 1868)
 - Conosanus obsoletus* (Kirschbaum, 1858)
 - Euscelis alsius* Ribaut, 1952
 - E. distinguendus* (Kirschbaum, 1858)
 - **E. incisus* (Kirschbaum, 1858)
 - E. lineolatus* Brulle, 1832
 - **Streptanus aemulans* (Kirschbaum, 1868)
 - **S. albanicus* (Horvath, 1916)
 - **Artianus manderstjernii* (Kirschbaum, 1868)
 - ?*Phycottetix truncatipennis* (Rambur, 1840)
 - Melilaius matsumuri* (Metcalf, 1955)
 - **Oxytettigella viridinervis* (Kirschbaum, 1868)
 - Aglena ornata* (Herrich-Schaffer, 1838)
 - **Paramesus major* Haupt, 1927
 - **P. paludosus* Ribaut, 1952
 - **Paralimnus phragmitis* (Boheman, 1847)
 - **P. picturatus* Haupt, 1930
 - **P. zachvatkini* Emeljanov, 1964
 - Arocephalus longiceps* (Kirschbaum, 1868)
 - **A. (Ariellus) languidus* (Flor, 1861)
 - *cf. *A. (Ariellus) punctum-siculus* D'Urso, 1978
 - A. (Ariellus) superpunctum* R. et A., 1980
 - **Psamnotettix adriaticus* Wagner, 1959
 - **P. alienus* (Dahlbom, 1850)
 - *cf. *P. cephalotes* (Herrich-Schaffer, 1834)
 - **P. confinis* (Dahlbom, 1850)
 - **P. helvolus* (Kirschbaum, 1868)
 - **P. inexpectatus* Remane, 1965
 - **P. ornaticeps* (Horvath, 1897)
 - **P. pelikani* Dlabola, 1965
 - **P. percictus* Logvinenko, 1978
 - **P. pictipennis* (Kirschbaum, 1868)
 - **P. poecilus* (Flor, 1861)
 - **P.* sp. n.
Ebarrius cognatus (Fieber, 1869)
 - **E. interstinctus* (Fieber, 1869)
 - **Adarrus bellevoyei* (Puton, 1877)
 - A. ellisi* Dlabola, 1980
 - **A. falcatus* Linnauvori, 1952
 - A. graecornatus* Remane et Asche, 1980
 - **A. multinotatus* (Boheman, 1847)
 - **A. reductus* (Melichar, 1897)
 - **Kazachstanicus aitosicus* Dlabola, 1965
 - Jassargus graecobitus* Remane et Schulz, 1976
 - **J. neglectus* (Then, 1896)
 - J. obtusivalvis* (Kirschbaum, 1868)
 - **J. reptetus* (Fieber, 1869)
 - **Diplocolenus abdominalis* (Fabricius, 1803)
 - **D. bohemani* (Zetterstedt, 1840)
 - **D. frauenfeldi* (Fieber, 1869)
 - **D. nigritrons* (Kirschbaum, 1868)
 - **Arthaldeus pascuellus* (Fallen, 1826)
 - **A. striifrons* (Kirschbaum, 1868)
 - **Sorhoanus medius* (Mulsant et Rey, 1855)
 - **Rhoanarus hypochlorus* (Fieber, 1869)
 - **Calamotettix* sp.
 - **Mocuellus quadricornis* Dlabola, 1949
 - **Nanosius chloroticus* (Melichar, 1896)
 - Pseupalus graecanarus* Remane et Asche, 1980
- Subfamily: **TYPHLOCYBINAЕ** Kirschbaum, 1868
- Alebra albostriella* (Fallen, 1826)
 - **A.* sp. n.
- **A.* sp. n.
 - **A. wahlbergi* (Boheman, 1845)
 - *cf. *Erythria aureola* (Fallen, 1806)
 - **Liguropia juniperi* (Lethierry, 1876)
 - ?*Emelyanoviana mollicula* (Boheman, 1845)
 - ?*Dikraneura variata* Hardy, 1850
 - **Micantulina micantula* (Zetterstedt, 1840)
 - M. stigmatipennis* (Mulsant et Rey, 1855)
 - M. teurci* (Cerutti, 1938)
 - cf. *Wagneriala sinuata* (Then, 1897)
 - **Kybos butleri* (Edwards, 1908)
 - *?*K. populi* (Edwards, 1908)
 - K. virgator* (Ribaut, 1933)
 - Empoasca decadens* Paoli, 1932
 - E. decipiens* Paoli, 1930
 - E. solani* (Curtis, 1846)
 - **E. vitis* (Gothe, 1875)
 - **Jacobiasca lybica* (Bergevin et Zanon, 1922)
 - **Chlorita paolii* (Ossiannilsson, 1939)
 - C. spp.*
 - Fagocyba cruenta* (Herrich-Schaffer, 1838)
 - **Edwardsiana crataegi* (Douglas, 1876)
 - ?*E. cretica* Dworakowska, 1971
 - **E. flavescens* (Fabricius, 1794)
 - E. gratiosa* (Boheman, 1852)
 - **E. hippocastani* (Edwards, 1888)
 - **E. platanicola* (Vidano, 1961)
 - ?*E. rosae* (Linnaeus, 1758)
 - **E. rosaesugans* (Cerutti, 1939)
 - **Eupterycyba jucunda* (Herrich-Schaffer, 1837)
 - **Linnauvoriola sexmaculata* (Hardy, 1850)
 - ?*Ficocyba sicaria* (Horvath, 1897)
 - Lindbergina loewi* (Lethierry, 1884)
 - **L. aurovittata* (Douglas, 1875)
 - **L. spoliata* (Horvath, 1907)
 - L. cretica* Asche, 1980
 - Ribautiana alces* (Ribaut, 1931)
 - R. cruciata* (Ribaut, 1931)
 - **R. tenerrima* (Herrich-Schaffer, 1834)
 - **R. ulmi* (Linnaeus, 1758)
 - **Typhlocyba quercus* (Fabricius, 1777)
 - Eurhadina concinna* (Germar, 1831)
 - **E. kirschbaumi* Wagner, 1937
 - **E. ribauti* Wagner, 1935
 - **E.* sp. n.
 - **Eupteryx adspersa* (Herrich-Schaffer, 1838)
 - **E. andalusica* Ferrari, 1882
 - **E. atropunctata* (Goeze, 1778)
 - **E. aurata* (Linnaeus, 1758)
 - **E. calcarata* Ossiannilsson, 1936
 - **E. castelvecchica* Dlabola, 1967
 - **E. cyclops* Matsumura, 1906
 - E. cypria* (Ribaut, 1948)
 - E. filicum* (Newman, 1853)
 - ?*E. graeca* (Lindberg, 1949)
 - ?*E. gravesteini* Dlabola 1957
 - E. gyaudagica* Dlabola, 1957
 - **E. heydenii* (Kirschbaum, 1868)
 - E. insulana* (Ribaut, 1948)
 - E. melissae* Curtis, 1837
 - **E. notata* Curtis, 1837
 - **E. ribauti* Dworakowska, 1972
 - E. stachydearum* (Hardy, 1850)
 - **E. tenella* (Fallen, 1806)
 - **E. thoulessi* Edwards, 1926
 - E. utricae* (Fabricius, 1803)
 - ?*E. vicaria* Linnauvori, 1968

- **E. vittata* (Linnaeus, 1758)
 **E. zelleri* (Kirschbaum, 1868)
 **Aguriahana germari* (Zetterstedt, 1840)
 **A. stellulata* (Burmeister, 1841)
 **Zygiella albifrons* Horvath, 1897
Z. pulchra Low, 1885
 **Alnetoidia alneti* (Dahlbom, 1850)
Hauptidia duffelsi Dlabola, 1979
 **H. provincialis* (Ribaut, 1931)
H. soosi (Dlabola, 1952)
 cf. *Kropka unipunctata* (Dlabola, 1957)
 **Zyginidia adamczewskii* Dworakowska, 1970
 **Z. mocsaryi* (Horvath, 1910)
 **Z. pullula* (Bohemian, 1845)
- **Z. ribauti* Dworakowska, 1970
Z. sohrab Zachvatkin, 1947
 **Zygina angusta* Lethierry, 1874
 **Z. discolor* Horvath, 1897
 **Z. titihide* Ferrari, 1882
Hypericiella hyperici (Herrich-Schaffer, 1836)
 **Arboridia cantoreanica* Dworakowska, 1970
 **A. parvula* (Bohemian, 1845)
 **A. ribauti* (Ossiannilsson, 1937)
 **A. velata* (Ribaut, 1952)
Frutiodia bisignata (Mulsant et Rey, 1855)
 **F. sp.*
Tamaricella spp.
Fruticidia nigricostalis (Matsumura, 1908)
- Dlabola, J.
 zur
 geb
 21-
 Dlabola, J.
 lien
 Län
 25-
 Dlabola, J.
 dena
 Cica
 Nat.
 Dlabola, J.
 Med
 weite
 (Hon
 Acta

Table 2. Species of Cicadomorpha reported from Greece. One asterisk indicates species of uncertain generic position and two asterisks species erroneously reported from Greece.

CICADIDAE

- Cicada mordoganensis* Bulard, 1979
Cicadetta aestuans (Fabricius, 1794)
C. albipennis Fieber, 1876
C. brullei Fieber, 1876
C. flaveola (Brulle, 1832)
Euryphorà cantans (Fabricius, 1794)

CERCOPIDAE

- Triecphorella geniculata* (Horvath, 1881)

CICADELLIDAE

- ***Oncopsis tristis* (Zetterstedt, 1840)
Idiocerus poecilus (Herrich-Schaffer, 1835)
Neoaliturus inscriptus (Haupt, 1927)
Balclutha hebe (Kirkaldy, 1906)
Phlogotettix cyclops (Mulsant et Rey, 1855)
Osbornellus consanguineus Dlabola, 1967
 ***Anoplotettix ibericus* Remane, 1966
A. malickyi Dlabola, 1984
A. cruciatus Dlabola, 1984
A. rodosicus Dlabola, 1974

Acknowledgements

The authors wish to express their sincere gratitude to Prof. R. Remane for his interest, criticism and valuable help during the identification of the species. We thank also Drs H. Abdul-Nour, H. Duffels and W.H. Gravestein for comments.

Thanks are also due to the Fachbereich of Biologie (Zoologie), Philipps-Universität, Marburg for hospitality and to «Deutscher Akademischer Austauschdienst» (DAAD) for financial support to the first author for a period of six months.

Bibliography

- Asche, M., 1980. *Rhytidostylus kaloneroi* nov. spec., eine neue Cicadellidae aus Nordwest-Griechenland (Homoptera Cicadina Cicadellidae Deltocephalinae).

- Selenocephalus heraki* Dlabola, 1977
S. moreanus Dlabola, 1977
S. planus (Turton, 1802)
Stenometopius fraudulentus (Horvath, 1903)
Euscelidius mundus (Haupt, 1927)
Streptanus jostovi Dlabola, 1957
Parasemus ancorifer Linnauvori, 1953
Goldeus helenicus (Dlabola, 1974)
Psammotettix provincialis (Ribaut, 1925)
Arocephalus laconicus (Dlabola, 1980)
 **Cicada longicornis* (Poda, 1761)

TYPHLOCYBINAЕ

- Chlorita beieri* Dlabola, 1959
C. tessellata Lethierry, 1884
Edwardsiana iranicola Zachvatkin, 1947
Eupteryx collina (Flor, 1861)
Zyginidia scutellaris (Herrich-Schaffer, 1838)
Zygina krueperi Fieber, 1884
Z. nivea (Mulsant et Rey, 1855)
 **Anomia lepida* Fieber, 1884
 **Chlorita punctum* Lethierry, 1884

– Marburger Ent. Publ. 1 (3): 123-132.
 Asche, M., 1980. *Lindbergina cretica* nov. spec., eine neue Typhlocybidae von Kreta (Homoptera Auchenorrhyncha Cicadellidae Typhlocybinae). – Marburger Ent. Publ. 1 (4): 37-44.

Dlabola, J., 1974. Zur Taxonomie und Chorologie einiger mediterraner Zikaden (Homoptera: Auchenorrhyncha). – Acta Zool. Acad. Scient. Hung. 20 (3-4): 289-308.

Dlabola, J., 1974. Übersicht der Gattungen *Anoplotettix*, *Goldeus* und *Thamnotettix* mit Beschreibungen von 7 neuen mediterranen Arten (Homoptera Auchenorrhyncha). – Acta faun. ent. Mus. Nat. Pragae, 15: 103-130.

Dlabola, J., 1977. Neue Zikadentaxone von *Mycterodus*, *Erythria*, *Selenocephalus* und *Goldeus*. – Acta Zool. Sci. Nat. Hung., 23: 279-292.

- Dlabola, J., 1977. Chorologische Ergänzungen zur Zikadenfauna des Mittelmeergebietes. Sbor. Nár mus. Praha, 33: 21-40.
- Dlabola, J., 1979. Neue Zikaden aus Anatolien, Iran und aus südeuropäischen Ländern. – Acta zool. Acad. Sci. Hung., 25: 235-257.
- Dlabola, J., 1980. Neue griechische Zikadenarten der Fam. Cixiidae, Issidae und Cicadellidae. – Acta faun. ent. Mus. Nat. Pragae, 16: 5-13.
- Dlabola, J., 1984. Neue Zikadenarten aus Mediterraneum und dem Iran mit weiteren Beiträgen zur iranischen Fauna (Homoptera - Auchenorrhyncha). – Acta Mus. Nat. Pragae, 40: 21-64.
- Drosopoulos, S., 1980. Hemipterological studies in Greece. Part II Homoptera - Auchenorrhyncha. A catalogue of the reported species. – Biol. Gallo-Hellenica, 9: 187-194.
- Drosopoulos, S., Asche, M. and Hoch H., 1983. Contribution to the Planthopper fauna of Greece (Homoptera, Auchenorrhyncha, Fulgoromorpha, Delphacidae). – Annls. Inst. phytopath. Benaki (N.S.), 14: 19-68.
- Drosopoulos, S. and Loukas, M., 1987. Genetic differentiation between coexisting color-types of the *Alebra albostriella* group (Homoptera: Cicadellidae). (subm. to J. Hered).
- Janković, Lj. and Papović R., 1985. New and rare species in the Yugoslav leafhopper fauna from the family Cicadellidae (Hom. Auchenorrhyncha). – Proc. Fauna SR Serbia. Serbian Academy of Sciences and Arts, Belgrade, 3: 111-147.
- Nast, J., 1972. Palaearctic Auchenorrhyncha (Homoptera). An annotated check list. – Polish Scient. Publ., Warszawa, 550 pp.
- Remane, R. und Asche, M., 1980. Neue Zikaden-Taxa aus dem Tribus Paralimnini DISTANT, 1908, aus dem Mittelmeergebiet (Homoptera Cicadina Cicadelloidea Cicadellidae). – Marburger Ent. Publ. 1 (4): 67-166.
- Schulz, K., 1976. Zur Kenntnis der Gattung *Jassargus* Zachvatkin (Homoptera Auchenorrhyncha). Inaugural-Dissertation, der Philipps-Universität Marburg/Lahn, 256 pp.