

***Parasogata* gen. n., a new genus of the tribe Delphacini with descriptions of two new species from China (Hemiptera, Fulgoromorpha, Delphacidae)**

Zheng-Xiang Zhou^{1,2}, Lin Yang¹, Xiang-Sheng Chen^{1,3,4}

1 Institute of Entomology, Guizhou University, Guiyang, Guizhou, 550025, P.R. China **2** College of Agriculture, Anshun University, Anshun, 561000, P.R. China **3** Guizhou Key Laboratory for Plant Pest Management of Mountainous Region, Guizhou University, Guiyang, Guizhou, 550025, P.R. China **4** Special Key Laboratory for Development and Utilization of Insect Resources of Guizhou, Guizhou University, Guiyang, Guizhou, 550025, P.R. China

Corresponding author: *Xiang-Sheng Chen* (chenxs3218@163.com)

Academic editor: *Mike Wilson* | Received 6 May 2018 | Accepted 10 November 2018 | Published 13 December 2018

<http://zoobank.org/17F67E8D-764D-4802-9E8E-4E256742C773>

Citation: Zhou Z-X, Yang L, Chen X-S (2018) *Parasogata* gen. n., a new genus of the tribe Delphacini with descriptions of two new species from China (Hemiptera, Fulgoromorpha, Delphacidae). ZooKeys 806: 73–85. <https://doi.org/10.3897/zookeys.806.26394>

Abstract

A new planthopper genus *Parasogata* **gen. n.** (Delphacidae: Delphacinae: Delphacini) was described and illustrated with two new species *P. binaria* **sp. n.** and *P. furca* **sp. n.** from south China. A key to species of the new genus is also given.

Keywords

Delphacid, distribution, Fulgoroidea, new taxa, planthopper

Introduction

The planthopper tribe Delphacini Leach, 1815 is the largest clade of Delphacidae, occurring in all ecoregions (excluding Antarctica) and including approximately 1652 species in 319 genera (Bourgoin 2018). In China, 259 species in 135 genera are known (Ding 2006; Dong and Qin 2012; Hou et al. 2013, 2014a, b; Qin 2005, 2006, 2007; Qin et al. 2006, 2008, 2009a, b, 2010, 2011, 2012, 2014; Ren et al. 2015).

Here, a new genus, *Parasogata* gen. n., with two new species, *P. binaria* sp. n. and *P. furca* sp. n., are described and illustrated from China. The new genus is assigned to the Delphacini because the spinal formula of the hind leg 5–7–4, tibial spur large, thin, flattened and bearing a row of fine, black-tipped teeth on the posterior margin; genital diaphragm developed, suspensorium present. The similarities and affinities of the new genus with similar genera are compared and discussed. A key to the species of the new genus is also provided.

Materials and methods

Terminology of morphological and measurements follow Yang and Yang (1986) and the morphological terminology of female genitalia follows Bourgoïn (1993). Measurements of body length equal the distance between the apex of vertex and tip of tegmen. All measurements are in millimeters (mm). Dry specimens were used for the description and illustration. Color pictures for adult habitus were obtained by KEYENCE VHX-1000. External morphology was observed under a stereoscopic microscope Leica Mz 12.5 and characters were measured with an ocular micrometer. The genital segments of the examined specimens were macerated in 10% KOH and drawn from preparations in glycerin jelly using Olympus CX41 and Leica MZ 12.5 stereomicroscope. Illustrations were scanned with Canon CanoScan LiDE 200 and imported into Adobe Photoshop 6.0 for labeling and plate composition.

The type specimens of the new species are deposited in the Institute of Entomology, Guizhou University, Guiyang, Guizhou Province, China (GUGC).

Taxonomy

Genus *Parasogata* gen. n.

<http://zoobank.org/C168D41B-DB7F-4841-BCCC-EF3299BD2EC6>

Figs 9–60

Type species. *Parasogata binaria* sp. n.

Diagnosis. This genus is readily recognized by its large size and vertex, pronotum and mesonotum bearing an uninterrupted white fascia. The genus is most similar to *Sogata* Distant, 1906 but separately by the phallus being up-curved (down-curved in *Sogata* (Ding 2006: figs 281–283)), with a row processes at subapically (without process in *Sogata*).

Description. General color of male yellowish white to brown (Figs 9–12, 35–38). Vertex, pronotum and mesonotum with an uninterrupted white fascia (Figs 9, 35). Vertex, frons, face, antennae yellowish brown to yellowish white (Figs 9–12, 35–38). Pronotum and mesonotum yellowish white (Figs 13, 39). Forewings and hindwings hyaline (Figs 9–12, 35–38). Legs yellowish white (Figs 10, 36). Abdomen yellow (Figs 10, 36). Head including eyes narrower than pronotum (Figs 13, 15, 39, 41). Vertex

subquadrate, anterior margin arched, lateral carinae with slightly concave, submedian carinae uniting at apex. Frons with single median carina, longer in middle line than wide at widest part, widest at apex (Figs 14, 16, 40, 42). Y-shaped carina feeble (Figs 13, 15, 39, 41). Antennae cylindrical, with basal segment shorter than second, reaching frontoclypeal suture (Figs 14, 16, 40, 42). Pronotum with lateral carinae almost attaining hind margin (Figs 13, 15, 39, 41). Posttibial spur with 29–32 distinct teeth along hind margin.

Male genitalia. Anal segment collar-shaped, lateroapical angles produced into processes (Figs 24–25, 50–51). Pygofer in profile wider ventrally than dorsally, laterodorsal angles not produced, in posterior view with opening wider than long, lateral margins well defined, lateral quadrate areas strongly sclerotized, medioventral process absent (Figs 21–23, 47–49). Diaphragm broad (Figs 23, 49). Aedeagus long, tubular, with a row processes at subapically, slightly upward apically (Figs 26, 52). Genital styles simple, widely divergent apically (Figs 27–28, 53–54). Suspensorium large (Figs 29, 55).


Etymology. This generic name “*Parasogata*” refers to its strong similarity to *Sogata*. The name is to be treated as feminine.

Distribution. China.

Remarks. The genus *Parasogata* gen. n. resembles *Sogata* Distant, 1906, *Neometopina* Yang, 1989, *Neunkanodes* Yang, 1989 and *Lisogata* Ding, 2006 in vertex, pronotum and mesonotum with an uninterrupted white fascia, frons with median carina single (Figs 1–8), but differs from these genera by anal segment with two pairs of processes, or with a pair of forked processes (without process in *Lisogata*); aedeagus not forked at half of apex (with forked at half of apex in *Neometopina* and *Neunkanodes*); aedeagus with processes and decurved dorsad apically (without process and decurved ventrad apically in *Sogata*) (Table 1).

Table 1. Differences among *Parasogata* and similar genera.

	<i>Parasogata</i> gen. n.	<i>Neometopina</i>	<i>Sogata</i>	<i>Neunkanodes</i>	<i>Lisogata</i>
Size (mm)	4.72–5.20	4.62–4.82	4.20–4.70	4.40–4.90	4.40–4.70
Frons color	Black with median carina yellowish white	Brown with median carina yellowish white	Black with median carina yellowish white	Brown with median carina yellowish white	Yellow
Y-shaped carina	Feeble	Distinct	Feeble	Feeble	Feeble
Lateral carinae of pronotum	Almost attaining hind margin	Almost attaining hind margin	Almost attaining hind margin	Conspicuous not attaining hind margin	Almost attaining hind margin
Number of teeth of hind tibial spur	29–32	20–23	18–23	23–24	30–38
Hind margin of male pygofer	Not produced	Not produced	Not produced	Produced caudad, lobe-like	Not produced
Processes of male anal segment	Two pairs or one pair with bifurcation	One pair	One pair	One pair	None
Apex of aedeagus	Unforked	Forked	Unforked	Forked	Unforked
Inner basal angle of genital styles	None	None	None	Protruding	None



Figures 1–8. Dorsal and frontal view **1,2** *Neometopina penghuensis* Yang, 1989 **3,4** *Sogata dohertyi* Distant, 1906 **5,6** *Neunkanodes formosana* Yang, 1989 **7,8** *Lisogata zhejiangensis* Ding, 2006. Scale bar: 0.5 mm.

Revised couplets to the key to Chinese Delphacini by Ding (2006)

- 70 Pygofer in profile with posterior margin produced caudad in a lobe (Ding 2006: fig. 314)..... *Neunkanodes* Yang
- Pygofer in profile not produced **71a**
- 71a Aedeagus without processes, apically decurved ventrally (Ding 2006: figs 281–283)..... *Sogata* Distant, 1906
- Aedeagus with distinct processes, apically recurved dorsally (Figs 26, 52).....
..... *Parasogata* gen. n.

Key to species of genus *Parasogata* gen. n. (male)

- 1 Pronotum brown except median carina yellowish white (Figs 13, 17); anal segment with two pairs of processes, each with apex not forked (Figs 24–25) ***P. binaria* sp. n.**
- Pronotum yellow except median carina yellowish white (Figs 39, 43); anal segment with a pair of processes, each with apex forked (Figs 50–51) ***P. furca* sp. n.**

***Parasogata binaria* sp. n.**

<http://zoobank.org/ACC2E082-45A0-4653-9B55-52B8E0182662>

Figs 9–34


Type material. Holotype: ♂, CHINA, Yunnan: Daweishan National Natural Reserve (22°81'N, 103°79'E), 18 Aug. 2017, Y.-J. Sui. Paratypes: 1♂1♀, same data as holotype; 1♂2♀♀, same data as holotype except, 22 Aug. 2017, Q. Luo; 6♂♂5♀♀, same data as holotype except, 19 Aug. 2017, N. Gong.

Measurements. Body length (from apex of vertex to apex of forewing): male 5.10–5.20 mm (n = 9); female 5.90–6.00 mm (n = 8); forewing length: male 4.30–4.42 mm (n = 9); female 5.10–5.12 mm (n = 8).

Diagnosis. Big-sized species with general color yellowish white to yellowish brown, anal segment with two pairs of spinose processes; aedeagus with ten processes subapically and with irregular teeth on ventral side of apex, constriction and bluntly rounded at apex (Fig. 26).

Description. *Coloration.* General color yellowish white to yellowish brown. Head yellowish brown (Figs 13–16). Vertex yellowish white, except along lateral margin dark brown (Figs 13–16). Frons black, except median carinae yellowish white and lateral margins yellowish brown (Fig. 14). Clypeus and genae yellowish brown (Fig. 14). Rostrum yellowish brown, with apex brown. Eyes generally yellow to brown (Figs. 9–16), ocelli dark brown (Figs 10, 12, 14, 16). Antennae yellow (Figs 13–16). Pronotum and mesonotum yellowish brown, except media carinae yellowish white (Figs 13, 15). Forewings with veins dark brown (Figs 9–12). Hindwings pale white, with veins brown. Legs yellowish white to pale yellow, tibiae pale yellow basally, tarsomeres yellowish white (Figs 10, 12). Abdomen brown, except lateral margins yellow (Figs 9–12).

Structure. Head including eyes narrower than pronotum, ratio 0.77:1 (Figs 13, 15). Vertex with anterior margin arched, lateral carinae slightly concave, submedian carinae uniting at apex, longer than wide at base, ratio 1.28:1, narrower at apex than at base, ratio 0.64:1 (Figs 13, 17). Frons longer in middle line than wide at widest part, ratio 2.28:1 (Figs 14, 18), lateral carinae nearly straight (Figs 14, 18). Postclypeus wider at base than frons at apex, slightly longer than wide at base (Figs 14, 18). An-




Figures 9–16. *Parasogata binaria* sp. n. **9, 10** Male habitus (dorsal and lateral views) **11, 12** Female habitus (dorsal and lateral views) **13** Male head and thorax, dorsal view **14** Male front **15** Female head and thorax, dorsal view **16** Female front. Scale bar: 0.5 mm.

tennae cylindrical, basal segment longer than wide, ratio 1.55:1, shorter than second, ratio 0.42:1 (Figs 14, 18). Pronotum shorter than vertex, ratio 0.75:1 (Figs 13, 17). Mesonotum longer than pronotum and vertex combined, ratio 1.25:1 (Figs 13, 17). Posttibial spur with approximately 30–32 distinct teeth along hind margin. Forewings longer than widest part, ratio 3.48:1, widest at apical 1/4 (Figs 10, 12, 19).

Male genitalia. Anal segment with two pairs of spinose processes, upper pair smaller (Figs 24–25). Pygofer quadrate in posterior view (Figs 21–23). Diaphragm broad, transparent (Fig. 23). Aedeagus with ten processes subapically and with irregularity teeth at ventral of apex, constriction and blunt rounded at apex (Fig. 26). Genital styles with inner margin arched and outer margin concave in caudal view, distinctly constricted at apex (Figs 27–28). Suspensorium large and arrow-shaped (Fig. 29).

Female genitalia. Gonocoxa VIII at base of inner margin arched (Fig. 31). Gonapophyses VIII with apex sharp, with ventral margin membranous at half of apex, dorsal




Figures 17–29. *Parasogata binaria* sp. n. male **17** Head and thorax, dorsal view **18** Front **19** Forewing **20** Hindwing **21** Genitalia, lateral view **22** Genitalia, caudal view **23** Diaphragm, caudal view **24** Anal segment, left view **25** Anal segment, caudal view **26** Aedeagus, left view **27** Genital style, caudal view **28** Genital style, left view **29** Suspensorium. Scale bars: 0.5 mm (**17–23**); 0.2 mm (**24–25**); 0.1 mm (**26–29**).

aspect with several small teeth apically (Fig. 32). Gonapophyses IX long, sclerotized, curved basally, narrowing towards apex, with approximately 17 teeth, abruptly reduced and indistinct at apex (Fig. 33). Gonoplasts twisted, long and stripe-shaped (Fig. 34).

Report hosts. None.

Distribution. China (Yunnan).

Etymology. The specific epithet is from the Latin word *binaria* (bipartite), referring to the anal segment with two pairs of processes.



Figures 30–34. *Parasogata binaria* sp. n., female **30** Abdomen, ventral view **31** Gonocoxa VIII **32** Gonapophysis VIII **33** Gonapophysis IX **34** Gonoplac. Scale bars: 0.5 mm (**30**); 0.2 mm (**31–34**).

Remarks. The species is similar to *Sogata dohertyi* (Distent, 1906), but can be distinguished by anal segment with processes cross (not cross in *Sogata dohertyi*), aedeagus with a row processes (without process in *Sogata dohertyi*).

***Parasogata furca* sp. n.**

<http://zoobank.org/85CEB0C9-63CD-42F2-B312-9AD19FAD4F40>


Figs 35–60

Type material. Holotype: ♂, **CHINA, Guizhou:** Wangmo County, Zhexiang (24°97'N, 106°15'E), 7 Jul. 2016, H.-X. Li and L.-J. Yang. Paratypes: 3♂♂, same data as holotype: 2♂♂1♀, **Yunnan:** Yuanjiang County, Donggezhen (23°69'N, 101°82'E), 26 Aug. 2014, Z.-X. Zhou.

Measurements. Body length (from apex of vertex to apex of forewing): male 4.72–4.84 mm (n = 6), female 5.10–5.22 mm (n = 1); forewing length: male 3.61–3.93 mm (n = 6); female 4.42–4.51 mm (n = 1).

Diagnosis. Big-sized species with General color yellow, anal segment with a pair of spinose processes, forked apically (Figs 50–51); aedeagus with eight processes and with many irregularity ventral teeth at subapically (Fig. 52).


Description. *Coloration.* Head yellow. Vertex yellowish white to black (Figs 39, 41). Frons black except middle carinae yellowish white and lateral margin yellowish brown (Figs 40, 42). Clypeus and genae yellow (Figs 40, 42). Rostrum yellowish brown,



Figures 35–42. *Parasogata furca* sp. n. **35, 36** Male habitus (dorsal and lateral views) **37, 38** Female habitus (dorsal and lateral views) **39** Male head and thorax, dorsal view **40** Male front **41** Female head and thorax, dorsal view **42** Female front. Scale bar: 0.5 mm.


with apex brown. Eyes generally yellow to brown (Figs 35–42), ocelli yellowish brown (Figs 36, 38, 40, 42). Antennae yellow (Figs 35–42). Pronotum and mesonotum with carinae yellowish brown (Figs 39, 41). Forewings with veins dark brown (Figs 35–38). Hindwings pale white, veins brown. Legs yellowish white to pale yellowish; tibiae yellow basally, tarsomeres yellowish white (Figs 36, 38). Abdomen yellow (Figs 35–38).

Structure. Head including eyes narrower than pronotum, ratio 0.79:1 (Figs 39, 43). Vertex with anterior margin transverse, lateral carinae with slightly concave, longer than wide at base, ratio 1.22:1, narrower at apex than at base, ratio 0.55:1 (Figs 39, 43). Frons longer in middle line than wide at widest part, ratio 1.83:1, lateral carinae straight (Figs 40, 44). Postclypeus wider at base than frons at apex, slightly longer than wide at base (Figs 41, 46). Antennae cylindrical, basal segment longer than wide, ratio 1.55:1, shorter than second, ratio 0.38:1 (Figs 39–44). Pronotum shorter than vertex, ratio 0.67:1 (Figs 36, 38, 40, 42, 44). Mesonotum longer pronotum and vertex combined, ratio 1.21:1 (Figs 39, 43). Forewings longer than widest part, ratio 3.85:1, widest at apical 1/4 (Figs 35–38, 45). Posttibial spur with approximately 29–32 distinct teeth along hind margin.



Figures 43–55. *Parasogata furca* sp. n., male **43** Head and thorax, dorsal view **44** Front **45** Forewing **46** Hindwing **47** Genitalia, lateral view **48** Genitalia, caudal view **49** Diaphragm, caudal view **50** Anal segment, caudal view **51** Anal segment, left view **52** Aedeagus, left view **53** Genital style, left view **54** Genital style, caudal view **55** Suspensorium. Scale bars: 0.5 mm (**43–49**); 0.2 mm (**50, 51**); 0.1 mm (**52–55**).

Male genitalia. Anal segment with a pair of spinose processes, forked apically (Figs 50–51). Pygofer quadrate in caudal view (Figs 47–49). Diaphragm broad, transparent, dorsal margin arched (Fig. 49). Aedeagus with eight processes and with many irregularity ventral teeth at subapically (Fig. 52). Genital styles with lateral margins arched in caudal view, with two lateral margins almost parallel in profile (Figs 53–54). Suspensorium large and X-shaped, with a process at each lateral margin (Fig. 55).



Figures 56–60. *Parasogata furca* sp. n., female **56** Abdomen, ventral view **57** Gonocoxa VIII **58** Gonapophysis VIII **59** Gonapophysis IX **60** Gonoplac. Scale bars: 0.5 mm (**56**); 0.2 mm (**57–60**).

Female genitalia. Gonocoxa VIII with base of inner margin slightly concave (Fig. 57). Gonapophyses VIII with apex sharp, ventral margin membranous at half of apical, in dorsal margins with several small teeth at half of apical (Fig. 58). Gonapophyses IX long, sclerotized, curved basally and narrowing towards apex, serrated caudad in distal, with approximately 18 teeth, abruptly reduced and indistinct at apex (Fig. 59). Gonoplacs twisted (Fig. 60).

Report hosts. None.

Distribution. China (Guizhou, Yunnan).

Etymology. The specific epithet is from the Latin word *furca* (forked), indicating the anal segment produced lateroapical angles forked.

Remarks. This species is similar to *Parasogata binaria* sp. n., but can be distinguished by the anal segment with a single pair of processes (two pairs of processes in *Parasogata binaria* sp. n.), suspensorium with dorsal margin hunch-up (with dorsal margin concave in *Parasogata binaria* sp. n.).

Acknowledgments

We wish to express our sincere thanks to Prof. M.D. Webb (The Natural History Museum, London, UK) and J.F. Campodonico (University of Chile, Chile) for helpful suggestions on the revision of the early draft of the manuscript. We are grateful to

all collectors of specimens. This work was supported by the National Natural Science Foundation of China (No. 31472033), the program of Science and Technology Innovation Talents Team, Guizhou Province (No.20144001) and the Program of Excellent Innovation Talents, Guizhou Province (No. 20154021).

References

- Bourgoin T (1993) Female genitalia in Hemiptera Fulgoromorpha, morphological and phylogenetic data. *Annales de la Société Entomologique de France (Nouvelle Série)* 29: 225–244.
- Bourgoin T (2018) FLOW (Fulgoromorpha Lists On The Web): a world knowledge base dedicated to Fulgoromorpha. Version 8, updated 20 November 2018. <http://hemiptera-databases.org/flow/> [accessed 25 November 2018]
- Ding JH (2006) *Fauna Sinica. Insecta Vol. 45. Homoptera Delphacidae*. Editorial Committee of Fauna Sinica, Chinese Academy of Science. Science Press, Beijing, 776 pp.
- Dong AP, Qin DZ (2012) A new species in the Oriental delphacid genus *Miranus* Chen & Ding (Hemiptera: Fulgoroidea) from China. *Entomotaxonomia* 34(1): 45–49.
- Hou XH, Yang L, Chen XS (2013) Revision of the genus *Neometopina* (Hemiptera, Fulgoromorpha, Delphacidae) from China. *ZooKeys* 307: 97–104. <https://doi.org/10.3897/zookeys.307.4660>
- Hou XH, Chen XS (2014a) Revision of the planthopper genus *Nycheuma* Fennah (Hemiptera, Fulgoromorpha, Delphacidae). *ZooKeys* 462: 47–57. <https://doi.org/10.3897/zookeys.462.6657>
- Hou XH, Yang L, Chen XS (2014b) Revision of the Oriental genus *Neunkanodes* Yang (Hemiptera, Fulgoromorpha, Delphacidae) with descriptions of two new species. *Zootaxa* 3795(2): 174–180. <https://doi.org/10.11646/zootaxa.3795.2.6>
- Qin DZ (2005) A new genus and a new species of Delphacini (Hemiptera, Fulgoroidea, Delphacidae) from China. *Acta Zootaxonomica Sinica* 30(4): 791–793.
- Qin DZ (2006) Synopsis of two genera of Delphacidae, with descriptions of three new species (Hemiptera, Fulgoroidea). *Acta Zootaxonomica Sinica* 31(2): 392–397.
- Qin DZ, Zhang YL (2006) A new genus and a new species of Delphacini (Hemiptera, Fulgoroidea, Delphacidae) from China. *Zootaxa* 1204: 61–68.
- Qin DZ (2007) Two new species of the Chinese endemic delphacid genus *Neuterthron* Ding (Hemiptera: Fulgoromorpha) from Yunnan and Shaanxi Provinces. *Zootaxa* 1547: 59–64.
- Qin DZ, Chen XS, Lin YF (2008) Revision of the Delphacid genus *Altekon* Fennah (Hemiptera, Fulgoroidea), with description of a new record species from China. *Acta Zootaxonomica Sinica* 33(4): 813–815.
- Qin DZ, Zhang YL (2009a) A new species of *Longtania* Ding from China and redescription of the male genitalia of *Platyptibia ferruginea* Ding (Hemiptera, Fulgoroidea, Delphacidae). *Zootaxa* 1979: 62–68.
- Qin DZ, Chen XD, Men QL (2009b) A new record species of the genus *Megadelphax* Wanger and description of *Eurybregma nigrolineata* Scott (Hemiptera, Fulgoroidea, Delphacidae) from China. *Acta Zootaxonomica Sinica* 34(3): 654–658.

- Qin DZ, Wang SY, Lin YF (2010) Two new record species of Delphacidae (Hemiptera, Fulgoroidea) from China. *Acta Zootaxonomica Sinica* 35(4): 925–929.
- Qin DZ, Chen XD, Lin YF (2011) New record of the genus *Trichodelphax* Vilbaste from China with redescription of *T. Lukjanovitshi* (Hemiptera, Fulgoroidea, Delphacidae). *Acta Zootaxonomica Sinica* 36(1): 202–204.
- Ren FJ, Xie Q, Qin DZ (2014) *Kakuna taibaiensis* sp. n. and a newly recorded species of *Dicranotropis* (Hemiptera, Fulgoroidea, Delphacidae) from China. *ZooKeys* 545: 67–74.
- Ren FJ, Xie Q, Qin DZ (2015) *Mestus cruciatus*, a new delphacid species from southwest China with some remarks on the genus (Hemiptera, Fulgoromorpha, Delphacidae). *ZooKeys* 444: 119–130.
- Yang JT, Yang CT (1986) Delphacidae of Taiwan (I) Asiracinae and the tribe Tropidocephalini (Homoptera: Fulgoroidea). *Taiwan Museum Special Publication* 6: 1–79.