

Pest Alert

Florida Department of Agriculture and Consumer Services,
Division of Plant Industry

Texas Phoenix Palm Decline

Susan Halbert, Susan.Halbert@FreshFromFlorida.com, Taxonomic Entomologist, Florida Department of Agriculture and Consumer Services, Division of Plant Industry

INTRODUCTION: Texas Phoenix palm decline (TPPD) (Figs. 1-7) is a disease of palms that is caused by a phytoplasma (Harrison and Elliot 2009). The phytoplasma is in the taxonomic group of organisms that produce lethal yellows or palm decline in palms (16 Sr-IV group of phytoplasmas). This group of organisms is vast and varied in host range and vector associations. TPPD was noticed initially in Corpus Christi, Texas in 2001 (Harrison et al. 2002) because decline symptoms were more common on *Phoenix* spp. than was expected for known U.S. phytoplasma diseases of palms. This disease now is known to cause decline in *Phoenix sylvestris*, *Phoenix dactylifera*, *Phoenix canariensis*, *Phoenix roebelenii* (Jeyaprakash et al. 2011), *Sabal palmetto* (Harrison and Elliot 2008; Harrison et al. 2009), and *Syagrus romanzoffiana* (Harrison et al. 2008). The entire host range of the pathogen is uncertain at this time. The disease first was noticed in Florida in the Ruskin area (Hillsborough County) in late 2006 and has been observed since then in Hillsborough, Manatee, Sarasota, Pinellas, Polk, Hardee, DeSoto, Highlands, Charlotte, Lee, Lake, Orange, Palm Beach, Indian River, Alachua and Duval counties (Fig. 8).



Figure 1. Sabal palm decline. Photography credit: Susan Halbert, FDACS-DPI.



Figure 2. Sabal palm decline. Note the red color on dying leaves. Photography credit: Susan Halbert, FDACS-DPI.



Figure 3. Sabal palms in various stages of decline. Photography credit: Susan Halbert, FDACS-DPI.



Figure 4. Sabal palms in various stages of decline. Photography credit: Susan Halbert, FDACS-DPI.



Figure 5. Dying *Phoenix sylvestris*. Note that the spear lead has died. This indicates that the apical meristem is dead. Photography credit: Monica Elliott, University of Florida, IFAS, Ft. Lauderdale.




Figure 6. *Phoenix sylvestris* with Texas Phoenix palm decline. Note the dead spear leaf. Photography credit: Monica Elliott, University of Florida, IFAS, Ft. Lauderdale.



Figure 7. Subtle early symptoms of Texas Phoenix palm decline in a nursery setting. Photography credit: Monica Elliott, University of Florida, IFAS, Ft. Lauderdale.




Figure 8. Distribution of Texas Phoenix palm decline in Florida, August 2014. NOTE: Larger map on last page.



DESCRIPTION: The earliest symptom is a discoloration of the lower (oldest) leaves of the palms (Fig. 7). Discoloration begins at the tips of the leaflets. Subsequently, reproductive parts of the plant (if present) will die, resulting in dropping of fruits and flowers. In Phoenix palms, the spear leaf dies after approximately $\frac{1}{4}$ to $\frac{1}{3}$ of the lower canopy has turned brown (Figs. 5, 6). In cabbage palms (*Sabal palmetto*), this may not occur. The disease can be difficult to recognize in the field, because nutritional problems (potassium deficiency, for example) and certain fungal diseases can look similar to the effects of the phytoplasma infection. If it is not the season for fruits and flowers, the diagnostic characteristics involving those parts cannot be used. In taller Phoenix palms, it can be difficult to impossible to see the spear leaf. Typically, infected cabbage palms will have at least the bottom $\frac{1}{3}$ of the canopy dead and bronzed brown, and a much paler dead spear leaf (Fig. 1). A ring of leaves surrounding the spear leaf typically remains green for some time after the spear leaf dies (Fig. 4). Eventually, all the leaves collapse and fall, leaving the stem erect (Fig. 3).

TRANSMISSION: The disease is thought to be transmitted by an insect vector, probably a planthopper (superfamily Fulgoroidea). The species is not known, but there are three species that are found routinely on palms in the areas where the disease is spreading (Halbert *et al.* 2014). One is a large flatid planthopper, *Ormenaria rufifascia* (Walker) (Fig. 9); another is a cixiid planthopper, *Haplaxius crudus* (Van Duzee) [formerly *Myndous crudus* Van Duzee] (Fig. 10); and the third one is a derbid planthopper, *Omolicna joi* Wilson *et al.* (Fig. 11).



Figure 9. Flatid planthoppers, *Ormenaria rufifascia* on *Sabal palmetto* in Florida. Photography credit: Susan Halbert, FDACS-DPI.



Figure 10. Cixiid planthopper on palms, *Haplaxius crudus*, in Florida. Photography credit: David C. Ziesk.



Figure 11. Derbid planthopper, *Omolicna joi*, on palms in Florida. Photography credit: Lyle J. Buss, University of Florida.

The cixiid is the known vector of lethal yellows in South Florida (Howard *et al.* 1983). It occurs throughout the Florida peninsula as far north as Gainesville and can be found during most of the year, but especially in the winter (Fig. 12) (Halbert *et al.* 2014). The derbid is known from Florida north of Broward and Collier counties. Flight activity is in the fall. The flatid is known from all of the Florida peninsula and is most abundant in the late spring.




Figure 12. (Right) Seasonal flight activity of three planthopper species on Florida palms.

REPORTING AND SAMPLING: Homeowners who suspect TPPD should contact their local IFAS County Extension Office. Telephone numbers and addresses can be found at the following website: <http://solutionsforyourlife.ufl.edu/map/index.html>. Samples can be sent to the Fort Lauderdale Research and Education Center. Information on sampling protocol and applicable fees can be found here: <http://frec.ifas.ufl.edu/pdfs/LY-TPPD-Trunk-Sampling.pdf>.

HOSTS: *Phoenix sylvestris*, *Phoenix canariensis*, *Phoenix dactylifera*, *Phoenix roebelenii*, *Syagrus romanzoffiana* and *Sabal palmetto*.


DISTRIBUTION: Texas and Florida, USA.

FLORIDA DISTRIBUTION: Hillsborough, Manatee, Sarasota, Pinellas, Polk, Hardee, DeSoto, Highlands, Orange, Lake, Charlotte, Lee, Palm Beach, Alachua, Broward and Duval counties.

REFERENCES:

- Halbert, S.E., S.W. Wilson, B. Bextine, and S.B. Youngblood. 2014.** Potential planthopper vectors of palm phytoplasmas in Florida with a description of a new species of the genus *Omolocna* (Hemiptera: Fulgoroidea). Florida Entomologist 97: 90-97. <http://journals.fcla.edu/flaent/article/view/83056/79944> [Accessed 2014 August 26].
- Harrison, N.A., M. Womack, and M.L. Carpio. 2002.** Detection and characterization of a lethal yellowing (16SrIV) group phytoplasma in Canary Island date palms affected by lethal decline in Texas. Plant Disease 86: 676-681.
- Harrison, N.A. and M.L. Elliott. 2008.** A Lethal Phytoplasma Disease of *Sabal palmetto* on Florida's Central West Coast. University of Florida. http://frec.ifas.ufl.edu/palm_prod/pdfs/Sabal-palmetto-Infected-with-Phytoplasma-in-Florida.pdf [Accessed 2014 July 23].
- Harrison, N.A. and M.L. Elliott. 2009.** Texas Phoenix palm decline. University of Florida, IFAS. http://manatee.ifas.ufl.edu/comm-hort/pdf/palms/Texas_Phoenix_Palm_Decline_2009.pdf [Accessed 2014 July 23].
- Harrison, N.A., E.E. Helmick, and M.L. Elliott. 2008.** Lethal yellowing-type diseases of palms associated with phytoplasmas newly identified in Florida, USA. Annals of Applied Biology 153: 85-94. <http://onlinelibrary.wiley.com/doi/10.1111/j.1744-7348.2008.00240.x/> [Accessed 2014 July 23].
- Harrison, N.A., E.E. Helmick, and M.L. Elliot. 2009.** First report of a phytoplasma-associated lethal decline for *Sabal palmetto* in Florida, USA. Plant Pathology 58: 792.
- Howard, F.W., R.C. Norris, and D.L. Thomas. 1983.** Evidence of transmission of palm lethal yellowing agent by a planthopper, *Myndus crudus* (Homoptera: Cixiidae). Tropical Agriculture (Trinidad) 60: 168-171.
- Jeyaprakash, A., B.D. Sutton, S.E. Halbert, and T.S. Schubert. 2011.** High-fidelity PCR facilitates detection and identification of a Texas Phoenix palm phytoplasma strain from pigmy date palm, *Phoenix roebelenii* in Florida. Plant Disease 95: 1475.

Texas Phoenix Palm Decline (TPPD) in Florida



FDACS, DPI, CAPS
G. Gardner, L. Whilby, W. Dixon
Map for illustrative purposes only



GCS North American 1983
PCS Albers Equal Area Conic
Data Source: CAPS
Date: 10/8/2014