



Overexpression of *Arabidopsis* Nucleotide-Binding and Leucine-Rich Repeat Genes *RPS2* and *RPM1(D505V)* Confers Broad-Spectrum Disease Resistance in Rice

Zhaowu Li, Jianzhong Huang, Zhangying Wang, Fen Meng, Siyuan Zhang, Xiaoqiu Wu, Zhihong Zhang and Zhiyong Gao*

State Key Laboratory of Hybrid Rice, Key Laboratory for Research and Utilization of Heterosis in Indica Rice of Ministry of Agriculture, College of Life Sciences, Wuhan University, Wuhan, China

OPEN ACCESS

Edited by:

Victor Flors,
University of Jaume I, Spain

Reviewed by:

Jordi Gamir,
Estación Experimental del Zaidín
(EEZ), Spain
Marc Tad Nishimura,
Colorado State University,
United States

*Correspondence:

Zhiyong Gao
zygao@whu.edu.cn

Specialty section:

This article was submitted to
Plant Microbe Interactions,
a section of the journal
Frontiers in Plant Science

Received: 19 January 2019

Accepted: 19 March 2019

Published: 05 April 2019

Citation:

Li Z, Huang J, Wang Z, Meng F, Zhang S, Wu X, Zhang Z and Gao Z (2019) Overexpression of *Arabidopsis* Nucleotide-Binding and Leucine-Rich Repeat Genes *RPS2* and *RPM1(D505V)* Confers Broad-Spectrum Disease Resistance in Rice. *Front. Plant Sci.* 10:417. doi: 10.3389/fpls.2019.00417

The nucleotide-binding domain leucine-rich repeat (NLR) immune receptors play important roles in innate plant immunity. The activation of NLRs is specifically induced by their cognate effectors released from pathogens. Autoactive NLRs are expected to confer broad-spectrum resistance because they do not need cognate effectors to activate their immune responses. In this study, we demonstrated that the *NLR* genes *RPS2* and *RPM1(D505V)* from *Arabidopsis* were autoactive in *Oryza sativa* and conferred broad-spectrum resistance to fungal pathogen *Magnaporthe oryzae*, bacterial pathogen *Xanthomonas oryzae pv. oryzae* (*Xoo*), and pest brown planthopper (BPH, *Nilaparvata lugens* Stål). These results revealed that interfamily transfer of dicot NLRs to monocot species could be functional. The transgenic plants displayed early and strong induction of reactive oxygen species (ROS), callose deposition, and expression of defense-related genes after challenged with *M. oryzae*. The transcriptome analysis showed that the expressions of some defense-related genes were primed to adapt the transformed autoactive NLRs in the transgenic plants. This study indicates that autoactive NLRs are a promising resource for breeding crops with broad-spectrum resistance and provides new insights for engineering disease resistance.

Keywords: *AtRPS2*, *AtRPM1*, rice, *Magnaporthe oryzae*, *Xanthomonas oryzae*, *Nilaparvata lugens*

INTRODUCTION

Plants are constantly threatened by a wide range of pathogens in nature, and plants have evolved a complicated and multifaceted innate immune system to combat pathogen attacks. The first layer of the plant immune system is mediated with pattern recognition receptors (PRRs). PRRs perceive the pathogen-associated molecular patterns (PAMPs), the highly conserved molecular signatures across a wide range of pathogens, and activate the PAMP-triggered immunity (PTI). PTI contributes a basal immunity during pathogen infection (Jones and Dangl, 2006; Couto and Zipfel, 2016;

Zipfel and Oldroyd, 2017). In most cases, PTI is sufficient to prevent pathogen invasion and disease development. Successful pathogens deploy virulent proteins which are named effectors to suppress PTI, and correspondingly, plants have evolved immune receptors to recognize the effectors and initiate rapid and strong defense responses, named effector-triggered immunity (ETI) (Bent and Mackey, 2007; Tsuda and Katagiri, 2010; Spoel and Dong, 2012). PTI is conserved in different plant species and acts as a basal defense, and ETI provides a race-specific resistance and amplifies PTI signaling to induce robust resistance responses. Therefore, PTI and ETI form a two-tier defense system to protect plants against various pathogens (Pieterse et al., 2014; Cui et al., 2015).

Most of the effector-recognizing immune receptors have a nucleotide-binding domain and a leucine-rich repeat domain, and these receptors are called NLR receptors. NLR receptors either directly or indirectly recognize effectors (Cesari et al., 2014). Only a few NLR receptors directly bind to their corresponding effectors. More NLRs recognize effectors *via* an indirect recognition. One indirect recognition model is the “guardee” strategy. RIN4 is guarded by NLR proteins RPM1 and RPS2 in *Arabidopsis* and is targeted by effectors AvrRpm1, AvrB, and AvrRpt2. When AvrB and AvrRpm1 are delivered into the plant cell, they induce the phosphorylation of RIN4, and the phosphorylated RIN4 activates RPM1 signaling (Mackey et al., 2003). RPS2 interacts with RIN4, and its activity is inhibited by RIN4. AvrRpt2 can cleave RIN4 to stimulate the activity of RPS2 (Mackey et al., 2002; Axtell and Staskawicz, 2003).

NLR genes are important resources for breeding resistant plants. Several NLR genes have been cloned and employed in practical applications, and obtained great economic and social benefits (Park et al., 2012; Wang et al., 2016). The genome of the dicotyledonous plant *Arabidopsis thaliana* contains about 150 NLR genes that encode CC-NB-LRR and TIR-NB-LRR two types of NLR proteins, and the genome of the monocotyledonous plant *Oryza sativa* contains about 480 NLR genes that encode only CC-NB-LRR proteins (Yang et al., 2006). Although the large amount of NLR genes has been found from genomes of different plant species, NLRs with known activation mechanisms are quite limited. Because NLRs specially recognize their cognate effectors to activate the immune responses, lacking the knowledge on their cognate effectors restricts the successful applications of those NLRs. Autoactive NLRs do not need effectors to activate their immune responses, and are potential effective resources for breeding broad-spectrum resistant plants. There are two ways to obtain autoactive NLRs. RPS2 is inhibited by RIN4 in *Arabidopsis*, it is autoactive in *rin4* knock-out mutant or transiently expressed in *Nicotiana benthamiana* (Day et al., 2005). Interfamily transfer of NLR genes such as RPS2 to the plant species without their suppressive genes such as RIN4 can obtain autoactive NLRs. NLRs have conserved domains to regulate the switch between inactive and active states. Mutations in certain domains can lead to autoactive NLRs. Activation of RPM1 requires phosphorylated RIN4, and it does not autoactive in *rin4* knock-out mutant. However, RPM1(D505V) is autoactive in *rin4* knock out mutant (Gao et al., 2011).

Rice (*Oryza sativa* L.) is an important staple food crop that feeds more than half of the world's population. However, its production is threatened by many pathogens and insect pests (Ke et al., 2017). Rice blast disease, which is caused by the hemibiotroph ascomycete pathogen *M. oryzae*, is one of the most destructive diseases. *M. oryzae* infects rice stems, nodes, leaves, and panicles, resulting in 10–30% yield losses in many rice growing areas (Wu et al., 2015; Bundo and Coca, 2016; Wang B.H. et al., 2017). Bacterial blight is another devastating disease of rice, which is caused by *Xanthomonas oryzae* *pv.* *oryzae* (Xoo) (Gu et al., 2009). Brown planthopper (BPH) is a rice-specific herbivore and is one of the most destructive insects of rice (Cheng et al., 2013; Liu et al., 2015). *M. oryzae*, Xoo, and BPH cause massive yield decrease and are a huge threat to food security (Zhang and Wang, 2013; Jing et al., 2017). Therefore, it is an urgent issue to develop durable and broad-spectrum of rice cultivars against diseases and insects.

Some NLR proteins from both *Arabidopsis* and rice could induce programmed cell death (PCD) when they were transiently expressed in the leaves of *N. benthamiana* (Wulff et al., 2011). The results suggest that NLRs share conserved signal pathways among different species, and it is possible that NLR proteins from *Arabidopsis* can be used in rice to confer resistance. Interfamily transfer of NLR genes was seldom reported (Maekawa et al., 2012; Narusaka et al., 2013). We transformed the *Arabidopsis* NLR genes *AtRPS2* and *AtRPM1(D505V)* into the *Japonica* rice cultivar Nipponbare under the control of *Zea mays ubiquitin* promoter. The results showed that these transgenic rice plants increased resistance against fungal pathogen *M. oryzae*, bacterial pathogen Xoo and insect pest BPH. Evidence is provided to show that the dicot NLR genes can be functional and confer the broad-spectrum disease and pest resistance in monocot species. Our observations revealed that interfamily transfer NLR genes will broaden the resources for breeding multi-resistance crops.

MATERIALS AND METHODS

Plant Materials and Growth Conditions

The rice used in this study was *Oryza sativa* ssp. *Japonica* var. Nipponbare. All of the rice plants were grown with soil in a greenhouse or in the experimental fields at Wuhan. The greenhouse was set with a cycle of 14 h light at 28°C/10 h dark at 25°C, and the relative humidity was 50–60%.

Plasmid Construction and Rice Transformation

AtRPS2 and *AtRPM1(D505V)* were cloned into a modified pCAMBIA1380 vector with an *ubiquitin* promoter. The constructs were transformed into the *Agrobacterium tumefaciens* strain EHA105. The *Agrobacterium*-mediated transformation and regeneration of transgenic plants were carried out as previously described (Chen et al., 2015). With hygromycin selection and segregation analysis, the single insertion stable transgenic plants and further progeny were used for subsequent experimental studies.

Rice Blast Inoculations and Disease Resistance Evaluations

The *M. oryzae* isolates used in this study were grown on oatmeal agar for 1 week at 28°C and then transferred to blue light for 2 days to enhance sporulation. Fungal spores were suspended in 0.02% Tween-20 at a concentration of 5×10^5 /mL. Three to four-leaf stage rice plants were used for spray inoculation, and the inoculated plants were incubated under a dark moist condition (28°C, 99% relative humidity) for 24 h and then under a 12 h light/dark cycle with 80% humidity. Disease severity was evaluated at 7 days after inoculation by counting the lesion numbers, and measuring the lengths of the three largest lesions per leaf. For panicle blast inoculation, the upper-middle part of panicles was injected with 1 mL of conidial suspension at a concentration of 5×10^5 /mL, and panicle blast resistance was assessed at 14 days after inoculation by measuring the proportion of diseased panicles.

Punch Inoculation and Root Infection Assay

For punch inoculation, detached rice leaves were lightly wounded with an ear punch, and 8 μ L of spore suspension (5×10^5 /mL) was added to the wound sites. The inoculated leaves were incubated in a Petri dish contained sterile water with 0.1% 6-Benzylaminopurine (6-BA) to keep moisture. Lesions were measured at 7 days after inoculation. The fungal biomass in the infected rice leaf was quantified using the threshold cycle value (C_T) of *M. oryzae* *Pot2* DNA against the C_T of rice genomic *actin* DNA (Park et al., 2008).

Root infection assays were carried out with moist vermiculite. The mycelial plugs were put into a 50 mL centrifuge tube filled with 25 cm of autoclaved moist vermiculite and covered with 3 mL of vermiculite. Rice seeds were surface-sterilized for 10 min and washed six times with autoclaved water. Three sterilized seeds were placed on the top of the inoculated vermiculite and covered with another 3 mL of vermiculite. The tubes were sealed with Parafilm to retain moisture and incubated for 3 weeks (22°C, 16 h light/8 h dark photoperiod) to examine lesions.

Rice Sheath Infection Assay

To observe the infection process of *M. oryzae*, the spore suspensions of GC1-4 was inoculated on 6 cm-long leaf sheaths with the concentration of 5×10^5 /mL, and then the inoculated leaf sheaths were incubated under the same condition as spraying assay. The leaf sheaths were observed with a laser scanning confocal microscope (Leica SP8-X) at 24 and 48 hpi.

Xoo Inoculation and Bacterial Blight Disease Assay

Xoo strain PXO99 was grown on PSA medium (300 g/L Potato, 5 g/L peptone, 15 g/L sucrose, 0.5 g/L $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$, 2 g/L $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$, 20 g/L agar, pH 6.8–7.0) for 2 days, then bacteria were suspended in sterile water and diluted to $\text{OD}_{600} = 0.5$. The three youngest fully expanded leaves of the 6 weeks old plants were inoculated with PXO99 by the

leaf-clipping method. Lesion length and bacterial growth were measured at 14 dpi as previously described (Gu et al., 2009).

BPH Resistance Evaluations

The BPH insects used for resistance evaluations were reared on the susceptible rice cultivar TN1. For honeydew excretion and BPH weight gain assay, BPH nymphs were weighed and released to a pre-weighed Parafilm sachet and tied to the rice leaf sheath. The honeydew of each Parafilm sachet and the insects were weighed after 48 h. The weight change of BPH was recorded as the weight gain of BPH, and the weight change of the sachet was recorded as the honeydew excretion.

To assess the survivals and durations of transgenic plants and Nipponbare after BPH infestation, 10 rice seedlings at 4 weeks old were infested with BPH at a rate of eight insects per seedling. The transgenic plants were examined and photographed when all Nipponbare seedlings were dead.

In host selection test, two 30 days old plants were grown in one plastic bucket. One plant is Nipponbare and the other is a transgenic plant. Twenty BPH nymphs were released in the bucket, and the number of BPH nymphs that settled on each plant was recorded at 3, 6, 12, 24, and 48 h after insect release.

RNA Extraction and qRT-PCR Analysis

The total RNA was extracted using Trizol reagent (TaKaRa, Dalian, China), and cDNA synthesis was performed according to the manufacturer's protocol (Thermo Fisher Scientific, Lithuania). qRT-PCR was carried out in a Bio-Rad CFX96 real-time system with SYBR green PCR kit. The *OsActin* gene was used as the internal control for data normalization. The primers used for qRT-PCR are listed in **Supplementary Table S6**.

RNA-Seq Analysis

About 2 cm of leaf strips from 4 week-old Nipponbare and transgenic plants were collected, and total RNA of the samples was extracted by TRIzol reagent (TaKaRa, Dalian, China). The RNA quality was quantified using an Agilent 2100 bioanalyzer (Agilent, Santa Clara, CA, United States), and then sequenced at an Illumina HiSeqTM 2500 platform. The adaptors and low-quality sequences were removed from the raw data. Differentially expressed genes (DEGs) were identified using the criteria of fold change > 2 and $p < 0.05$.

DAB Staining

Hydrogen peroxide was monitored by DAB staining as previously described (Shimono et al., 2012). Pieces of leaf sheaths (3 cm) were immersed in 1 mg/mL DAB solution (pH 3.8), followed by vacuum filtration for 10 min and incubation at room temperature for 10 h in the dark. Samples were bleached by boiling in ethanol for 15 min, and images were observed with a microscope (Olympus BX51, Tokyo, Japan).

Callose Deposition Analysis

For examining callose deposition, rice seedlings were inoculated with *M. oryzae* GC1-4 for 24 and 48 h and washed in an ethanol:acetic acid (3:1 v/v) solution for 5 h. The seedlings were

incubated in 70% ethanol for 2 h, in 50% ethanol for 2 h, and in water overnight. After rinsing three times with water, the seedlings were incubated with 10% NaOH for 1 h to make all tissues transparent. After being washed three times with water, the seedlings were stained with 0.01% aniline blue for 4 h (Hu et al., 2017). Representative images of callose deposition were observed using a laser scanning confocal microscope (Leica SP8-X) under the UV channel.

Assessment of Agronomic Performance

The Nipponbare and transgenic plants were grown in a standard field in Wuhan, Hubei, China. When plants were harvested, the middle six plants in the central row of each plot were collected for agronomic trait measurement. The measured agronomic traits include: plant height, number of panicles per plant, filling rate per plant, 1000-grain weight (grams).

RESULTS

Overexpression of *AtRPS2* and *AtRPM1(D505V)* Confers Resistance Against *M. oryzae* in Transgenic Rice

AtRPS2 is autoactive in plants without RIN4. *AtRPM1* is not autoactive without RIN4 because the activation of *AtRPM1* needs the phosphorylated RIN4. However, *AtRPM1(D505V)* is autoactive without RIN4. Both *AtRPS2* and *AtRPM1(D505V)* can cause HR when transiently expressed in *N. benthamiana*. Since the autoactive NLRs usually result in broad-spectrum disease resistance, *AtRPS2* and *AtRPM1(D505V)* transgenic rice plants under the control of *ZmUbi* promoter were generated to determine the function of *AtRPS2* and *AtRPM1(D505V)* in rice. We obtained 51 independent *AtRPS2* transgenic T₁ lines and 54 independent *AtRPM1(D505V)* lines. Among the T₁ lines, 13 *AtRPS2* transgenic lines and 15 *RPM1(D505V)* transgenic lines grew so tiny that no seeds formed. Other lines showed various dwarf growth or normal growth. The transgenes were fused with HA tag, so the protein levels of *RPS2* and *RPM1(D505V)* in the transgenic lines could be determined. The detectable protein expressions of *RPS2* or *RPM1(D505V)* were found in the leaves of the dead T₁ lines, while no detectable protein expressions in the survived T₁ lines. The expression level of the genes in those transgenic lines were measured by qRT-PCR. The significant high lethal ratio and the dwarf growth of the transgenic plants indicate the autoactivation of *AtRPS2* and *AtRPM1(D505V)* in rice. Seven independent lines with stable single insertion and normal growth were selected for *AtRPS2* and *AtRPM1(D505V)* transgenic plants, respectively (Supplementary Tables S1, S2). Two lines per construct, *Ubi::AtRPS2*-2, -9 and *Ubi::AtRPM1(D505V)*-D, -E were chosen for experiments. They are lines with the highest expression level or the lowest expression level of transgenic genes based on qRT-PCR analyses (Supplementary Figures S1A,B). These transgenic lines and the transgenic background plant Nipponbare were used to evaluate the disease resistance against blast fungus. Five different *M. oryzae* isolates GC1-4, YA2-3, TF11-3, ZX2-6, and HN-8, which were collected from different


cities or provinces of China and highly virulent to Nipponbare, were selected. Both *AtRPS2* and *AtRPM1(D505V)* transgenic plants showed obviously less disease severity compared to Nipponbare (Figure 1A). The lesion lengths and lesion numbers were dramatically reduced in the transgenic plants with these five isolates (Figures 1B,C). Similar results were obtained in detached leaf assays. The transgenic plants developed smaller lesions, and the lesion lengths on the transgenic plants were reduced to 40–50% compared to Nipponbare. Fungal DNA quantification indicated that the *AtRPS2* and *AtRPM1(D505V)* transgenic plants only harbored ~35% *M. oryzae* compared to Nipponbare (Supplementary Figure S2).

The panicle blast resistance was tested with the transgenic plants. The whitish dead panicles were observed in Nipponbare at 14 days post-inoculation (dpi), while none of the *AtRPS2* and *AtRPM1(D505V)* transgenic plants exhibited disease symptoms (Figures 2A,B). The semi-quantitative RT-PCR assay indicated that the fungal biomass in Nipponbare was much higher than those in the transgenic plants (Figure 2C). In the root infection assay, the roots of Nipponbare showed obvious necrotic symptoms, while the roots of the transgenic plants displayed less necrotic symptoms (Figure 2D). These results suggest that *AtRPS2* and *AtRPM1(D505V)* induce strong blast resistance in leaves, panicles, and roots of rice.

Leaf sheaths were inoculated with conidial suspensions of *M. oryzae* isolate GC1-4 to microscopically monitor the growth of the pathogen in plants. High levels of invasive hyphae and extensions to neighboring cells were observed in Nipponbare at 24 h post-inoculation (hpi), while only a small percentage of spores developed invasive hyphae and some plant cells displayed cell death in the transgenic plants. Almost all the penetration sites showed high levels of invasive hyphae in Nipponbare at 48 hpi, while the infection sites generated severe HR death response in the transgenic plants (Figure 3).

Overexpression of *AtRPS2* and *AtRPM1(D505V)* Confers Enhanced Defense Responses in Transgenic Rice


Reactive oxygen species (ROS), such as H₂O₂, are often correlated with plant defense responses (Liu et al., 2016; Yuan et al., 2017). In this study, 3,3-diamino-benzidine (DAB) was used to detect H₂O₂ in the leaves of the plants. While the leaves of Nipponbare and the transgenic plants had no H₂O₂ staining without the inoculation of *M. oryzae*, the transgenic plants generated much higher amounts of H₂O₂ than Nipponbare at 24 and 48 hpi (Figures 4A,B). We analyzed the activities of peroxidases (POD), catalases (CAT), and superoxide dismutases (SOD) in the plants. Before pathogen inoculation, the activities of these enzymes had no obvious differences between Nipponbare and the transgenic plants. Inoculation of *M. oryzae* increased the activities of the three enzymes, and the transgenic plants had higher POD and CAT activities than Nipponbare at 24 hpi (Figures 4C,D). The SOD activities in transgenic plants were higher than Nipponbare but did not display significant differences after *M. oryzae* inoculation (Figure 4E). These results indicate that expression of *AtRPS2* and *AtRPM1(D505V)* in



rice can increase H_2O_2 accumulation corresponding to the inoculation of *M. oryzae*.

Callose deposition is another important defense response and can prevent pathogens from penetrating into the plant cells (Shangguan et al., 2018). The callose depositions on the leaves of rice were stained with aniline blue at 0, 24, and 48 hpi, and the transgenic plants displayed more callose depositions than Nipponbare (Figures 5A,B). We determined the expression patterns of callose biosynthesis genes, *OsGSL5*, *OsGSL6*, and *OsGSL7*. *M. oryzae* induced the highest expressions of *OsGSL5* and *OsGSL6* in *AtRPS2* and *AtRPM1* transgenic plants at 12 hpi, but the two genes did not have significant expression changes in Nipponbare (Figures 5C,D). *OsGSL7* displayed similar expression patterns in all transgenic plants and Nipponbare (Figure 5E). We also determined the expression patterns of callose hydrolase genes, *OsGNS6*, *OsGNS8*, and


OsGNS9. *OsGNS6* was induced by the inoculation of *M. oryzae* in the transgenic plants and Nipponbare, but the transgenic plants had lower expression levels than Nipponbare at 12 and 24 hpi (Figure 5F). *OsGNS8* displayed similar expression patterns in the transgenic plants and Nipponbare, but the transgenic plants significantly reduced the gene expression at 12 hpi (Figure 5G). *OsGNS9* displayed different expression patterns between the transgenic plants and Nipponbare. The expression of *OsGNS9* was constantly downregulated in the transgenic plants after *M. oryzae* inoculation, while the expression of *OsGNS9* was dramatically upregulated in Nipponbare at 12 and 24 hpi (Figure 5H). Therefore, the early induced *OsGSL5* and *OsGSL6*, as well as the constantly suppressed *OsGNS9* promote the synthesis of callose and reduce the hydrolysis of the callose as well, resulting in quick and strong callose depositions in the transgenic plants.



The *AtRPS2* and *AtRPM1(D505V)* Transgenic Plants Are Primed for Defense Responses

The transgenic plants did not display constitutively higher contents of H_2O_2 and callose deposition, suggesting that

AtRPS2 and *AtRPM1(D505V)* did not constitutively activate the plant immunity in rice. The early and strong inductions of H_2O_2 and callose deposition indicate that the transgenic plants are primed for defense responses. We analyzed the expression of some defense-responsive genes in the transgenic plants and Nipponbare. Upon *M. oryzae* inoculation, the



transgenic plants displayed higher expression levels of *PR2* than Nipponbare at 12 and 24 hpi (**Figure 6A**). *PR3* and *PR5* were induced to higher levels in the transgenic plants than in Nipponbare at 12 hpi, and the two genes displayed obviously constitutive expression in the *AtRPM1(D505V)* transgenic plants (**Figures 6B,C**). The expression level of *WRKY45* in Nipponbare was not induced upon *M. oryzae* challenge, while its expression was induced to higher levels in the transgenic plants at 12 and 24 hpi (**Figure 6D**). These results suggest that the expression of *AtRPS2* and *AtRPM1(D505V)* leads to the quick and strong induction of the defense-related genes upon pathogen infection.

We compared the global gene expressions between the transgenic plants and Nipponbare without pathogen inoculation. Compared to Nipponbare, the *AtRPS2* transgenic plants and the *AtRPM1(D505V)* transgenic plants had 106 and 245 upregulated genes, and 86 and 148 downregulated genes, respectively (**Supplementary Tables S3, S4**). We analyzed these DEGs using the gene ontology (GO) analyses. In *AtRPS2* transgenic plants, the expression of genes involved in “metabolic process,” “defense response,” and “zeatin activity” showed significant enrichment, and in *AtRPM1(D505V)* transgenic plants, the expression of genes involved in “metal ion homeostasis,”

“defense response,” and “metal ion binding” showed significant enrichment (**Supplementary Figures S3, S4**). Because *AtRPS2* and *AtRPM1(D505V)* transgenic plants displayed similar disease resistance, we selected the commonly regulated genes in both transgenic plants. 97 genes were commonly regulated in both transgenic plants, and 50 genes had annotations (**Supplementary Table S5**). Most of the genes are defense-related genes, including eight putative *NLR* genes and four *receptor-like kinase* genes. These results indicate that expression of *AtRPS2* and *AtRPM1(D505V)* causes the plant to be primed for disease resistance.

Overexpression of *AtRPS2* and *AtRPM1(D505V)* Confers Resistance to Bacterial Pathogen *Xoo* and BPH in Transgenic Rice

Because *AtRPS2* and *AtRPM1(D505V)* enhanced resistance to blast disease and could trigger stronger defense responses, we tested whether the *AtRPS2* and *AtRPM1(D505V)* transgenic plants showed enhanced resistance to bacterial blight disease, which was caused by *Xoo*. The transgenic plants showed remarkably shorter lesion lengths caused by *Xoo* strain






FIGURE 4 | *AtRPS2* and *AtRPM1*(D505V) Trigger Strong ROS Induction During *M. oryzae* Infection. **(A)** DAB staining of the transgenic plants and Nipponbare at 24 and 48 hpi. There were many tawny shading cells in the transgenic plants, but in Nipponbare, the infected cells mostly exhibited light yellow color. Red arrows indicate dead plant cells. Scale bars = 50 μ m. **(B)** H_2O_2 contents after inoculation with *M. oryzae* isolate GC1-4 in Nipponbare and transgenic plants. FW, fresh weight. **(C-E)** The enzymatic activities of POD, CAT, and SOD were assayed after *M. oryzae* inoculation. Data are shown as means \pm SD ($n = 3$). Asterisks denote significant differences (one way ANOVA, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). Similar results were obtained in three independent experiments.



PXO99 compared to Nipponbare at 14 dpi (Figure 7A). The average lesion length of the transgenic plants was 3.5 cm, while the average lesion length of Nipponbare was 8.2 cm

(Figure 7B). Similarly, the bacterial growth rates in the transgenic plants were significantly less compared to Nipponbare (Figure 7C). These data suggest that overexpression of *AtRPS2*



and *AtRPM1(D505V)* in rice enhanced resistance to bacterial pathogen *Xoo*.

We also determined whether the transgenic plants developed enhanced resistance to the phloem-sucking insect BPH. We measured the weight gains of BPH and the honeydew excretions of the insects after feeding on the plants 2 days. The insects gained less weight and produced less honeydew on the transgenic plants (Figures 8A,B). Host choice test showed that Nipponbare was more attractive to BPH after 12–48 h release (Figure 8C). Furthermore, the transgenic plants survived from the BPH infection when all the Nipponbare plants had been killed by the insects (Figure 8D). BPH weight change and honeydew excretion reflect the antibiosis of the plants to the insects, and the host choice reflects the antixenosis of the plants to the insects. The results suggest that *AtRPS2* and *AtRPM1(D505V)* also confer the enhanced resistance to BPH.


AtRPS2 and *AtRPM1(D505V)* Transgenic Plants Exhibit Fitness Costs

The constitutive expression of *AtRPS2* or *AtRPM1(D505V)* conferred the enhanced disease resistance, but it also brought in the fitness cost. We grew the transgenic plants and Nipponbare in the experimental fields to examine the agronomic traits. At mature stage, the *AtRPS2* and *AtRPM1(D505V)* transgenic plants

were shorter than Nipponbare (Figures 9A,B). In addition, the transgenic plants possessed more infertile pollen (Figure 9C). The yield agronomic traits including plant height, tiller numbers per plant, seed setting rates and 1000-grain weights were measured. There was no difference between the transgenic plants and Nipponbare in 1000-grain weight, while the plant height, tiller numbers per plant, seed setting rates were significantly lower in the transgenic plants (Figures 9D–G).

DISCUSSION

The transfer of the PRR genes and resistance-related genes from *Arabidopsis* to other plants has been proposed as a strategy for achieving broad-spectrum resistance. The *Arabidopsis* PRR EFR recognizes the bacterial elongation factor Tu and elf18. Expression of *AtEFR* in tomatoes and *Nicotiana benthamiana* conferred resistance to various bacterial pathogens (Lacombe et al., 2010), and expression of *AtEFR* in rice and wheat enhanced resistance to bacterial pathogen *Xoo* and *Pseudomonas syringae* pv. *Oryzae*, respectively (Lu et al., 2015; Schoonbeek et al., 2015). The expression of *AtRPW8.1* in rice conferred rice blast and bacterial blight resistance (Li et al., 2018). NPR1 is a positive regulator of the systemic acquired resistance (SAR) in *Arabidopsis* (Fu and Dong, 2013). Expression



of *AtNPR1* in rice enhanced resistance to the fungal pathogen *Rhizoctonia solani* and bacterial pathogen *Xoo* (Fitzgerald et al., 2004; Molla et al., 2016). Therefore the successful usage of *Arabidopsis* PRRs and resistance-related genes in other species indicates that the immunity pathways among different plants are relatively conserved.

Different from previous research, we transformed the *Arabidopsis* autoactive NLR genes *AtRPS2* and *AtRPM1(D505V)*

into the rice cultivar Nipponbare. The autoactivity of the two genes in rice can be proven by the tiny growth appearing on the transgenic plants with high expressions of the two genes. Because *Agrobacterium*-mediated gene transfer randomly integrates the transgenic gene to the rice genome, some transgenic plants with low expression of the two genes grew normally. The cognate effector of *AtRPS2* is *AvrRpt2* from *Erwinia amylovora*, and the cognate effectors of *AtRPM1* are *AvrRpm1* and *AvrB* from




FIGURE 8 | *AtRPS2* and *AtRPM1(D505V)* Confer Resistance to BPH in Transgenic Plants. **(A,B)** Weight gain and Honeydew excretion of BPH on transgenic plants and Nipponbare after feeding 2 days. Data were shown as means \pm SD ($n = 20$). **(C)** Host choice test of BPH on transgenic plants and Nipponbare plants. Data are shown as means \pm SD ($n = 10$). Asterisks denote significant differences (one way ANOVA, $*P < 0.05$, $**P < 0.01$). Similar results were obtained in three independent experiments. **(D)** Representative images of the transgenic plants and Nipponbare plants after BPH infestation. Note that the Nipponbare seedlings were dead thoroughly, but the transgenic plants still survived.




FIGURE 9 | Phenotypes and agronomic traits of the *AtRPS2*, *AtRPM1(D505V)* transgenic plants and Nipponbare plants. **(A,B)** The morphological phenotypes of *AtRPS2*, *AtRPM1(D505V)* transgenic plants and Nipponbare. **(C)** Comparison of pollen fertility between the transgenic plants and Nipponbare plants. The heavily stained pollen is fertile and the lightly stained pollen is infertile. **(D–G)** Plant height (cm), tiller numbers per plant, seed setting rate, and 1000-grain weight. Data are shown as means \pm SD ($n = 6$). Asterisks represent significant differences (one way ANOVA, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

Pseudomonas syringae. The two NLRs should specially fight against pathogens contained the cognate effectors, however, the autoactive *AtRPS2* and *AtRPM1(D505V)*, which did not need their cognate effectors to activate their function in rice, conferred enhanced resistance to fungal pathogen *M. oryzae*, bacterial pathogen *Xoo*, and insect pest BPH (Figures 1, 7, 8). In addition, the low expression transgenic lines display strong resistance, which means that we could choose the low expression autoactive transgenic lines to minimize fitness cost. Our study suggests that NLRs share conserved signaling pathways among different species, and autoactive NLRs confer plants broad-spectrum resistances.

Because *AtRPS2* and *AtRPM1(D505V)* were constitutively expressed in rice, we previously expected the transgenic plants to have constitutive defense responses, such as high H₂O₂ content and callose depositions. However, both *AtRPS2* and *AtRPM1(D505V)* transgenic plants did not display detectable higher H₂O₂ content or more callose depositions compared to the wild type plant Nipponbare (Figures 4, 5). The early and strong defense responses were induced in transgenic plants by the inoculation of *M. oryzae*, which ultimately suppressed the spread of invasive hyphae.

Compared with the global gene expressions in Nipponbare, two putative receptor-like kinase genes *OS11G0274100* (similar to stem rust resistance protein) and *OS11G0470500* were upregulated 8- and 3-fold, respectively, in the *AtRPM1(D505V)* transgenic plant, and two putative receptor-like kinase genes *OS10G0162836* (Wall-associated receptor kinase 1, WAK1) and *OS11G0226201* (similar to HSL2) were downregulated 4- and 33-fold, respectively (Supplementary Table S5). WAK1 is a candidate receptor of oligogalacturonides (OGs), components released from the plant cell wall (Brutus et al., 2010). HSL2 reduces leaf pectin content and leaf robustness, and HSL2 knockdown lines increase *Arabidopsis* resistance to *Pst* DC3000 (Wang X. et al., 2017). These data suggest that the cell wall structure and some PRRs associated with cell wall or on the plasma membrane have constitutively changed in the transgenic plants. Both *AtRPS2* and *AtRPM1(D505V)* transgenic plants displayed the same resistance, and it was consistent with the data that they share many commonly regulated genes.

The fitness costs resulting from autoactive NLRs hurdle the practical application of such NLRs. We used the *Zea mays ubiquitin* promoter to control the gene expression of *AtRPS2* or *AtRPM1(D505V)* in rice. The constitutive and high expression promoter was used to produce sufficient protein levels in rice, thus determining the autoactivity of the NLRs in the transgenic plants and subsequent resistance analysis. We obtained a series of transgenic plants with phenotypes ranging from dwarf plants to wild-type-like plants. The random insertions of the transfer genes into the genome can produce transgenic plants with different expression levels, allowing the choice of transgenic plants with less fitness costs while maintaining broad-spectrum resistance. The induction, but not the constitutive presence, of the defense response such as high contents of H₂O₂ and callose deposition indicates that plants have some mechanisms to suppress inappropriate

immunity and keep balances between normal growth and defense responses. It is reasonable to hypothesize that plants must make corresponding changes at the transcriptional and physiological level to deal with the stress from the interfamily transferred NLRs.

Conditional expression of the autoactive NLRs is a promising method to reduce the fitness costs. The TBF1-cassette, a pathogen responsive translational control system, has been successfully used to conditionally regulate the protein level of NPR1, conferring broad-spectrum disease resistance but also avoiding fitness costs (Xu et al., 2017). The transcription activator-like (TAL) effectors can specially bind to the promoter sequences of their target genes. Engineering promoters to trap pathogen TAL effectors from pathogens can specially activate resistance genes at the infection sites (Zeng et al., 2015). Additionally, the global transcription analysis of the plants inoculated with target pathogens can find some pathogen-induced genes. The promoters from these genes can be used to restrict the expression of the autoactive NLRs at infection sites.

AUTHOR CONTRIBUTIONS

ZL and ZG designed the study and wrote the manuscript. ZL carried out most experiments and analyzed data. JH, ZW, SZ, and XW contributed to obtain the transgenic plans. FM and ZZ contributed to rice blast assay.

FUNDING

This study was supported by The National Key Research and Development Program of China (2016YFD0100600 to ZG) and National Nature Science Foundation of China (31270315 to ZG).

ACKNOWLEDGMENTS

We thank Dr. Guangcun He, Jinxiong Sun, and Jinqiang He from Wuhan University for guidance on BPH resistance assays, and Dr. Shaoqing Li for providing *Xoo* strain PXO-99. We also thank Dr. Xin Yuan and Xiaoxiao Liu for critical reading and thoughtful inputs to the manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2019.00417/full#supplementary-material>

FIGURE S1 | Relative expression levels of *AtRPS2* and *AtRPM1(D505V)* in the transgenic plants. The expression levels of *AtRPS2* (A) and *AtRPM1(D505V)* (B) in seven independent stable transgenic lines were detected by qRT-PCR assay. The *OsActin* gene was used as an internal control. Data are shown as means \pm SD ($n = 3$).

FIGURE S2 | Disease resistance assays with detached leaves of the *AtRPS2*, *AtRPM1(D505V)* transgenic plants and Nipponbare. **(A)** Detached leaves of 4 weeks old plants were lightly wounded with an ear punch, and 8 μ L of spore suspension (5×10^5 /mL) of *M. oryzae* isolate was added to the wound sites. Pictures were taken at 7 dpi, bars = 2 cm. **(B,C)** Lesion length and fungal growth were measured at 7 dpi. Data are shown as means \pm SD ($n = 20$). Asterisks denote significant differences (one way ANOVA, $***P < 0.001$). Similar results were obtained in three independent experiments.

FIGURE S3 | GO analysis of differentially expressed genes between *AtRPS2* transgenic plants and Nipponbare.

FIGURE S4 | GO analysis of differentially expressed genes between *AtRPM1(D505V)* transgenic plants and Nipponbare.

TABLE S1 | Segregation analysis of T₁ *AtRPS2* transgenic plants.

TABLE S2 | Segregation analysis of T₁ *AtRPM1(D505V)* transgenic plants.

TABLE S3 | Differentially expressed genes (DEGs) between Nipponbare and *AtRPS2* transgenic plants.

TABLE S4 | Differentially expressed genes (DEGs) between Nipponbare and *AtRPM1(D505V)* transgenic plants.

TABLE S5 | Co-regulated genes in *AtRPS2* and *AtRPM1(D505V)* transgenic plants.

TABLE S6 | List of qRT-PCR primers used in this study.

REFERENCES

- Axtell, M. J., and Staskawicz, B. J. (2003). Initiation of RPS2-specified disease resistance in *Arabidopsis* is coupled to the avrrpt2-directed elimination of RIN4. *Cell* 112, 369–377. doi: 10.1016/S0092-8674(03)00036-9
- Bent, A. F., and Mackey, D. (2007). Elicitors, effectors, and r genes: the new paradigm and a life time supply of questions. *Ann. Rev. Phytopathol.* 45, 399–436. doi: 10.1146/annurev.phyto.45.062806.094427
- Brutus, A., Sicilia, F., Macone, A., Cervone, F., and De Lorenzo, G. (2010). A domain swap approach reveals a role of the plant wall-associated kinase 1 (WAK1) as a receptor of oligogalacturonides. *Proc. Natl. Acad. Sci. U.S.A.* 107, 9452–9457. doi: 10.1073/pnas.1000675107
- Bundo, M., and Coca, M. (2016). Enhancing blast disease resistance by overexpression of the calcium-dependent protein kinase OsCPK4 in rice. *Plant Biotechnol. J.* 14, 1357–1367. doi: 10.1111/pbi.12500
- Cesari, S., Bernoux, M., Moncuquet, P., Kroj, T., and Dodds, P. N. (2014). A novel conserved mechanism for plant NLR protein pairs: the “integrated decoy” hypothesis. *Front. Plant Sci.* 5:606. doi: 10.3389/fpls.2014.00606
- Chen, J., Peng, P., Tian, J., He, Y., Zhang, L., Liu, Z., et al. (2015). Pike, a rice blast resistance allele consisting of two adjacent NBS-LRR genes, was identified as a novel allele at the pik locus. *Mol. Breed.* 35:117. doi: 10.1007/s11032-015-0305-6
- Cheng, X., Zhu, L., and He, G. (2013). Towards understanding of molecular interactions between rice and the brown planthopper. *Mol. Plant* 6, 621–634. doi: 10.1093/mp/sst030
- Couto, D., and Zipfel, C. (2016). Regulation of pattern recognition receptor signalling in plants. *Nat. Rev. Immunol.* 16, 537–552. doi: 10.1038/nri.2016.77
- Cui, H., Tsuda, K., and Parker, J. E. (2015). Effector-triggered immunity: from pathogen perception to robust defense. *Annu. Rev. Plant Biol.* 66, 487–511. doi: 10.1146/annurev-arplant-050213-040012
- Day, B., Dahlbeck, D., Huang, J., Chisholm, S. T., Li, D., and Staskawicz, B. J. (2005). Molecular basis for the RIN4 negative regulation of RPS2 disease resistance. *Plant Cell* 17, 1292–1305. doi: 10.1105/tpc.104.030163
- Fitzgerald, H. A., Chern, M.-S., Navarre, R., and Ronald, P. C. (2004). Overexpression of (At)NPR1 in rice leads to a bth- and environment-induced lesion-mimic/cell death phenotype. *Mol. Plant Microbe Interact.* 17, 140–151. doi: 10.1094/MPMI.2004.17.2.140
- Fu, Z. Q., and Dong, X. (2013). Systemic acquired resistance: turning local infection into global defense. *Annu. Rev. Plant Biol.* 64, 839–863. doi: 10.1146/annurev-arplant-042811105606
- Gao, Z., Chung, E. H., Eitas, T. K., and Dangl, J. L. (2011). Plant intracellular innate immune receptor resistance to *Pseudomonas syringae* pv. maculicola 1 (RPM1) is activated at, and functions on, the plasma membrane. *Proc. Natl. Acad. Sci. U.S.A.* 108, 7619–7624. doi: 10.1073/pnas.1104410108
- Gu, K., Tian, D., Qiu, C., and Yin, Z. (2009). Transcription activator-like type III effector AvrXa27 depends on OsTFIIAgamma5 for the activation of Xa27 transcription in rice that triggers disease resistance to *Xanthomonas oryzae* pv. *oryzae*. *Mol. Plant Pathol.* 10, 829–835. doi: 10.1111/j.1364-3703.2009.00567.x
- Hu, L., Wu, Y., Wu, D., Rao, W., Guo, J., Ma, Y., et al. (2017). The coiled-coil and nucleotide binding domains of brown planthopper resistance14 function in signaling and resistance against planthopper in rice. *Plant Cell* 29, 3157–3185. doi: 10.1105/tpc.17.00263
- Jing, S., Zhao, Y., Du, B., Chen, R., Zhu, L., and He, G. (2017). Genomics of interaction between the brown planthopper and rice. *Curr. Opin. Insect Sci.* 19, 82–87. doi: 10.1016/j.cois.2017.03.005
- Jones, J. D., and Dangl, J. L. (2006). The plant immune system. *Nature* 444, 323–329. doi: 10.1038/nature05286
- Ke, Y., Deng, H., and Wang, S. (2017). Advances in understanding broad-spectrum resistance to pathogens in rice. *Plant J.* 90, 738–748. doi: 10.1111/tpj.13438
- Lacombe, S., Rougon-Cardoso, A., Sherwood, E., Peeters, N., Dahlbeck, D., van Esse, H. P., et al. (2010). Interfamily transfer of a plant pattern-recognition receptor confers broad-spectrum bacterial resistance. *Nat. Biotechnol.* 28, 365–369. doi: 10.1038/nbt.1613
- Li, Y., Zhang, Y., Wang, Q. X., Wang, T. T., Cao, X. L., Zhao, Z. X., et al. (2018). Resistance to powdery mildew8.1 boosts pattern-triggered immunity against multiple pathogens in arabidopsis and rice. *Plant Biotechnol. J.* 16, 428–441. doi: 10.1111/pbi.12782
- Liu, Q., Yang, J., Yan, S., Zhang, S., Zhao, J., Wang, W., et al. (2016). The germin-like protein OsGLP2-1 enhances resistance to fungal blast and bacterial blight in rice. *Plant Mol. Biol.* 92, 411–423. doi: 10.1007/s11103-016-05214
- Liu, Y., Wu, H., Chen, H., Liu, Y., He, J., Kang, H., et al. (2015). A gene cluster encoding lectin receptor kinases confers broad-spectrum and durable insect resistance in rice. *Nat. Biotechnol.* 33, 301–305. doi: 10.1038/nbt.3069
- Lu, F., Wang, H., Wang, S., Jiang, W., Shan, C., Li, B., et al. (2015). Enhancement of innate immune system in monocot rice by transferring the dicotyledonous elongation factor Tu receptor EFR. *J. Integr. Plant Biol.* 57, 641–652. doi: 10.1111/jipb.12306
- Mackey, D., Belkhadir, Y., Alonso, J. M., Ecker, J. R., and Dangl, J. L. (2003). *Arabidopsis* RIN4 is a target of the type III virulence effector AvrRpt2 and modulates rps2-mediated resistance. *Cell* 112, 379–389. doi: 10.1016/S0092-8674(03)000400
- Mackey, D., Holt, B. F., Wiig, A., and Dangl, J. L. (2002). RIN4 Interacts with *Pseudomonas syringae* Type III effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. *Cell* 108, 743–754. doi: 10.1016/S0092-8674(02)00661-X
- Maekawa, T., Kracher, B., Vernaldi, S., van Themaat, E., and Schulze-Lefert, P. (2012). Conservation of NLR-triggered immunity across plant lineages. *Proc. Natl. Acad. Sci. U.S.A.* 109, 20119–20123. doi: 10.1073/pnas.1218059109
- Molla, K. A., Karmakar, S., Chanda, P. K., Sarkar, S. N., Datta, S. K., and Datta, K. (2016). Tissue-specific expression of *Arabidopsis* NPR1 gene in rice for sheath blight resistance without compromising phenotypic cost. *Plant Sci.* 250, 105–114. doi: 10.1016/j.plantsci.2016.06.005
- Narusaka, M., Kubo, Y., Hatakeyama, K., Imamura, J., Ezura, H., Nanasato, Y., et al. (2013). Interfamily transfer of dual NB-LRR genes confers resistance to multiple pathogens. *PLoS One* 8:e55954. doi: 10.1371/journal.pone.0055954
- Park, C. H., Chen, S., Shirsekar, G., Zhou, B., Khang, C. H., Songkumarn, P., et al. (2012). The *Magnaporthe oryzae* effector AvrPiz-t targets the RING E3 ubiquitin ligase APIP6 to suppress pathogen-associated molecular pattern-triggered immunity in rice. *Plant Cell* 24, 4748–4762. doi: 10.1105/tpc.112.105429
- Park, D.-S., Saylor, R. J., Hong, Y.-G., Nam, M.-H., and Yang, Y. (2008). A method for inoculation and evaluation of rice sheath blight disease. *Plant Dis.* 92, 25–29. doi: 10.1094/pdis-92-1-0025

- Pieterse, C. M., Zamioudis, C., Berendsen, R. L., Weller, D. M., Van Wees, S. C., and Bakker, P. A. (2014). Induced systemic resistance by beneficial microbes. *Annu. Rev. Phytopathol.* 52, 347–375. doi: 10.1146/annurev-phyto-082712102340
- Schoonbeek, H. J., Wang, H. H., Stefanato, F. L., Craze, M., Bowden, S., Wallington, E., et al. (2015). *Arabidopsis* EF-Tu receptor enhances bacterial disease resistance in transgenic wheat. *New Phytol.* 206, 606–613. doi: 10.1111/nph.13356
- Shangguan, X., Zhang, J., Liu, B., Zhao, Y., Wang, H., Wang, Z., et al. (2018). A mucin-like protein of planthopper is required for feeding and induces immunity response in plants. *Plant Physiol.* 176, 552–565. doi: 10.1104/pp.17.00755
- Shimono, M., Koga, H., Akagi, A., Hayashi, N., Goto, S., Sawada, M., et al. (2012). Rice WRKY45 plays important roles in fungal and bacterial disease resistance. *Mol. Plant Pathol.* 13, 83–94. doi: 10.1111/j.1364-3703.2011.00732.x
- Spoel, S. H., and Dong, X. (2012). How do plants achieve immunity? defence without specialized immune cells. *Nat. Rev. Immunol.* 12, 89–100. doi: 10.1038/nri3141
- Tsuda, K., and Katagiri, F. (2010). Comparing signaling mechanisms engaged in pattern-triggered and effector-triggered immunity. *Curr. Opin. Plant Biol.* 13, 459–465. doi: 10.1016/j.pbi.2010.04.006
- Wang, B. H., Ebbole, D. J., and Wang, Z. H. (2017). The arms race between *Magnaporthe oryzae* and rice: diversity and interaction of avr and r genes. *J. Integr. Agric.* 16, 2746–2760. doi: 10.1016/s2095-3119(17)617465
- Wang, X., Hou, S., Wu, Q., Lin, M., Acharya, B. R., Wu, D., et al. (2017). IDL6-HAE/HSL2 impacts pectin degradation and resistance to *Pseudomonas syringae* pv tomato DC3000 in *Arabidopsis* leaves. *Plant J.* 89, 250–263. doi: 10.1111/tpj.13380
- Wang, R., Ning, Y., Shi, X., He, F., Zhang, C., Fan, J., et al. (2016). Immunity to rice blast disease by suppression of effector-triggered necrosis. *Curr. Biol.* 26, 2399–2411. doi: 10.1016/j.cub.2016.06.072
- Wu, J., Kou, Y., Bao, J., Li, Y., Tang, M., Zhu, X., et al. (2015). Comparative genomics identifies the *Magnaporthe oryzae* avirulence effector AvrPi9 that triggers Pi9-mediated blast resistance in rice. *New Phytol.* 206, 1463–1475. doi: 10.1111/nph.13310
- Wulff, B. B. H., Horvath, D. M., and Ward, E. R. (2011). Improving immunity in crops: new tactics in an old game. *Curr. Opin. Plant Biol.* 14, 468–476. doi: 10.1016/j.pbi.2011.04.002
- Xu, G., Yuan, M., Ai, C., Liu, L., Zhuang, E., Karapetyan, S., et al. (2017). uORF-mediated translation allows engineered plant disease resistance without fitness costs. *Nature* 545, 491–494. doi: 10.1038/nature22372
- Yang, S., Feng, Z., Zhang, X., Jiang, K., Jin, X., Hang, Y., et al. (2006). Genome-wide investigation on the genetic variations of rice disease resistance genes. *Plant Mol. Biol.* 62, 181–193. doi: 10.1007/s11103-006-90123
- Yuan, H. M., Liu, W. C., and Lu, Y. T. (2017). CATALASE2 coordinates sa-mediated repression of both auxin accumulation and ja biosynthesis in plant defenses. *Cell Host Microbe* 21, 143–155. doi: 10.1016/j.chom.2017.01.007
- Zeng, X., Tian, D., Gu, K., Zhou, Z., Yang, X., and Luo, Y. (2015). Genetic engineering of the Xa10 promoter for broad-spectrum and durable resistance to *Xanthomonas oryzae* pv. *oryzae*. *Plant Biotechnol. J.* 13, 993–1001. doi: 10.1111/pbi.12342
- Zhang, H., and Wang, S. (2013). Rice versus *Xanthomonas oryzae* pv. *oryzae*: a unique pathosystem. *Curr. Opin. Plant Biol.* 16, 188–195. doi: 10.1016/j.pbi.2013.02.008
- Zipfel, C., and Oldroyd, G. E. D. (2017). Plant signalling in symbiosis and immunity. *Nature* 543, 328–336. doi: 10.1038/nature22009

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Li, Huang, Wang, Meng, Zhang, Wu, Zhang and Gao. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.