

Three true hopper species new for the Netherlands (Homoptera: Delphacidae & Cicadellidae)

C.F.M. (Kees) den Bieman
M.C. (Marco) de Haas
J.D.M. (Dick) Belgers

KEY WORDS

Auchenorrhyncha, Cicadomorpha, Climate change, Fulgoromorpha, *Toya*

Entomologische Berichten 82 (3): 90-93

In 2021, three species were identified as new for the Netherlands: the planthopper *Toya propinqua* and the leafhoppers *Cicadula placida* and *Edwardsiana alnicola*. The first two species clearly show, probably due to climate change, a range expansion from the south. The ecology and distribution of each species are discussed.

Introduction

Over the last 20 years, fieldwork has intensified, including the use of malaise and light traps. This has resulted in the addition of 51 true hopper species to the last fauna list of 2011 (Den Bieman et al. 2011) that are new for the Netherlands (Den Bieman et al. 2020, 2021a, 2021b, 2021c and references therein).

Three more species were identified in the Netherlands for the first time and presented in this article: *Toya propinqua* (Fieber, 1866), *Cicadula placida* (Horváth, 1897) and *Edwardsiana alnicola* (Edwards, 1924). Details on the distribution and ecology of these species are given.

The Dutch geographical coordinates (Amersfoortcoördinaten = AC) are given for each collection site. Unless stated otherwise, the material was collected by the authors and preserved in their collections. Identifications of all treated species are based on Biedermann & Niedringhaus (2004) and Ribaut (1952).

New species for the Dutch fauna

Delphacidae

Toya propinqua (figure 1)

Material Province of Zeeland: 3.5 km northwest of Kamperland, AC 35.1-401.1, 12.v.2021, 1 macropterous ♂.

This specimen was swept in a medium dry and sandy meadow next to a deciduous forest edge (figure 2), mown once a year and also frequented by pedestrians.

Toya is primarily a tropical genus represented by five species in Europe, mainly in the Mediterranean region. A critical evaluation of the European species is given in Della Giustina (2019). The discovery of the first *T. propinqua* in the Netherlands does not stand on its own: an increasing number of southern plant- and leafhopper species have reached the Netherlands in recent years. *Toya propinqua* occurs in most of the southern and central European countries: Albania, Austria, Bosnia Herzegovina,

1. Male of *Toya propinqua*, reared by H. Strübing. Photo: Gernot Kunz

1. Mannetje van *Toya propinqua*, uit een kweek van H. Strübing.





2. Collecting site of *Toya propinqua*. Photo: Kees den Bieman
2. Vindplaats van *Toya propinqua*.

Bulgaria, Croatia, Cyprus, Czech Republic, France, Germany, Greece, Hungary, Italy, Macedonia, Moldavia, Poland, Portugal, Romania, Slovakia, Slovenia, Southern European Russia, Spain, Switzerland, Ukraine and former Yugoslavia (Jach & Hoch 2013). Its occurrence north of the Alps is perhaps only irregularly (Nickel 2003). *Toya propinqua* is a pantropical species in the warmer regions of the Old World and the Nearctic (Jach & Hoch 2013, Wilson & Claridge 1991).

In Europe, *T. propinqua* is probably bivoltine with overwintering nymphs. Adults can be found from February till October (Della Giustina 2019, Nickel 2003). *Toya propinqua* is an eurytopic pioneer species, often collected on disturbed sites (Holzinger 2009, Nickel 2003).

It is an oligophagous species living on grasses, in particular *Cynodon dactylon*. It has also been observed on *Paspalum thunbergia*, *Digitaria*, *Zoysia japonica* and *Panicum repens* (Della Giustina 2019, Drosopoulos et al. 1983, Nickel 2003). This species is frequently collected in rice fields but there is little evidence of any significant damage to this crop (Wilson & Claridge 1991). However, dense populations can cause damage to young barley plants. Transmission of *Cynodon* chlorotic streak virus 16Sr is observed on maize and *Cynodon dactylon* (Della Giustina 2019, Lockhart et al. 1985).

In a study on samples from suction traps at 12 m height in

France, *T. propinqua* was collected in numbers, all macropterous, in 30% of the traps from May till October (Della Giustina & Balasse 1999). Passive dispersion over longer distances by air currents seems likely for the flying individuals. In Iceland, a brachypterous male was collected on a locality close to a domestic airstrip (Remane 2005), indicating that it has probably arrived by plane or due to passive long-distance migration with jet-streams (Söderman et al. 2009). The first specimen of this species collected in the Netherlands is probably also a migrant. A collecting trip to the same locality at July 21, 2021 was in vain. A second delphacid species living on the grass *Cynodon dactylon* is *Laodelphax striatellus* (Fallén, 1826). This is also a vagrant species that has rarely been collected in the Netherlands before 2003. However, since 2003 *L. striatellus* is collected once or twice a year and it is sometimes able to establish a semi-permanent population (Den Bieman 2017, personal observations). Perhaps the same will happen to *T. propinqua*.

Cicadellidae-Cicadellinae

Cicadula placida (figure 3)

Material Province of Gelderland: Wageningen, AC 174.7-442.0, light trap, 9.vii.2021, 1 ♂.

The light trap was placed in a city garden. This light trap has been used regularly since 2018, but only four *Cicadula* specimens were collected since that time: one male of *C. flori* (Sahlberg, 1871), one male of *C. placida* and two unidentified females. Strong attraction of *C. placida* on dispersal flight by light is shown by samples collected in light traps in the suburbs of Brno (Czech Republic) (Malenovský & Lauterer 2005, Malenovský et al. 2011).

The genus *Cicadula* comprises fifteen species in Europe. Three belong to the subgenus *Henriana*, the Mediterranean species *C. lineatopunctata* (Matsumura, 1908) and two species that occur also in the Netherlands: *C. placida* and *C. frontalis* (Herrich-Schäffer, 1835). Differences between these species are given by Ribaut (1952). The antennae of *C. placida* are shorter than those of *C. frontalis* and the aedeagi differ in shape. The appendages of the aedeagus of *C. placida* are inserted closer to the base than in *C. frontalis* (see also Lock 2019). In Italy specimens collected in June were light yellow-green (var. *inornata* Ribaut, 1952) and specimens collected in November brown (Guglielmino et al. 2017). The single Dutch specimen is green.

Cicadula placida has primarily a south-eastern European distribution: Austria, Bulgaria, Czech Republic, France, Greece,

3. *Cicadula placida*, Geidorf, Graz, Austria, 19.vi.2009. Photo: Gernot Kunz

3. *Cicadula placida*, Geidorf, Graz, Oostenrijk, 19.vi.2009.





4. Female of *Edwardsiana cf. alnicola*, Göttingen, Niedersachsen, Germany, 6.ix.2009. Photo: Gernot Kunz

4. Vrouwtje van *Edwardsiana cf. alnicola*, Göttingen, Nedersaksen, Duitsland, 6.ix.2009.

Hungary, Italy, Moldavia, Romania, Slovenia, South Eastern Russia and former Yugoslavia (Jach & Hoch 2013). Recently it was collected in the south-eastern part of Germany (Nickel 2011) and Belgium (Lock 2019). Since the identification of the first German sample in 2011, the species spread rapidly over the south of Germany (Achtziger et al. 2020). Outside Europe this species is known from a mountainous area in Iran (Mozaffarian & Wilson 2016).

The biology of *C. placida* has not been extensively studied. In Austria, adults were found from June till October, in Italy in June and July and again in October and November. The Italian data suggest that it is a bivoltine species, at least in the southern part of its distribution area. Eggs are the hibernating stage (Holzinger 2009, Guglielmino et al. 2017). *Cicadula placida* is collected on eutrophic shores of rivers and ponds, and open marshy sites in floodplain forests. These are all hygrophilous and thermophilous habitats. In Italy and France, it is collected, sometimes in high numbers, on *Phalaris arundinacea* and *Glyceria maxima* (and perhaps other Poacea species) (Guglielmino et al. 2017, Ribaut 1952). Holzinger (2009) indicates the same habitat type in Austria but mentions *Carex* as host plant. Malenovský et al. (2011) report *Carex acuta* as a host plant. The closely related *C. frontalis* lives on *Carex* species (Nickel 2003).

Cicadellidae-Typhlocybae

Edwardsiana alnicola (figure 4)

Material Province of Noord-Brabant: Udenhout, De Brand, AC 137.4-404.8, malaise trap, 13-20.vi.2020, 1 ♂, leg. IWG KNNV Tilburg; same location, malaise trap, 4-11.vii.2020, 1 ♂, leg. IWG KNNV Tilburg.

The taxonomically difficult genus *Edwardsiana* is species rich with at least 51 species in Europe (Jach & Hoch 2013) of

which 23 species are listed for the Netherlands (Den Bieman & De Haas 2018, Den Bieman & Van Klink 2015, Den Bieman et al. 2011, 2020). *Edwardsiana alnicola* is characterized by the shape and the orientation of the aedeagus appendages (Biedermann & Niedringhaus 2004).

Edwardsiana alnicola is known from Austria, Czech Republic, Denmark, Estonia, Finland, Germany, Great Britain, Italy, Latvia, Norway, Poland, Romania, Slovenia, Sweden and Switzerland (Jach & Hoch 2013, Löcker 2003). The species has not been recorded in France, but it was also collected in Belgium recently (Lock in preparation). Throughout its geographical range, it is a rather rare or local species and therefore poorly known (Nickel 2003, Walczak et al. 2018). After the first records in 2020, extensive field research in 2021 was however unsuccessful in collecting *E. alnicola* in the nature reserve De Brand, the same location where the two Dutch specimens were found previously. This suggests that *E. alnicola* is also rare in this area.

De Brand is partly a very wet marshy deciduous forest, which corresponds to the habitat that is described for *E. alnicola* (Holzinger 2009, Nickel 2003). Specific habitat requirements are not known but the extreme localisation and rarity of the species indicate that such factors exist (Kirby 1992). It lives on *Alnus glutinosa* and *A. incana*, both very common species in the Netherlands (<http://verspreidingsatlas.nl>) and in De Brand. Adults are found from June till August, *E. alnicola* is a bivoltine species, and it hibernates in the egg stage (Holzinger 2009, Nickel 2003).

Acknowledgements

We are highly indebted to Gernot Kunz for the beautiful photos. We want to thank André van Eck for providing the Auchenorrhyncha specimens from a malaise trap in nature reserve De Brand.

Literature

- Achtziger R, Chen P-P, Funke L, Hartung V, van Klink R, Malenovský I, Nickel H, Nieser N, Rösch V, Walter S, Zeman Š & Jessat M 2020. Zur Zikaden- und Wanzenfauna von Wasserbüffelweiden und angrenzenden Flächen im Altenburger Land (Thüringen) –Ergebnisse der Sammelexkursionen zur 26. Jahrestagung des Arbeitskreises Zikaden Mitteleuropas 2019 (Hemiptera: Auchenorrhyncha, Heteroptera). *Mauritiana* (Altenburg) 38: 137-162.
- Biedermann R & Niedringhaus R 2004. Die Zikaden Deutschlands. Bestimmungstabellen für alle Arten. WABV Fründ.
- Della Giustina W & Balasse H 1999. Gone with the wind: Homoptera Auchenorrhyncha collected by the French network of suction traps in 1994. *Marburger entomologische Publikationen* 3: 7-42.
- Della Giustina W 2019. Les Delphacidae de France et des pays limitrophes (Hemiptera, Fulgoromorpha). *Faune de France* 100: 1-831.
- Den Bieman CFM 2017. The first population of the small brown planthopper *Laodelphax striatellus* (Homoptera: Delphacidae) in the Netherlands. *Entomologische Berichten* 77: 219-222.
- Den Bieman CFM & De Haas M 2018. Vier nieuwe dwergcicaden voor Nederland (Hemiptera: Auchenorrhyncha: Cicadellidae). *Entomologische Berichten* 78: 102-106.
- Den Bieman CFM & Van Klink R 2015. Een forse uitbreiding van de Nederlandse

- dwergcicadenfauna met vijftien soorten (Hemiptera: Auchenorrhyncha: Cicadellidae). Entomologische Berichten 75: 211-226.
- Den Bieman K, Biedermann R, Nickel H & Niedringhaus R 2011. The planthoppers and leafhoppers of Benelux. Identification keys to all families and genera and all Benelux species not recorded from Germany. WABV Fründ.
- Den Bieman CFM, De Haas M, Belgers JDM, De Somer N, Lommen G & Reinboud M 2020. A remarkable extension of the Dutch planthopper and leafhopper fauna with nine species (Homoptera: Auchenorrhyncha: Cicadellidae, Cixiidae & Tettigometridae). Entomologische Berichten 80: 138-146.
- Den Bieman CFM, Tanis M, Drukker EF & De Waart S 2021a. The leafhopper fauna of green roofs including the Mediterranean leafhopper *Circulifer haematoceps* new for the Netherlands (Auchenorrhyncha: Cicadellidae). Entomologische Berichten 81: 46-51.
- Den Bieman CFM, de Haas MC, Langbroek AJ & Van Leeuwen J 2021b. New and interesting true hoppers for the Netherlands (Homoptera: Cixiidae, Delphacidae & Cicadellidae). Entomologische Berichten 81: 244-251.
- Den Bieman CFM, De Haas MC & Soethof R 2021c. The leafhopper genus *Rhopalopyx* in the Netherlands (Hemiptera: Auchenorrhyncha: Cicadellidae). Nederlandse Faunistische Mededelingen 57: 95-106.
- Drosopoulos S, Asche M & Hoch H 1983. Contribution to the planthopper fauna of Greece (Homoptera, Auchenorrhyncha, Fulgoromorpha, Delphacidae). Annales de l'Institut phytopathologique Benaki (N.S.) 14: 19-68.
- Guglielmino L, Scarici L, De Santis A & Bückle C 2017. Study on flora and Auchenorrhyncha biocenoses (Insecta Hemiptera) in moist areas considered restricted relics of the ancient Lacus Velinus in the provinces of Terni and Rieti (Umbria and Latium, Italy). Biodiversity Journal 8: 151-184.
- Holzinger WE 2009. Rote Liste der Zikaden (Hemiptera: Auchenorrhyncha) Österreichs. In: Rote Liste gefährdeter Tiere Österreichs 3: 41-317.
- Jach M & Hoch H 2013. Fauna Europaea: Cicadomorpha, Cicadellidae. Fauna Europaea version 2.6.2. Available on: www.faunaeur.org [accessed January 21, 2022].
- Kirby P 1992. A review of the scarce and threatened Hemiptera of great Britain. Joint Nature Conservation Committee no. 2.
- Lockhart BEL, Khaless N, El Maataoui M & Lastra R 1985. Cynodon Chlorotic Streak Virus, a previously undescribed plant rhabdovirus infecting Bermuda grass and Maize in the Mediterranean area. Phytopathology 75: 1094-1098.
- Löcker H 2003. Arborikole Zikaden-Gilden in Slowenien (Hemiptera, Auchenorrhyncha). Beiträge zur Zikadenkunde 6: 7-38.
- Lock K 2019. Fifty leafhoppers new to Belgium (Hemiptera: Cicadellidae). Belgian Journal of Entomology 88: 1-28.
- Malenovský I & Lauterer P 2005. Leafhoppers, planthoppers and psyllids (Hemiptera: Cicadomorpha, Fulgoromorpha, Psylloidea) in ruderal habitats: material attracted by light in the suburbs of Brno (Czech Republic). Acta Musei Moraviae Scientiae Biologicae 90: 195-207.
- Malenovský I, Baňář P & Kment P 2011. A contribution to the faunistics of the Hemiptera (Cicadomorpha, Fulgoromorpha, Heteroptera, and Psylloidea) associated with dry grassland sites in southern Moravia (Czech Republic). Acta Musei Moraviae, Scientiae Biologicae 96: 41-187.
- Mozaffarian F & Wilson MR 2016. A checklist of the leafhoppers of Iran (Hemiptera: Auchenorrhyncha: Cicadellidae). Zootaxa 4062: 1-63
- Nickel H 2003. The leafhoppers and planthoppers of Germany (Hemiptera, Auchenorrhyncha). Patterns and strategies in a highly diverse group of phytophagous insects. Pensoft Publishers.
- Nickel H 2011. Zikaden (Hemiptera: Auchenorrhyncha). In: Biologische Vielfalt im Nationalpark Bayerischer Wald (Jehl H, Müller J, Bässler C & Pöhlmann R eds): 141-144. Sonderband der Wissenschaftlichen Schriftenreihe des Nationalparks Bayerischer Wald Grafenau.
- Remane R 2005. Zur Zikadenfauna Islands, einer nur geologisch „alten“ Vulkaninsel: Analyse eines 2001 von Dr. M. v. Tschirnhaus gesammelten Materials (Insecta Rhyncota Auchenorrhyncha). Marburger entomologische Publikationen 3: 111-123.
- Ribaut H 1952. Homptères Auchenorrhynques II (Jassidae). Faune de France 57: 1-474.
- Söderman G, Gyllenfors G & Endrestöl A 2009. An annotated catalogue of the Auchenorrhyncha of Northern Europe (Insecta, Hemiptera, Fulgoromorpha et Cicadomorpha). Cicadina 10: 33-69.
- Walczak M, Kaszyca N & Taszakowski A 2018. Materiały do znajomości piewików (Hemiptera: Fulgoromorpha, Cicadomorpha) Beskidu Wschodniego. Acta entomologica silesiana 26: 1-9.
- Wilson MR & Claridge MF 1991. Handbook for the identification of leafhoppers and planthoppers of rice. CAB International.

Accepted: March 29, 2022

Samenvatting

Drie nieuwe cicaden voor de Nederlandse fauna (Homoptera: Delphacidae & Cicadellidae)

Sinds 2011 zijn 51 cicadensoorten toegevoegd aan de Nederlandse faunalijs. Recent zijn weer drie soorten nieuw voor Nederland gevonden: *Toya propinqua*, *Cicadula placida* en *Edwardsiana alnicola*. *Toya propinqua* is een pantropische en mediterrane soort die steeds vaker in noordelijke landen gesignaleerd wordt. *Cicadula placida* breidt recent haar areaal vanuit Zuidoost-Europa uit. Beide zijn voorbeelden van zuidelijke soorten die hun areaal door het opwarmende klimaat noordwaarts uitbreiden. *Edwardsiana alnicola* is een soort die op els leeft en in haar hele areaal zeldzaam of zeer lokaal is. Specifieke habitateisen van deze soort zijn nog niet bekend. Van alle soorten wordt de verspreiding en de biologie beschreven.

C.F.M. (Kees) den Bieman
Ulvenhout, the Netherlands
cdbieman@planet.nl

M.C. (Marco) de Haas
Ede, the Netherlands

J.D.M. (Dick) Belgers
Wageningen, the Netherlands

