

The systematics of the Hemiptera

Sistemática de Hemiptera

DIMITRI FORERO¹

Abstract: The Order Hemiptera comprises four main clades: Sternorrhyncha, Auchenorrhyncha, Coleorrhyncha and Heteroptera. In this article the main phylogenetic hypotheses for Hemiptera and its subgroups are reviewed. Important references regarding catalogs and identification aids are provided for the major groups, making emphasis in recent on-line tools.

Key words: Catalogs. Diversity. Heteroptera. Phylogeny. On-line taxonomic tools. Taxonomy.

Resumen: El Orden Hemiptera comprende cuatro clados: Sternorrhyncha, Auchenorrhyncha, Coleorrhyncha y Heteroptera. En este artículo se revisan las principales hipótesis filogenéticas para Hemiptera y sus subgrupos. También se brindan referencias importantes para la identificación de los grupos principales, haciendo énfasis en herramientas recientes disponibles en la web.

Palabras clave: Catálogos. Diversidad. Filogenia. Heteroptera. Herramientas taxonómicas en línea. Taxonomía.

Introduction

Hemiptera is the fifth largest group of insects after Coleoptera, Diptera, Hymenoptera, and Lepidoptera (Schuh and Slater 1995; Grimaldi and Engel 2005; Cameron *et al.* 2006), and the most diversified group of non-endopterygote insects (Kristensen 1991), with approximately 82.000 described species (Arnett 2000). Their feeding habits range from phytophagy to predation, including ectoparasitism and hematophagy. Many of them are important pest species to cultivated crops and some are important vectors of human diseases.

This review deals with the phylogenetic systematic hypotheses for Hemiptera. Throughout the text, common phylogenetic terminology is used in cladistics that the non-familiarized reader may find confusing. The reader shall refer to general references for an explanation of those terms and the logic behind it (e.g., Hennig 1966; Kitching *et al.* 1998; Schuh 2000; Albert 2005).

The number of described species within Hemiptera is not exactly known. It is considered to have roughly 82.000 described species (Slater 1982; Hodkinson and Casson 1991; Arnett 2000), a number that must be seen as conservative. For instance, an estimate of the total world fauna using Asian tropical rain forest samples lies between 184.000 and 193.000 species (Hodkinson and Casson 1991). In some diversified groups, as Cicadellidae, the proportion of undescribed tropical taxa may be as high as 90% (Dietrich and Wallner 2002). Of the total described species, merely three families account for most of the diversity: Cicadellidae, with approximately 22.000 species (Auchenorrhyncha) (Metcalf 1962a, 1962b, 1962c, 1962d, 1963a, 1963b, 1963c, 1963d, 1964a, 1965a, 1965b, 1966a, 1966b, 1966c, 1966d, 1967, 1968; Oman *et al.*

1990; McKamey 2001), and Miridae and Reduviidae (Heteroptera), with more than 10.000 and 6.000, respectively (Schuh 1995; Maldonado 1990).

Characters defining Hemiptera

Hemiptera has long been recognized as a monophyletic group (Hennig 1969; Carver *et al.* 1991). It can be recognized by the particular structure of the mouthparts: the mandibles and maxillary laciniae are modified into concentric stylets, the mandibular enclosing the maxillary ones, both forming the food and salivary channels; the multisegmented sheetlike labium is covering the mandibular and maxillary stylets; and the maxillary and labial palpi are always absent (Weber 1930; Hennig 1969, 1981; Cobben 1978; Kristensen 1991).

Paraphyly of Homoptera

There was a long-standing tradition to treat the Homoptera and Heteroptera as separate groups usually having each the rank of order or as suborders within Hemiptera (e.g., Brues *et al.* 1954; Borror and White 1970; Borror *et al.* 1981). The former approach was common in North American entomology, in which Hemiptera included Heteroptera only, thus restricting the concept for Hemiptera (Schuh and Slater 1995).

Morphological evidence, nonetheless, pointed out that Homoptera was probably paraphyletic (e.g., Goodchild 1966; Schlee 1969d; Bourgoin 1986a, 1986b, 1993; Sweet 1996), or at least that evidence of its monophyly was not documented (Schlee 1969d). Some of the alleged characters supporting "Homoptera" mentioned were: enlarged foramen in the head, large sutures defining the mandibular plate, forewing larger than hindwing, reduced tarsomeres, and simple sperm (Bou-

¹ American Museum of Natural History, New York, NY 10024-5192, USA, and Department of Entomology, Comstock Hall, Cornell University, Ithaca, NY 14853-2601, USA. idf2@cornell.edu.

dreaux 1979; Hamilton 1981). Nonetheless, some of these characters are not synapomorphies (e.g., forewing–hindwing character), or may be based on reductions, which are difficult to homologize (e.g., number of tarsomeres).

Hennig (1969, 1981) doubted the monophyly of Homoptera, stating that the characters used for distinguish it from Heteroptera were symplesiomorphies. He recognized three groups within Hemiptera: Sternorrhyncha, Auchenorrhyncha, and Heteropteroidea (as “Heteropteroidea”), the latter clade formed by Coleorrhyncha + Heteroptera (Schlee 1969d). The paraphyly of Homoptera was further corroborated with 18S rDNA sequence analyses (Wheeler *et al.* 1993; Campbell *et al.* 1994, 1995; Sorensen *et al.* 1995; von Dohlen and Moran 1995).

Even though evidence is compelling towards a paraphyletic “Homoptera”, and that this has been adequately communicated to a more general audience (e.g., Carver *et al.* 1991; Kristensen 1991; Gullan 2001; Fagua 2005), it is still frequent to see references to “Homoptera” in areas such as Integrated Pest Management (e.g., Pedigo 1996), or in general entomological textbooks (e.g., Arnett 2000), practice that should be avoided.

Phylogenetic hypotheses

1. Paraneoptera

Paraneoptera is the group of Neoptera that includes Thysanoptera, Psocodea (“Phthiraptera” + “Psocoptera” [Lyal 1985; Yoshizawa and Johnson 2006]), and Hemiptera (Kristensen 1991; Wheeler *et al.* 2001). Kristensen (1991) listed for this clade the following putative synapomorphies: maxillary lacinia elongate and slender, detached from stipes; postclypeus enlarged; reduced number of tarsomeres (three or less); cerci absent; at most six Malpighian tubules; abdominal ganglia fused in one mass; gonangulum in the females fused with tergum nine (Scudder 1961); and sperm acrosome without perforatorium (Jamieson *et al.* 1999). Yoshizawa and Saigusa (2001) proposed nine additional synapomorphies from the front wing axillary sclerites for the group.

Some authors have included Zoraptera in Paraneoptera based on morphological characters (Hennig 1969, 1981; Kristensen 1981; Wheeler *et al.* 2001; Beutel and Weide 2005), in which the inclusive group of Psocodea + Thysanoptera + Hemiptera is denominated Acercaria, due to the absence of cerci (Börner 1904). The inclusion of Zoraptera in Paraneoptera, nonetheless, is highly controversial (Hennig 1969, 1981; Yoshizawa 2007), and has not always been followed (e.g., Kristensen 1991; Grimaldi and Engel 2005). The putative synapomorphies of Zoraptera + Acercaria are all reductions (number of Malpighian tubules, number of tarsomeres, and fusion of abdominal ganglia) (Hennig 1969, 1981; Kristensen 1981; but see Hünefeld 2007), and thus, difficult to homologize. Potential synapomorphies have been proposed for Zoraptera + Acercaria based on head capsule characters (Beutel and Weide 2005), and on genitalic characters both for Zoraptera + Psocoptera, and for Zoraptera + Hemiptera (Hünefeld 2007), but have not been tested in a phylogenetic analysis. Nonetheless, some authors had rejected the inclusion of Zoraptera in Paraneoptera, based on phylogenetic analyses of 18S rDNA sequences, which place Zoraptera closely related to Dictyoptera (Wheeler *et al.* 2001; Yoshizawa and Johnson 2005), and on wing base morphology (Yoshizawa 2007). These evidences support the hypothesis that Zoraptera is not a member of the Paraneoptera.

Relationships among members of Paraneoptera (in the restricted sense of Kristensen 1991) are still unclear. Some phylogenetic hypotheses show a polytomy among Psocodea, Thysanoptera, and Hemiptera (e.g., Kristensen 1991; Beutel and Gorb 2001). In other instances Hemiptera is the sister group of Thysanoptera + Psocodea (e.g., Whiting *et al.* 1997; Gorb and Beutel 2001; Wheeler *et al.* 2001; Willmann 2004), or the sister group of Thysanoptera (e.g., Hennig 1969, 1981; Kristensen 1981; Whiting *et al.* 1997; Wheeler *et al.* 2001; Yoshizawa and Saigusa 2001; Cranston and Gullan 2003). The latter clade, Hemiptera + Thysanoptera, is named Condylgnatha (Börner 1904), and is supported by a few synapomorphies (Kristensen 1981), namely the modification of the mandibles and maxillary laciniae into stylets (Hennig 1969, 1981; Heming 1980), specialized sclerotized rings between antennal flagellomeres (Seeger 1975), and the particular conformation of the axillary sclerites of the forewing (Yoshizawa and Saigusa 2001). The character of the flagellomeres rings should be further evaluated since it was found only in Thysanoptera and in Pentatomidae (Heteroptera) (Seeger 1975).

No phylogenetic hypotheses have been proposed yet for Paraneoptera using molecular characters with a comprehensive taxon sampling. Crespi *et al.* (1996) analyzed 18S rDNA sequences of Thysanoptera with some Hemiptera as outgroups which resulted in a monophyletic Thysanoptera, but in a combined analysis with COI sequences it produced a polytomy of the two suborders of Thysanoptera with Hemiptera. Morris and Mound (2003) in an analysis with an extended Thysanoptera taxon sampling, with some Hemiptera and Psocodea as outgroup species, found a monophyletic Thysanoptera but a paraphyletic Hemiptera with respect to Psocodea. Johnson *et al.* (2004) while studying the relationships of Psocodea using 18S DNA, found moderate support for the Condylgnatha, but again, with an extremely limited taxon sampling for Hemiptera and Thysanoptera.

In those phylogenetic analyses of Hexapoda including paraneopteran representatives, usually the taxon sample is small (e.g., Whiting *et al.* 1997) or some groups are missing completely (e.g., Thysanoptera: Kjer 2004), rendering the conclusions of paraneopteran relationships inadequate. When representatives of all the three major clades are included, the results are either an unresolved polytomy (Kjer *et al.* 2006) or unlikely poly- or paraphyletic groups (e.g., Hemiptera paraphyletic respect to Thysanoptera: Whiting *et al.* 1997; Paraneoptera and Hemiptera polyphyletic: Wheeler *et al.* 2001; Paraneoptera paraphyletic with Psocodea as basal-most Neoptera and Polyneoptera as sister group of Heteropteroidea + Auchenorrhyncha: Whiting 2002; Psocodea paraphyletic respect to Thysanoptera: Kjer *et al.* 2006). The relationships among members of Paraneoptera are an open field of research.

2. Hemiptera

Hemiptera is an ancient lineage with fossils known since the Early Permian (Kukalová-Peck 1991; Shcherbakov and Popov 2002). Hemiptera is a monophyletic group, based on the unique structure of the mouthparts (see above), which consist of a labium enclosing the maxillary and mandibular stylets (Kristensen 1991). Yoshizawa and Saigusa (2001) proposed another potential synapomorphy, the fork of the anterior axillary fold-line of the forewing, a character that is independent from that one of the mouthparts, which reinforces the monophyly of Hemiptera.

As discussed above, Hemiptera was traditionally divided in two groups: “Homoptera” and Heteroptera. Compelling evidence, nonetheless, suggest that Hemiptera is composed of three main clades: Sternorrhyncha, Auchenorrhyncha, and Heteropteroidea, in which Sternorrhyncha is the sister group of Auchenorrhyncha + Heteropteroidea (e.g., Hennig 1969, 1981; Kristensen 1975; Wootton and Betts 1986; Carver *et al.* 1991; Wheeler *et al.* 1993). Of these, only the monophyly of Auchenorrhyncha is still debatable (see below). Zrzavý (1990) termed the clade Auchenorrhyncha + Heteropteroidea as Euhemiptera.

Shcherbakov and Popov (2002) proposed an alternative classification for Hemiptera, mostly following Börner (1904), in which they recognized five suborders and an “ancestral” group, the Archescytinoidea. In their view, the Auchenorrhyncha (as Cicadina) is paraphyletic, and the extant hemipterans that represent the sister group of the Heteropteroidea (as Peloridiina + Cimicina) are the Cicadomorpha. They presented a phylogram depicting relationships among these groups (Shcherbakov and Popov 2002: see their fig. 179), which despite a list of characters supposedly supporting the nodes, can hardly be viewed as a phylogenetic hypothesis. However, Bourgoïn and Campbell (2002), based on unpublished molecular data sets (18S rDNA), also argue that Auchenorrhyncha is paraphyletic, and that the sister group of Heteropteroidea is Cicadomorpha, similar to the proposal of Shcherbakov and Popov (2002).

Yang (2004) proposed a scheme of relationships within Hemiptera based on morphological characters that differ from that presented by Wheeler *et al.* (1993), mainly in that Psylloidea is the sister group of Heteropteroidea, Sternorrhyncha and Cicadomorpha are paraphyletic, and in the internal relationships of the Heteroptera infraorders (Leptopodomorpha, Cimicomorpha, and Pentatomomorpha are in a polytomy, Enicocephalomorpha is the sister group of this clade, and the remaining infraorders forms a basal group in an unresolved polytomy). Nonetheless, several flaws in the analysis, such as choosing *a priori* a “functional” outgroup (within Hemiptera) and not stating how the analysis was done, prevent his discussion of relationships being useful.

Sorensen *et al.* (1995) proposed unnecessary new names for clades already named (e.g., in Wheeler *et al.* 1993), in part, because Auchenorrhyncha was found to be paraphyletic respect to Heteroptera, and also because they wanted to recognize what they considered suborders of Hemiptera. They proposed: Clypeorrhyncha (for extant Cicadomorpha), Archaeorrhyncha (for Fulgoromorpha), Peloridiomorpha (for Coleorrhyncha [Peloridiidae]), Neohemiptera (for Fulgoromorpha + Heteropteroidea), and Prosorrhyncha (for Heteropteroidea) (Table 1). Nonetheless, these names may lead to confusion, in particular if Auchenorrhyncha is not paraphyletic (see below for discussion). Even more, the names Cicadomorpha, Fulgoromorpha, and Heteropteroidea, independently of the outcome of the phylogenetic relationships within Hemiptera, convey a clear meaning already used in Hemiptera literature. Bourgoïn and Campbell (2002) argue in favor of dropping the names proposed by Sorensen *et al.* (1995). Because the ICZN (1999) do not regulate the application of names above family rank, it depends on the researchers to favor the use of certain names.

Vibrational communication is diverse in Hemiptera, with species producing sound in terrestrial as well as aquatic ecosystems, being substrate communication common (Crocroft and Rodríguez 2005). There may be stridulation, tymbal, and

percussional vibration (Claridge 1985; Hoy and Robert 1996; Gogala 2006; Hoch *et al.* 2006; Soulier-Perkins *et al.* 2007). It has been suggested that a tymbal-like communication is synapomorphic for Auchenorrhyncha + Coleorrhyncha + Heteroptera (Hoch *et al.* 2006).

3. Sternorrhyncha

Sternorrhyncha is a monophyletic group (Schlee 1969c; von Dohlen and Moran 1995), which includes four main groups: Psylloidea, Aleyrodoidea, Aphidoidea, and Coccoidea (e.g., Schlee 1969b; Carver *et al.* 1991; von Dohlen and Moran 1995; Bourgoïn and Campbell 2002).

Goodchild (1966) based on the alimentary tract morphology regarded Aphidoidea as the sister group of Coccoidea + (Psylloidea + Aleyrodoidea). Schlee (1969a, 1969b) based on external morphology and male genitalic characters considered two sister groups within Sternorrhyncha: Psylloidea + Aleyrodoidea (Psylliformes), and Aphidoidea + Coccoidea (Aphidiformes). In some analyses Psylloidea is considered the sister group to the rest of Sternorrhyncha (e.g. Campbell *et al.* 1994, 1995). Despite these competing hypotheses, no modern comprehensive phylogenetic analysis is available for the group (but see review of previous phylogenetic hypotheses in Schlee 1969b).

The following subgroups of Sternorrhyncha are presented in alphabetical order because no consensus has been achieved about their phylogenetic relationships.

3a. Aleyrodoidea. The Aleyrodoidea, or whiteflies (Figs. 1A, B, C), includes 1,556 valid species in Aleyrodidae, the only included family, which is subdivided into three subfamilies: Aleurodicinae, Aleyrodinae, and the controversial Udamoselinae (Mound and Halsey 1978; Martin and Mound 2007). Manzari and Quicke (2006) analyzed the phylogenetic relationships of the large and widespread Aleyrodinae, which contains most of the economically important pest species (Byrne and Bellows 1991; Martin and Mould 2007). Their analyses based on pupal case characters (Manzari and Quicke 2006) found that only 45% of the genera with multiple representatives were monophyletic, and that most of the tribes and economically important genera (e.g., *Bemisia* and *Trialeurodes*) are not monophyletic. They did not propose any new taxonomic scheme at the tribal level arguing that the data set used is not enough to provide adequate resolution.

3b. Aphidoidea. Aphidoidea has three families, Phylloxeridae, Adelgidae, and Aphididae, commonly called aphids (Figs. 1D, E), with some 4,500 described species (Remaudière and Remaudière 1997; Blackman and Eastop 2006). Some authors prefer to give them rank of superfamilies (e.g., Shcherbakov and Popov 2002).

Heie (1987) proposed a scheme of relationships for the Phylloxeridae and Aphididae based on morphology of extant and extinct taxa. Von Dohlen and Moran (2000) proposed a phylogenetic scheme for Aphididae using 12S and 16S mtDNA. They found little support and structure at deeper nodes, only recovering recognized tribes but not subfamilies. Martínez-Torres *et al.* (2001), also using a mitochondrial gene, found similar results. Ortiz-Rivas *et al.* (2004) analyzed the relationships of Aphididae based on a single nuclear gene (long wavelength opsin) and the topologies of their analyses were better resolved than in previous analysis. They identified three main clades, with Aphidinae and Lachninae always

monophyletic. The relationships within Aphidinae were examined by von Dohlen *et al.* (2006) with three genes: leucine tRNA, COII mtDNA, and EF1 α .

3c. Coccoidea. The group has about 7300 described species (Miller and Ben-Dov 2006), and the 20 or more families (Figs. 1F, G) are usually divided into two groups: Archaeococcoidea and Neococcoidea (Koteja 1974; Miller and Kosztarab 1979; Gullan and Kosztarab 1997). Gullan and Cook (2007) review the phylogeny and classification of the coccoids in detail.

Miller (1984), Miller and Hodgson (1997), and Foldi (1997) proposed phylogenetic hypothesis based in morphological characters for the group. Nonetheless, their analyses had either a small taxon sampling or poor nodal support that renders the results non-conclusive.

Cook *et al.* (2002) assessed the phylogeny of the Coccoidea using rDNA sequences of the small subunit. They found that the neococcoidea are a monophyletic group and that Pseudococcidae is the sister group to the rest of them. Eriococcidae was retrieved as a paraphyletic group in their analyses, corroborating previous views based on morphology (Cox and Williams 1987). The monophyly and relationships of the archaeococcoids are in doubt (Gullan and Cook 2007).

Two out of three of the largest families of Coccoidea have been assessed for phylogenetic relationships. Downie and Gullan (2004) assessed the phylogeny of the Pseudococcidae using DNA sequences of EF1 α , 28S and 18S. They found that the three major clades of the analysis correspond to the subfamilies Pseudococcinae, Phenacoccinae and Rhizoecinae, with Sphaerococcinae as polyphyletic. Recently, Hardy *et al.* (2008) reassessed the relationships of the Pseudococcidae, using an expanded taxon sampling and adding morphological data. They recognized two main clades to which they apply the names Phenacoccinae and Pseudococcinae, including in the former the Rhizoecini.

Morse and Normark (2006) studied the phylogenetic relationships of Diaspididae using EF1 α and 28S rDNA sequences. Most of the traditional groups recognized in classifications were broadly recovered, although none of them was strictly monophyletic. The remaining family, Coccidae, has never been subject of a comprehensive phylogenetic analysis. Qin and Gullan (1995) evaluated the relationships of one of the subfamilies, Ceroplastinae (Coccidae), finding that most of the genera were not monophyletic, and that most of the species should be grouped into the genus *Ceroplastes*.

3d. Psylloidea. Psylloidea (Fig. 1H) has more than 3000 described species (Hodkinson and Casson 1991; Hollis 2004; Burckhardt 2005). The classification of this group of sap-sucking insects has been extremely artificial (Burckhardt and Lauterer 1989). White and Hodkinson (1985) first proposed a phylogenetic scheme of the Psylloidea using mostly nymphal characters. In their phylogenetic analysis they regard Psyllidae as the sister group of all other psylloids. Of the eight families recognized by them, two, Aphalaridae and Spodilyaspididae, were accepted in their classification as paraphyletic following an "evolutionary" approach (White and Hodkinson 1985: 264). Ouvrard (2002) reassessed the relationships of the Psylloidea using morphological characters of the adult thorax and 18S rRNA sequences. His results differ from those of White and Hodkinson (1985), although it is difficult to compare and interpret them because Ouvrard did not present a list of characters, matrix, or list of species used in his analysis.

Nonetheless, he concluded that Aphalaridae and Psyllidae are paraphyletic.

Recently, Ouvrard *et al.* (2008) identified potential morphological synapomorphies for Psylloidea in relation to the wing base structure: absence of subalare, a median notal wing process, and of the anterior arm of the third axillary sclerite; no articulation with the second axillary sclerite; a weakly-sclerotized third axillary sclerite; and presence of a two-horned basalare. Hodkinson (1989) provided a review of the faunal elements present in the Neotropical region.

4. Auchenorrhyncha

Auchenorrhyncha has been traditionally divided in two main groups, Cicadomorpha and Fulgoromorpha (e.g., Hennig 1969, 1981; Carver *et al.* 1991). Nonetheless, it has been extensively debated if Auchenorrhyncha is a monophyletic group or not. Goodchild (1966) based on the morphology and histology of the digestive tract considered Auchenorrhyncha as paraphyletic because he considered two groups: Fulgoromorpha + Heteroptera and Cicadomorpha + Sternorrhyncha, with the Coleorrhyncha in a polytomy with both. Hamilton (1981) regarded Auchenorrhyncha paraphyletic because he considered Fulgoromorpha as sister group to Cicadomorpha + Aphidiformes, although he also considered Homoptera as monophyletic. Wootton and Betts (1986) using characters of wing morphology also doubted the monophyly of Auchenorrhyncha. Bourgoïn proposed that Auchenorrhyncha is paraphyletic, based on head morphology (Bourgoïn 1986a, 1986b), and on male (Bourgoïn and Huang 1990) and female genitalia (Bourgoïn 1993), because Fulgoromorpha is apparently more related to Heteropteroidea than to Cicadomorpha. Nonetheless, the particular aristate antennae (e.g., Grimaldi and Engel 2005) and evidence from the fore wing base sclerites (Yoshizawa and Saigusa 2001), suggest that Auchenorrhyncha is in fact monophyletic.

Much of the recent evidence for the paraphyly of Auchenorrhyncha come from molecular sets alone. Campbell *et al.* (1995) using 18S rDNA sequences found a polytomy among Fulgoromorpha, Cicadomorpha, and Heteropteroidea when using the full dataset (their fig. 3). When using a restricted data set (homoplasious sites removed), either Fulgoromorpha or Cicadomorpha become sister groups of Heteropteroidea (Campbell *et al.* 1995, their fig. 4), but when Peloridiidae was removed from this restricted data set, Fulgoromorpha become the sister group of Heteroptera. Sorensen *et al.* (1995) also using 18S rDNA found that Auchenorrhyncha is not monophyletic. Despite their small dataset, they found that Auchenorrhyncha is either paraphyletic or polyphyletic respect to Heteroptera. Bourgoïn *et al.* (1997) in an analysis of the Fulgoromorpha found that this group is the sister group to Heteroptera, rendering Auchenorrhyncha paraphyletic. They also found that Cicadomorpha was not monophyletic if Psyllidae was included as part of the outgroup, showing that their analysis is sensitive to taxon sampling. Ouvrard *et al.* (2000) found a polytomy among Fulgoromorpha, Cicadomorpha, and Heteropteroidea, similar to the results of Campbell *et al.* (1995). Urban and Cryan (2007) using an extended data set of four genes and 83 species of Fulgoromorpha found a monophyletic Auchenorrhyncha, which was the sister group of Heteroptera.

4a. Cicadomorpha. Within this group three superfamilies are recognized: Cercopoidea (spittlebugs or froghoppers) (Figs.



Figure 1. Habitus illustrations of different groups of Hemiptera (not at the same scale). **A.** *Trialeurodes vaporariorum* (Westwood) (Aleyrodidae). **B.** *Bemisia argentifolii* Bellows & Perring (Aleyrodidae). **C.** *Aleurodicus dugesii* Cockerell (Aleyrodidae). **D.** *Adelges cooleyi* (Gillette) (Adelgidae). **E.** *Aphis nerii* Fonscolombe (Aphididae). **F.** *Pseudococcus longispinus* (Targioni-Tozzetti) (Pseudococcidae). **G.** *Aspidaspis artostaphyli* Cockerell & Robbins (Diaspididae). **H.** *Russelliana solanicola* Tuthill (Psyllidae). Peru. Lateral and dorsal views. **I.** *Sphenorhina melanoptera* (Germar) (Cercopidae). Ecuador. **J.** *Prosapia bicincta* (Say) (Cercopidae). Panama. **K.** Cicadidae. Panama. Recently molted adult and exuviae. **L.** *Cladonota* sp. (Membracidae). Panama. **M.** *Heteronotus* sp. (Membracidae). Peru. **N.** *Ferrariana trivittata* (Signoret) (Cicadellidae). Panama. **O.** *Platygonia spatulata* (Signoret) (Cicadellidae). Panama. **P.** *Proconia* sp. (Cicadellidae). Ecuador.

II, J), Cicadoidea (Fig. 1K), and Membracoidea (leafhoppers and treehoppers) (Figs. 1L, M, N, O, P), with approximately 35.000 described species (Cryan 2005; Dietrich 2005).

The monophyly of Cicadomorpha has never been doubted based on morphological characters; nonetheless, all possible relationships have been proposed among its superfamilies: Membracoidea has been proposed the sister group of Cercopoidea + Cicadoidea (Campbell *et al.* 1995; Ouvrard *et al.* 2000; Bourgoïn and Campbell 2002); Cercopoidea as sister group of Membracoidea + Cicadoidea (Evans 1963; Hamilton 1999); and Cicadoidea as sister group of Membracoidea + Cercopoidea (Hamilton 1981; Sorensen *et al.* 1995; von Dohlen and Moran 1995). More recently, Cryan (2005) criticized previous analyses among the superfamilies arguing both poor taxon and data sampling. He proposed a phylogeny of Cicadomorpha based on three nuclear gene sequences. He found that including all data sets is better than analyzing individual gene data sets, and that 18S is a relatively poor molecular marker for recovering the relationships among superfamilies. Although some morphological evidence apparently placed Membracoidea as a derived taxon (e.g., Hamilton 1981), independent character analysis supports Membracoidea as a basal Cicadomorpha group, for instance, Liang and Fletcher (2002) with antennal characters, and Rakitov (2002) with structural characters of brochosomes, proteinaceous particles secreted by glandular regions of the Malpighian tubules of almost all cicadellids.

Membracoidea currently includes the extant families Aetalionidae, Cicadellidae, Melizoderidae, Membracidae, and Myerslopiidae (Hamilton 1999; Dietrich 2005).

Dietrich and Deitz (1993) analyzed the relationships of the Membracoidea using morphological characters, focusing mostly in non-cicadellid taxa (Aetalionidae and Membracidae). In their analysis Cicadellidae is monophyletic and the sister group of (Melizoderidae + (Aetalionidae + Membracidae)). Dietrich and Deitz (1993) listed as synapomorphies for Cicadellidae: the mesonotum exposed posteriorly, the labium not reaching the metathoracic coxae, m-cu1 crossvein present, metatibia with distinct long setae, tarsomere I of hind leg without cucullate setae, sternum IX and subgenital plate not fused, and abdominal tergum with divided canthae; all of which are homoplastic characters in their analysis.

Hamilton (1999) presented a phylogeny of the extinct and extant families of Membracoidea, and placed Myerslopiidae as the sister group of the remaining families. In Dietrich *et al.* (2001b) analysis Myerslopiidae grouped with Cicadoidea taxa, whereas in Cryan's (2005) Myerslopiidae was recovered as the sister group to all remaining Membracoidea. Hamilton (1999) also considered Ulopidae as a separate group from Cicadellidae, and placed Cicadellidae as sister group of (Ulopidae + (Aetalionidae + Membracidae)), rendering Cicadellidae paraphyletic with respect to Membracidae. This scheme was first supported by Dietrich *et al.* (2001b) based on the analysis of 28S rDNA sequences, where the clade including Ulopinae and Megophthalminae appears as more closely related to Aetalionidae + Membracidae; and posteriorly by Cryan (2005).

Within Membracoidea, the Membracidae has received much attention on its suprageneric classification with recent phylogenetic analysis at the subfamily level using morphological characters (Dietrich and Deitz 1993; Dietrich *et al.* 2001a), molecular ones (Cryan *et al.* 2000), or in combined analysis (Cryan *et al.* 2004). The relationships within

subfamilies, tribes, or genera have also been assessed for a number of taxa (e.g., Aconophorini: Dietrich and Deitz, 1991; Centrotinae: Wallace and Deitz 2004, 2005; Darnini: Roy *et al.* 2007; Membracinae: Lin *et al.* 2004; Microcentrini: Cryan *et al.* 2003; Nicomiinae: Albertson and Dietrich 2005).

The speciose family Cicadellidae, commonly called leafhoppers and sharpshooters (Membracoidea), has more than 22.000 described species worldwide and 5.000 species in the Neotropical region (Freytag and Sharkey 2002), and Cicadellidae has about 36 subfamilies worldwide (Oman *et al.* 1990; Dietrich 2004). Phylogenetic hypotheses based on morphology were first proposed by Ross (1957), then by Hamilton (1983) and Dietrich (1999). Dietrich *et al.* (2001b) was the first attempt to recover the phylogeny of most cicadellid subfamilies and tribes using 28S rDNA sequences, and found that most of them were not monophyletic. Within Cicadellidae some groups had been subject of analyses (e.g., Evacanthinae: Dietrich 2004; Cicadellinae: Takiya *et al.* 2006; Deltocephalinae: Zahniser and Dietrich 2008).

The phylogenetic relationships within Cicadoidea have rarely been addressed at the family level, except in a few instances, with characters used in traditional classifications (e.g., Chou *et al.* 1997). Recently, Moulds (2005) proposed a phylogenetic hypothesis for the Cicadoidea with an extended morphological data set in which he recognized two families, Tettigarctidae and Cicadidae, the latter with three subfamilies. The only higher phylogenetic proposal for Cercopoidea is that of Cryan (2005). He shows that Machaerotidae + Clastopteridae is the sister group of Cercopidae + Aphrophoridae. Cryan's (2005) analysis supports a monophyletic Cercopidae, and although it shows a monophyletic Aphrophoridae, he doubts of its monophyly based on additional unpublished data. Interestingly, the recently described family Epygidae (Hamilton 2001), was found nested within the Aphrophoridae.

4b. Fulgoromorpha. This group, often referred as Fulgoroidea in the literature, and commonly named planthoppers (Figs. 2A, B, C), has more than 9.000 described species, and about 20 families (O'Brien and Wilson 1985).

Several hypotheses of relationships have been proposed for this group based on morphological characters, all of which support a monophyletic Fulgoromorpha (Muir 1923; Asche 1987; Emeljanov 1990; Bourgoïn 1993; Chen and Yang 1995). In all but one of these hypotheses the Tettigometridae is considered the sister group to the rest of the fulgoroids, with various levels of resolution and clade composition for the remaining taxa (reviewed by Urban and Cryan 2007). Only Bourgoïn (1993) considered the Tettigometridae as a derived taxon using morphological characters.

Bourgoïn *et al.* (1997) assessed the relationships of a small sample of fulgoromorph taxa, targeting in particular the position of Tettigometridae. They found that Fulgoromorpha is a monophyletic clade and that the relictual Tettigometridae is not basal as previously thought. Yeh *et al.* (1998) analyzed a restricted set of Fulgoromorpha with partial sequences of 16S rDNA. Yeh *et al.* (2005) expanded their dataset of 16S sequence data and analyzed the fulgoroids with neighbor joining algorithms. They obtained similar results to that of Bourgoïn *et al.* (1997) in placing Cixiidae and Delphacidae as the most basal groups, and in having the Tettigometridae in a derived position. Yeh and Yang (1999) using 28S rDNA sequences analyzed the relationships of some fulgoromorphs. They reached similar conclusions to Yeh *et al.* (2005). More recently,




Figure 2. Habitus illustrations of different groups of Hemiptera (not at the same scale). **A.** *Pyrops* sp. (Fulgoridae). Borneo. **B.** Flatidae. Peru. **C.** *Acanalonia* sp. (Acanaloniidae). Panama. **D.** *Peloridora kuscheli* China (Peloridiidae). Chile. **E.** *Systelloderes* sp. (Enicocephalidae). Costa Rica. **F.** *Nannocoris* sp. (Schizopteridae). Ecuador. **G.** *Rheumatobates crassifemur schroederi* Hungerford (Gerridae). Brazil. **H.** *Rhagovelia* sp. (Veliidae). Colombia. **I.** *Cryphocricos latus* Usinger (Naucoridae). Costa Rica. **J.** *Gelastocoris* sp. (Gelastocoridae). Costa Rica. **K.** *Saldula pallipes* (Fab.) (Saldidae). USA.

Urban and Cryan (2007) assessed the phylogenetic relationships of 71 species of Fulgoromorpha and nine outgroup species, with 18S, 28S, Histone 3, and *Wingless* nucleotide sequences. They obtained Cixiidae and Delphacidae as the sister group of the rest. Many of the families were recovered as monophyletic, but some were poly- or paraphyletic, mostly in agreement with previous morphological hypotheses.

5. Heteropteroidea

Heteropteroidea is the clade containing Coleorrhyncha + Heteroptera (Schlee 1969d; Hennig 1969, 1981; Kristensen 1975). Zrzavý (1992) changed the term coined by Schlee (1969d), Heteropteroidea, to Heteropteroidea, to avoid confusion with a superfamily rank. Heteropteroidea is supported by the following synapomorphies: four antennal segments, body with a flattened dorsal area to receive the wings (Hennig 1969, 1981), anal veins of forewing fused forming a Y-shaped vein (Wootton 1965), last antennal segments larger than proximal ones, abdominal tergites flat with sharp connexivum, abdominal spiracles directed ventrad, anal cone forming a single, exposed element (Schlee 1969d), and wing coupling mechanism of heteropterous type (China 1962). This clade is further supported by a few 18S rDNA characters (Wheeler *et al.* 1993).

6. Coleorrhyncha

Coleorrhyncha is a small group of Hemiptera that comprises 13 extant genera and 25 species in the only extant family Peloridiidae (Fig. 2D) (China 1962; Evans 1981). The distribution is restricted to the Southern Hemisphere and is considered relictual, being found in Eastern Australia, New Caledonia, New Zealand, Chile and Argentina (Patagonia) (Evans 1981; Estévez and de Remes-Lenicov 1989; Burckhardt and Agosti 1991; Burckhardt and Cekalovic 2002). Extant species are associated with *Nothofagus* forests, and little is known about their biology, except that they live on mosses (Hacker 1932; Evans 1981), and are found on lichens, hepatics, and sphagnum bogs (Evans 1981). The South American fauna includes three monotypic genera, *Pantinia*, *Peloridium*, and *Kuscheloides*, as well as *Pelorida* with three species (China 1962; Evans 1981), although some undescribed taxa may still exist (e.g., Burckhardt and Agosti 1991).

Coleorrhyncha was traditionally included as a member of the "Homoptera" (e.g., Myers and China 1929; China 1962; Evans 1963), but it is now considered the sister group of Heteroptera due to compelling morphological and molecular evidence (Schlee 1969d; Hennig 1969, 1981; Wheeler *et al.* 1993; Sweet 1996; Ouvrard *et al.* 2000). Peloridiidae has been considered a monophyletic group because of its particular morphology (e.g., Evans 1963). Surprisingly, Ouvrard *et al.* (2000) found a paraphyletic Peloridiidae in some analysis with 18S rRNA, but recovered it as monophyletic after eliminating all homoplasious sites.

Popov and Shcherbakov (1996) revised the fossil Peloridiidae and proposed a scheme of relationships among the extant members. Despite presenting a "matrix" of characters and a tree, these certainly do not represent a cladistic analysis of relationships (see also Schuh 1997).

7. Heteroptera

Heteroptera is defined by the following putative synapomorphies: presence of gula, presence of metathoracic scent apparatus, abdominal dorsal scent glands (functional in

nymphs), and strong reduction of the tentorium (Carayon 1971; Cobben 1978; Hennig 1969, 1981; Kristensen 1975). Although metathoracic scent glands are not easy to observe in Enicocephalomorpha, there is evidence for its presence (Carayon 1962, 1971; Cobben 1978). The presence of a frenum, which helps to attach the posterior claval margin of the forewing to the lateral margin of the scutellum, is also a potential synapomorphy for Heteroptera (Štys 1998). An open rhabdom structure in the ommatidia of Heteroptera is considered a synapomorphy for Heteroptera, in contrast to the fused rhabdoms in Auchenorrhyncha and Coleorrhyncha (Fischer *et al.* 2000). The absence of the tegula in the forewing is another potential synapomorphy for Heteroptera (Yoshizawa and Saigusa 2001).

Latreille (1810) was the first to recognize Heteroptera as it is conceived nowadays. Dufour (1833) first recognized major groups within Heteroptera. He based his scheme on the habitat occupied by the bugs: Geocorises (terrestrial bugs), Hydrocorises (water bugs), and Amphibicorises (superficial water bugs). Dufour's (1833) classification remained in use until the mid-20th century (e.g., China and Miller 1959). Reuter (1910) proposed a phylogenetic scheme for Heteroptera based on external morphological characters. Štys and Kerzhner (1975) and Schuh (1986) provided a review of the historical proposals of classification among Heteroptera groups that were not based on cladistic methodology.

Leston *et al.* (1954) recognized monophyletic groups within the terrestrial Heteroptera (= Geocorises) of Dufour (1833), proposing and listing the characters defining Cimicomorpha and Pentatomomorpha. They typified these suprageneric names based on genera. Štys and Kerzhner (1975) recognized seven infraorders within Heteroptera, and standardized their names. They followed Leston *et al.* (1954) in applying the -morpha suffix to their infraordinal names.

Schuh (1979) first proposed a phylogenetic scheme for the infraorders of Heteroptera reanalyzing the data of adults and nymphs presented by Cobben (1978). Schuh (1979) considered Enicocephalomorpha as the sister group of the remaining Heteroptera, with Leptopodomorpha + Nepomorpha as the sister group of Cimicomorpha + Pentatomomorpha. Zrzavý (1992) regarded Enicocephalomorpha the sister group of Dipsocoromorpha + Gerromorpha, together forming a basal heteropteran clade, but presented the relationships among Nepomorpha, Leptopodomorpha, and Cimicomorpha + Pentatomomorpha as unresolved. Wheeler *et al.* (1993) used 18S rDNA sequences and morphological characters to resolve the relationships of the Heteroptera infraorders. Their results mostly agree with Schuh's (1979) scheme. They only disagree in the position of Leptopodomorpha, which they consider the sister group of Cimicomorpha + Pentatomomorpha.

Cimicomorpha and Pentatomomorpha are considered sister groups (Schuh 1979; Wheeler *et al.* 1993). Beside the characters mentioned by Schuh (1979) and Wheeler *et al.* (1993), they also share a V-pattern in the rhabdomeres of the ommatidia, further supporting their relationship (Fischer *et al.* 2000).

Schuh (1986) summarized the phylogenetic hypotheses available at that time for all Heteroptera infraorders. The following sections update the information provided by him.

7a. Enicocephalomorpha. This relatively small group of true bugs contains approximately 450 described species (Schuh and Slater 1995). Enicocephalomorpha contains two families: Aenictopecheidae and Enicocephalidae (Štys 1989). This



Figure 3. Habitus illustrations of different groups of Hemiptera (not at the same scale). **A.** *Diactor* sp. (Coreidae). Panama. **B.** *Dinidor* sp. (Dinidoridae). Ecuador. **C.** *Edessa rufomarginata* (DeGeer) (Pentatomidae). Panama. **D.** *Lygaeus* sp. (Lygaeidae). Peru. **E.** *Harpactor* sp. (Reduviidae). Peru. **F.** *Zelurus* sp. (Reduviidae). Peru. **G.** *Agriocoris flavipes* (Fab.) (Reduviidae). Ecuador. **H.** *Zanchius* sp. (Miridae). Nepal. **I.** *Zanchius* sp. (Miridae). Nepal. **J.** *Pseudoloxops* sp. (Miridae). Nepal.

monophyletic clade was at one point considered part of the Reduviidae (Cimicomorpha) (e.g., Usinger 1943), but is now considered the sister group of the remaining Heteroptera (e.g., Schuh 1979; Wheeler *et al.* 1993).

Wygodzinsky and Schmidt (1991) provided a preliminary phylogenetic scheme of the New World taxa. They placed Aenictopecheidae (as subfamily) as sister group to the remaining taxa. No comprehensive cladistic phylogenetic analysis has been carried out on this group of Heteroptera.

7b. Dipsocoromorpha. Five families (Fig. 2F) are included in the most recent classification of the infraorder (e.g., Schuh and Slater 1995). Štys (1985) termed Euheteroptera the clade containing Dipsocoromorpha + (Gerromorpha + (Nepomorpha + (Leptopodomorpha + (Cimicomorpha + Pentatomomorpha))). The monophyly of this infraorder was proposed by Štys (1983) based on the structure of the male genitalia and legs. Nonetheless, these characters are homoplastic within the infraorder (see Wheeler *et al.* 1993). Despite abundant morphological documentation (e.g., McAtee and Malloch

1925; Hill 1987; Rédei 2007; Štys 1970; Wygodzinsky 1950), no modern phylogenetic analysis is available for this rarely collected group of true bugs.

7c. Gerromorpha. There are approximately 1900 described species in this group (Figs. 2G, H) (Andersen and Weir 2004b). Species of this infraorder inhabit the surface of marine or freshwater systems, belonging to this clade the only true marine insect, the genus *Halobates* (Andersen and Weir 1994, 1999). The single most important reference for the Gerromorpha, or semiaquatic bugs, is that of Andersen (1982), in which he reviewed the biology, the morphology, and phylogeny of the group. An update for the phylogenetic hypotheses of the infraorder was provided by Spence and Andersen (1994), as well as a review of their ecology and behavior. Štys (1985) termed Neoheteroptera the clade containing Gerromorpha + (Nepomorpha + (Leptopodomorpha + (Cimicomorpha + Pentatomomorpha))).

Beside the characters listed as synapomorphic for Gerromorpha (Andersen 1982; Wheeler *et al.* 1993; Schuh

Table 1. Equivalences of terms for the higher categories of Hemiptera. Synonyms in parentheses are terms created by Sorensen *et al.* (1995). See text for details.

Major groups	Subordinate groups
Sternorrhyncha	Aleyrodoidea Aphidoidea Coccoidea Psylloidea
Auchenorrhyncha	Cicadomorpha (= Clypeorrhyncha) Fulgoromorpha (= Archaeorrhyncha)
Heteropteroidea (= Prosorrhyncha)	
Coleorrhyncha (= Peloridiomorpha)	Peloridiidae
Heteroptera	Enicocephalomorpha Dipsocoromorpha Leptopodomorpha Gerromorpha Nepomorpha Cimicomorpha Pentatomomorpha

and Slater 1995), it can be added that the eyes dorsally have the ommatidia with a rhabdom pattern of R7 and R8 in tandem (Fischer *et al.* 2000). Andersen (1998) reviewed fossil species of Gerromorpha, and mapped them on his previous phylogeny of the group (Andersen 1982), in which he placed the extinct subfamily Electrobatinae (Gerridae).

Muraji and Tachikawa (2000) analyzed the relationships among the families of Gerroidea (Gerridae, Hermatobatidae, and Veliidae), including 30 species of the superfamily and two other species of Gerromorpha as outgroups, using 16S rDNA and 28S rDNA. Sequence data were analyzed with parsimony and neighbor-joining algorithms. They recovered a monophyletic Gerroidea, and some other clades within the superfamily already supported by morphological characters.

Andersen and Weir (2004b) reanalyzed the relationships of Gerromorpha using 56 morphological characters with a numerical cladistic approach. The topology presented is mostly similar to that of Andersen (1982), except that the relationships of the Veliidae subfamilies Rhagoveliinae, Perittopinae, and Veliinae are unresolved, as well as most of the subfamilies of Gerridae. They also presented arguments for the monophyly of Veliidae and discussed the inclusion of *Ocellovelia* in this family.

Damgaard *et al.* (2005) reanalyzed the relationships of Gerromorpha using the same two gene regions of Muraji and Tachikawa (2000), but with an expanded taxon sampling, and adding morphological characters from Andersen (1982). They assessed the effects of alignment and taxon sampling in their analysis, showing that there is little node stability with different parameters. They recovered as monophyletic the superfamily Gerroidea, and Gerridae, but Veliidae as paraphyletic. They also found support for the Halobatinae (Gerridae) which had ambiguous morphological support, and that *Ocellovelia* was placed as sister group of "Veliidae" + Gerridae only with molecular characters. Recent systematic work has focused on the relationships within families (e.g., Andersen 1999), subfamilies (e.g., Andersen 1995, 2000), and genera, in parti-

cular within Gerridae (e.g., Damgaard *et al.* 2000a, 2000b; Damgaard and Sperling 2001; Damgaard and Cognato 2005).

7d. Nepomorpha. This infraorder contains about 2000 species in eleven families (Figs. 2I, J) (Štys and Jansson 1988; Hebsgaard *et al.* 2004). The group corresponds to the Hydrocorisae of Dufour (1833). Štys (1985) termed Panheteroptera the clade containing Nepomorpha + (Leptopodomorpha + (Cimicomorpha + Pentatomomorpha)).

Wheeler *et al.* (1993) presented molecular evidence for a monophyletic Nepomorpha, rejecting a basal position within Heteroptera as argued by some authors (e.g., Reuter 1910; Mahner 1993). The relationships within the clade, nonetheless, were unresolved probably due to the limited taxon sampling. Mahner (1993) provided a phylogenetic analysis for Nepomorpha and all the subordinate groups. His scheme is similar to that of Rieger (1976) in the placement of Nepoidea [i.e., (Nepidae + Belostomatidae), see Schuh and Slater 1995: 110] as sister group to the remaining Nepomorpha, and in having as the most derived clade the Notonectoidea [i.e., (Notonectidae + (Helotrephidae + Pleidae))]. Mahner's (1993) scheme differs from Rieger's (1976) in the position of Corixidae, which is considered the sister group of (Ochteroidea + (Naucoroidea + Notonectoidea)); whereas in Rieger's scheme the position of the Ochteroidea and Corixidae are inverted [i.e., (Ochteroidea + (Corixidae + ("Naucoroidea" + Notonectoidea)))]]. In addition, Rieger (1976) data suggest a paraphyletic Naucoroidea, whereas Mahner (1993) treat this superfamily as monophyletic, but without placing the Potamocoridae in his analysis.

Hebsgaard *et al.* (2004) proposed a phylogenetic hypothesis for the Nepomorpha using both molecular (16S and 28S rDNA) and morphological characters. They found support for the monophyly of all Nepomorpha families, and superfamilies except Naucoroidea (i.e., Naucoridae, Aphelocheiridae, and Potamocoridae). Their analysis place Nepoidea as the most basal group, and Corixidae as sister group remaining Nepomorpha, similar to the proposal of Mahner (1993).

7e. Leptopodomorpha. This infraorder contains four families (Fig. 2K) and about 300 described species, nearly all of them in Saldidae (Schuh *et al.* 1987). Schuh and Slater (1995) listed the synapomorphies for the infraorder. An additional synapomorphy for Leptopodomorpha is the shape of the rhabdom resembling a “5” pattern on a dice (Fischer *et al.* 2000).

Schuh (1986) reviewed the phylogenetic proposals and classification scheme for the infraorder. In the cladistic analysis of Schuh and Polhemus (1980) the relationship of the higher groups is ((Saldidae + Aepophilidae) + (Omaniidae + Leptopodidae)). Polhemus (1985) presented a cladistic analysis of the Leptopodomorpha, based on his PhD dissertation data, in which the relationships of the higher groups are (Leptopodidae + (Omaniidae + (Saldidae + Aepophilidae))), differing from that of Schuh and Polhemus (1980) in the position of the Omaniidae. Polhemus (1985) presented in addition phylogenetic hypotheses for the subfamilies and tribes of Saldidae, for the genera of Saldini, and for the genera of the large Saldoidini. No more recent hypotheses with additional morphological or molecular characters has been proposed.

7f. Pentatomomorpha. This infraorder contains about 15,000 described species (Figs. 3A, B, C, D) (Henry 1997, Schuh and Slater 1995). Leston *et al.* (1954) first recognized Pentatomomorpha as a monophyletic group. It included those groups with abdominal trichobothria, the Trichophora of Tullgren (1918), plus Aradoidea (Aradidae + Termitaphididae). Although Sweet (1996) has given infraordinal status to Aradoidea, ranking it differently does not modify its placement as the sister group of the remaining Pentatomomorpha, as argued by Henry (1997). Sweet (2006) has even further suggested that the position of the Aradoidea may lay outside the Leptopodomorpha, Cimicomorpha, and Pentatomomorpha, which clearly contradicts the available morphological and molecular evidence (e.g., Wheeler *et al.* 1993; Grazia *et al.* in press).

Henry (1997) assessed the relationships of the infraorder in a morphological cladistic analysis. He included 53 taxa, using families, subfamilies, and tribes—in particular of the Lygaeoidea (*sensu* Schuh and Slater 1995)—as terminals using a *ground plan* approach. He chose Aradoidea as the outgroup for his analyses. He also included the Pentatomoidea as a single terminal taxon in the analysis providing evidence for its monophyly, but showing the difficult position of Thaumastellidae, usually included in this group. Henry’s main interest was to elucidate the relationships among the Lygaeoidea and to test the monophyly of this group, while looking for a sister group relationship for the Berytidae. In his analyses Pentatomoidea is the sister group of the remaining Pentatomomorpha, i.e., ((Coreoidea + Pyrrhocoroidea) + (Idiostoloidea + Lygaeoidea *sensu stricto*)). He found that Lygaeoidea is paraphyletic with respect to the Berytidae, Colobathristidae, and Piesmatidae, which were nested within Lygaeidae *sensu lato* (e.g., Schuh and Slater 1995). Henry raised several of the previous subfamilies and tribes of the Lygaeidae (*sensu lato*) to family status proposing a new classification for Pentatomomorpha reflecting the phylogenetic pattern found.

Li *et al.* (2005) analyzed the relationships of Pentatomomorpha groups using partial 18S and COI gene sequences. About half of the terminal taxa used in their analysis had sequences from GenBank produced by Wheeler and Schuh. They used parsimony, maximum likelihood, and distance methods to analyze the data. Sequence data were aligned using

the default parameters of Clustal and “by eye”, and variable regions excluded from the analyses. As results, they found support for a monophyletic Pentatomomorpha and Trichophora in all analyses. Other groups, nonetheless, are in disagreement with previous morphological hypotheses. In the combined parsimony analysis, Lygaeoidea (*sensu* Henry 1997) is paraphyletic with respect to Berytidae and Piesmatidae, the latter two nested within a paraphyletic Pyrrhocoroidea. Having a paraphyletic Lygaeoidea may not be surprising because of the limited taxon sampling within the superfamily, in which only four out of 15 groups (as in Henry 1997) were included. The Coreoidea was found paraphyletic in this combined parsimony analysis. Pentatomoidea was as well paraphyletic because one of the species of Pentatomidae was the sister group of the remaining Pentatomomorpha species less the Aradoidea. In the maximum likelihood combined analysis, the Pentatomoidea was monophyletic. All of the other groups, nonetheless, in this analysis were recovered again as poly- or paraphyletic, with a more scrambled topology compared to that of the parsimony analysis. Despite these surprising findings, Li *et al.* (2005) failed to adequately discuss their results in light of previous morphological hypotheses (e.g., Henry 1997).

More recently, Grazia *et al.* (in press) analyzed the relationships within Pentatomoidea. They employed 135 terminals, using 57 morphological characters, and sequence data from 18S rDNA, 16S rDNA, 28S rDNA, and COI gene regions. They found that Urostylidae *sensu lato* is not monophyletic, and that Urostylidae *sensu stricto* is the sister group of the remaining Pentatomoidea, whereas Saileriolinae should be elevated to family status and placed as the sister group of the remaining Pentatomoidea less Urostylidae *sensu stricto*. Cydnidae is monophyletic only in the morphological analysis, but in the molecular and combined ones it is paraphyletic, in particular because of Corimelaenidae and Thaumastellidae. Parastrachiinae, formerly considered a family different from the Cydnidae (e.g., Sweet and Schaefer 2002), forms a monophyletic group with Corimelaenidae. Pentatomidae is a strongly supported monophyletic group, and based on morphology it also includes Aphyliinae and Cryptocorinae, sometimes treated as families distinct from Pentatomidae (e.g., Schuh and Slater 1995; Packauskas and Schaefer 1998). The expanded scutellum that occurs in many pentatomoid groups is probably a convergent feature as suggested by their analyses. Clearly, as the authors point out, a broader taxon sampling of Pentatomoidea, and additional and more complete data sets will help clarify the status of several equivocal taxa, in particular of the Cydnidae.

7g. Cimicomorpha. After Leston *et al.* (1954) proposed the infraorder, only a few phylogenetic schemes were presented for the groups included. Schuh (1986) reviewed and discussed the schemes of relationships within Cimicomorpha proposed by Kerzhner (1981) and Schuh (1979). He also presented the cladogram of an unpublished analysis of Cimicoidea relationships (see Schuh 1986: 79). Schuh’s (1979) analysis differs mainly from that of Kerzhner (1981) in the placement of the Reduvioidea, which Schuh considered to be the sister group to the remaining Cimicomorpha, contrary to Kerzhner’s assertion of being a derived clade.

Schuh and Štys (1991) analyzed for the first time the relationships within Cimicomorpha using cladistic methods. They provided a historical review of the terminal taxa used and arguments for the monophyly of each. Schuh and Štys

(1991) used in their analysis a *ground plan* approach for certain characters, and families as terminal taxa instead of species. Their analysis shows that Reduvidae (i.e., Reduviidae + Pachynomidae) is the sister group to the remaining Cimicomorpha, agreeing with the hypothesis of Schuh (1979), and that Velocipedidae is the sister group of ((Microphysidae + Joppeicidae + Miroidea) + (Naboidea + Cimicoidea)).

Schuh *et al.* (in press) reanalyzed the relationships of Cimicomorpha with new characters. The taxon sampling comprised 92 taxa and eight outgroups, using species instead of composite taxa (vs. Schuh and Štys 1991). They prepared a morphological matrix of 73 characters coded for all taxa, and sequence data from 16S rDNA, 18S rDNA, 28S rDNA, and COI, coded for 83 taxa. They also included morphological and sequence data for the recently described family Curaliidae from the Eastern United States (Schuh *et al.* 2008). Their results support a monophyletic Geocorisae (i.e., Pentatomomorpha + Cimicomorpha), but not a monophyletic Cimicomorpha (including Thaumastocoridae) in all analyses. The Thaumastocoridae was found to be polyphyletic when added the molecular data, with the Thaumastocorinae placed as the sister group of Pentatomomorpha, and the Xylastodorinae nested within Cimicomorpha, although their placements are ambiguous. They found that Reduvidae is monophyletic and nested within Cimicomorpha. Schuh *et al.* (in press) modified the concept of Cimiciformes to include Joppeicidae, Microphysidae, Velocipedidae, and Curaliidae, the first two thus removed from Miriformes of Schuh and Štys (1991). The monophyly of Cimiciformes is supported with several morphological and molecular data. The new circumscription of the Miriformes includes Miridae and Tingidae only (=Miroidea), because the monophyly of Thaumastocoridae was found to be ambiguous (see also Schuh *et al.* 2006).

The two largest groups within Cimicomorpha, Miridae and Reduviidae have been subject of cladistic analyses. Schuh (1974, 1976) proposed different phylogenetic hypotheses of relationship for the Miridae, which were discussed by Schuh (1986). Schuh *et al.* (in press) found a monophyletic Miridae, but with the suprageneric relationships in little agreement with previous morphological hypotheses. Extensive taxon sampling, and additional morphological and molecular characters, may help resolve these incongruences. No other recent attempt of elucidating the higher relationships of the Miridae has been carried out.

Recently, Weirauch (2008) for the first time presented a comprehensive morphological cladistic analysis of Reduviidae using 162 characters and 75 ingroup taxa. She found that Reduviidae is monophyletic, and that Pachynomidae is its sister group, confirming previous views (Schuh 1979; Schuh and Štys 1991). She also found that Hammacerinae is the sister group of the remaining Reduviidae; and the monophyly of the Phymatinae complex, which occupies a basal position, and comprises Centrocneminae, Elasmodeminae, Holoptilinae, and Phymatinae, with Phimophorinae probably included in this clade (not included in her analysis). Other findings show that the Harpactorinae exhibit a relatively basal position; that Ectrichodiinae + Tribelocephalinae forms a monophyletic group; that Salyavatinae is paraphyletic with respect to Sphaeridopinae and together they form a monophyletic group; the Reduviinae is clearly paraphyletic, and the Triatominae is monophyletic, with some of these "Reduviinae" as sister groups. Future phylogenetic work including molecular markers may test some of the hypotheses presented by Weirauch.

Economic importance

Most of the groups of Hemiptera are phytophagous. Vast literature exists for groups containing pest species (e.g., aphids, coccoids, psyllids), not restricted to management but also treating their complex life cycles. A review of this literature is beyond of the scope of the paper and thus the reader shall refer to particular references or to some of the general references indicated below.

Many species of Fulgoromorpha and Cicadomorpha are considered pests of cultivated crops. Dietrich (2005) and Wilson (2005) provide keys to economically important groups and provide useful literature.

Schaefer and Panizzi (2000) reviewed the economic importance, both of beneficial and pest species, of several groups of Heteroptera. Most of the economically important groups fall in the group of phytophagous species attacking cultivated crops, although one, Triatominae (Reduviidae), has species that are important vectors of Chagas' disease (Lent and Wygodzinsky 1979).

Catalogs and identification aids

Far from being complete, this section provides basic references for catalogs and or identification aids, in particular for the Neotropical Region. One general book of entomology, "Insects of Australia" (CSIRO 1991), has useful identification keys that cover South American taxa.

Identification aids

Hodgson (1994) provided a review of the genera of the Coccidae of the world with illustrations of key characters. Williams and Granara de Willink (1992) provided a synopsis for the Pseudococcidae of Central and South America.

Wilson (2005) provided keys to the economic important species of Fulgoromorpha of the Southeastern United States, with important references on the taxonomy and biology on the group as a whole. Deitz and Dietrich (1993) provided a key to the families of Membracoidea. Deitz (1975) provided keys to the subfamilies of Membracidae. Dietrich (2005) provided an illustrated key to the Cicadomorpha families, and to the subfamilies and tribes of Cicadellidae (except tribes of Deltocephalinae).

Schuh and Slater (1995) is an indispensable source of information for the classification, biology, and faunistics of Heteroptera in general, with keys to subfamilies or tribes of all groups. Comprehensive regional treatments for water bugs have been published for some tropical areas (e.g., Andersen and Weir 2004a). In the Neotropical region, nonetheless, scattered publications provide help to identify regional faunas (e.g., Nieser 1975; Pereira *et al.* 2007). In most cases original literature must be consulted for identification purposes.

Printed catalogs and lists

Ben-Dov (1993) provided a catalog of Coccidae, Ben-Dov (1994) of Pseudococcidae and Putonidae, Miller and Gimpel (2000) of Eriococcidae, Ben-Dov and German (2003) of Diaspididae, Ben-Dov (2005) of Margarodidae, Miller *et al.* (2005) and Ben-Dov (2006) for several other Coccoidea families. Kondo (2001) provided a list of the Coccoidea of Colombia.

Funkhouser (1927) catalog of the Membracidae, Aetalionidae, and Melizoderidae, was complemented with the additions of Metcalf and Wade (1963, 1965a, 1965b), and McKamey (1998). Metcalf's catalog for the Homoptera (Fulgoromorpha, Cicadoidea, and Membracoidea) is the starting point for much of the literature in Auchenorrhyncha (Metcalf 1932, 1936, 1943, 1945, 1946, 1947a, 1947b, 1954a-c, 1955a, 1955b, 1956, 1957, 1958, 1960a, 1960b, 1961, 1962a-f, 1963a-g, 1964a, 1964b, 1965a, 1965b, 1966a-d, 1967, 1968). Duffels and van der Laan (1985) and Oman *et al.* (1990) updated to the catalogs of Cicadoidea and Cicadellidae, respectively. A checklist with illustrations of the types of Cercopidae species of the New World was published by Carvalho and Webb (2005). Freytag and Sharkey (2002) provided a preliminary list of Cicadellidae from Colombia.

Stonedahl and Dolling (1991) provided a list of catalogs, monographs, and other literature relevant to identification purposes of different groups of Heteroptera. References not included in Stonedahl and Dolling (1991) include: Andersen (1995), for a checklist of the Gerrinae of the World; an update of the catalog of the Miridae (Schuh 1995); the update of the lygaeoid catalog (Slater and O'Donnell 1995); the Heteroptera catalog of Australia (Cassis and Gross 1995, 2002); the multivolume catalog of the Heteroptera of the Palearctic Region (Aukema and Rieger 1995, 1996, 1999, 2001, 2006); and the annotated checklist of the Heteroptera of Panama (Froeschner 1999).

Online catalogs and keys

Although printed catalogs are an indispensable source of information, the information contained is rapidly outdated, and they are more useful if available in digital format, for instance, in mapping taxa distribution in real-time from relational databases, or in showing the most current classification for a group. Furthermore, catalogs can link species and higher taxa to their respective digital publications if available (e.g., Schuh 2006). Many catalogs are already available, for instance, Coccoidea (Sternorrhyncha) (Miller and Ben-Dov 2006); Fulgoromorpha (Bourgoin 2007), Cercopoidea (Soulier-Perkins 2007), (Auchenorrhyncha), and Miridae (Heteroptera) (Schuh 2006). In addition, some regional checklists may provide further assistance in the identification of specimens, for instance, of Neotropical Cicadellidae (Freytag and Gaiani 2002), and the Cicadellinae of Colombia (Vargas-Rojas *et al.* 2006).

In addition, digital keys are now available via the Internet, which can display not only text information, but also color images. Another advantage of some of these keys is that they can be interactive and character based, unlike dichotomous, traditional keys. Identification keys are available for several groups of Hemiptera. For instance, families of Coccoidea (Miller *et al.* 2004); tribes of Cicadellidae (Dmitriev 2006), the genera of Erythroneurini (Dmitriev and Dietrich 2006) and Proconiini (Takiya and Dmitriev 2007); the Deltocephalinae related tribes (Zahniser 2007); families of Heteroptera of Australia (Cassis *et al.* 2002); and the genera of Pentatomidae (Cassis *et al.* 2003) and Tingidae of Australia (Cassis and Bulbert 2004).

Acknowledgments

I am thankful to Randall T. Schuh (American Museum of Natural History, USA) and Daniela M. Takiya (Universidade Fe-

deral do Paraná, Brazil) for the suggestions that greatly improved the manuscript.

Arthur Anker (STRI, Panama) kindly let me use many of his photographs (Figs. 1I-P, 2A-C, 3A-G) to illustrate this paper. Images of figures 1A-H from figure 1 were obtained from www.insectimages.org, under a Creative Commons License. Tomohide Yasunaga (Research Associate, AMNH; Nagasaki, Japan), let me use his Nepalese Orthotylinae photographs (Fig. 3H-J). The remaining photographs (Fig. 2D-K) were taken with a Microptics-USA photographic system equipped with an Infinity Photo-Optical K-2 three-lens system and a Nikon D1X digital camera, from specimens deposited at the American Museum of Natural History. Gervásio Silva Carvalho (Pontifícia Universidade Católica do Rio Grande do Sul) kindly identified the Cercopidae photos.

Cited Literature

- ALBERT, V. A. 2005. Parsimony, phylogeny, and genomics. Oxford University Press, New York, USA. 229 p.
- ALBERTSON, J. L.; DIETRICH, C. H. 2005. Systematics and phylogeny of the treehopper subfamily Nicomiinae (Hemiptera: Membracidae). *Revista Brasileira de Zoologia* 22 (1): 231-283.
- ANDERSEN, N. M. 1982. The semiaquatic bugs (Hemiptera, Gerromorpha). Phylogeny, adaptations, biogeography, and classification. *Entomograph* 3: 1-455.
- ANDERSEN, N. M. 1995. Cladistics, historical biogeography, and a check list of Gerrinae water striders (Hemiptera, Gerridae) of the World. *Steenstrupia* 21: 93-123.
- ANDERSEN, N. M. 1998. Water striders from the Paleogene of Denmark with a review of the fossil record and evolution of semiaquatic bugs (Hemiptera: Gerromorpha). *Det Kongelige Danske Videnskabernes Selskab, Biologiske Skrifter* 50: 1-152.
- ANDERSEN, N. M. 1999. *Cryptovelia stysi* sp. n. from Borneo with a reanalysis of the phylogeny of the Mesoveliidae (Heteroptera: Gerromorpha). *Acta Societatis Zoologicae Bohemicae* 63 :5-18.
- ANDERSEN, N. M. 2000. A new species of *Tetraripis* from Thailand, with a critical assessment of the generic classification of the subfamily Rhagoveliinae (Hemiptera, Veliidae). *Tijdschrift voor Entomologie* 142: 185-194.
- ANDERSEN, N. M.; WEIR, T. A. 1994. The sea skaters, genus *Halobates* Eschscholtz (Hemiptera: Gerridae), of Australia: taxonomy, phylogeny and zoogeography. *Invertebrate Taxonomy* 8: 861-909.
- ANDERSEN, N. M.; WEIR, T. A. 1999. The marine Haloveliinae (Hemiptera: Veliidae) of Australia, New Caledonia and southern New Guinea. *Invertebrate Taxonomy* 13: 309-350.
- ANDERSEN, N. M.; WEIR, T. A. 2004a. Australian water bugs (Hemiptera-Heteroptera, Gerromorpha and Nepomorpha). Their Biology and Identification. *Entomograph* 14: 1-344.
- ANDERSEN, N. M.; WEIR, T. A. 2004b. Mesoveliidae, Hebridae, and Hydrometridae of Australia (Hemiptera: Heteroptera: Gerromorpha), with a reanalysis of the phylogeny of semiaquatic bugs. *Invertebrate Systematics* 18: 467-522.
- ARNETT, R. H. 2000. American insects: a handbook of the insects of America north of Mexico. CRC Press, Boca Raton, USA. 1.003 p.
- ASCHE, M. 1987. Preliminary thoughts on the phylogeny of Fulgoromorpha (Homoptera Auchenorrhyncha). p. 47-53. In: *Proceedings of the 6th Auchenorrhyncha Meeting, Turin, Italy.*
- AUKEMA, B.; RIEGER, C. 1995. Catalogue of the Heteroptera of the Palearctic Region. Vol. 1. Enicocephalomorpha, Dispsocomorpha, Nepomorpha, Gerromorpha and Leptopodomorpha. The Netherlands Entomological Society, Amsterdam, the Netherlands. 222 p.

- AUKEMA, B.; RIEGER, C. 1996. Catalogue of the Heteroptera of the Palearctic Region. Vol. 2. Cimicomorpha I. The Netherlands Entomological Society, Amsterdam, the Netherlands. 361 p.
- AUKEMA, B.; RIEGER, C. 1999. Catalogue of the Heteroptera of the Palearctic Region. Vol. 3. Cimicomorpha II. The Netherlands Entomological Society, Amsterdam, the Netherlands. 576 p.
- AUKEMA, B.; RIEGER, C. 2001. Catalogue of the Heteroptera of the Palearctic Region. Vol. 4. Pentatomomorpha I. The Netherlands Entomological Society, Amsterdam, the Netherlands. 346 p.
- AUKEMA, B.; RIEGER, C. 2006. Catalogue of the Heteroptera of the Palearctic Region. Vol. 5. Pentatomomorpha II. The Netherlands Entomological Society, Amsterdam, the Netherlands. 550 p.
- BEN-DOV, Y. 1993. A systematic catalogue of the soft scale insects of the World (Homoptera: Coccoidea: Coccidae) with data on geographical distribution, host plants, biology, and economic importance. Flora & Fauna Handbook, No. 9. Sandhill Crane Press, Gainesville, USA. 536 p.
- BEN-DOV, Y. 1994. A systematic catalogue of the mealybugs of the world (Insecta: Homoptera: Coccoidea: Pseudococcidae and Putonidae) with data on geographical distribution, host plants, biology and economic importance. Intercept Limited, Andover, UK. 868 p.
- BEN-DOV, Y. 2005. A systematic catalogue of the scale insect family Margarodidae (Hemiptera: Coccoidea) of the World. Intercept Limited, Wimborne, UK. 400 p.
- BEN-DOV, Y. 2006. A systematic catalogue of eight scale insect families (Hemiptera: Coccoidea) of the world Aclerodidae, Asterolecaniidae, Beesoniidae, Carayonemidae, Conchaspidae, Dactylopiidae, Kerriidae and Lecanodiaspididae. Elsevier, Amsterdam, the Netherlands. 368 p.
- BEN-DOV, Y.; GERMAN, V. 2003. A systematic catalogue of the Diaspididae (armoured scale insects) of the World, subfamilies Aspidiotinae, Comstockiellinae, and Odonaspidinae. Intercept Limited, Andover, UK. 1111 p.
- BEUTEL, R. G.; GORB, S. N. 2001. Ultrastructure of attachment specializations of hexapods (Arthropoda): evolutionary patterns inferred from a revised ordinal phylogeny. Journal of Zoological Systematics and Evolutionary Research 39 (4): 177-207.
- BEUTEL, R. G.; WEIDE, D. 2005. Cephalic anatomy of *Zorotypus hubbardi* (Hexapoda: Zoraptera): new evidence for a relationship with Acercaria. Zoomorphology 124: 121-136.
- BLACKMAN, R. L.; EASTOP, V. F. 2006. Aphids on the World's Herbaceous Plants and Shrubs. The Natural History Museum, two volumes, John Wiley & Sons, Chichester, UK. 1439 p.
- BÖRNER, C. 1904. Zur Systematik der Hexapoden. Zoologischer Anzeiger 27: 511-533.
- BORROR, D. J.; DE LONG, D. M.; TRIPLEHORN, C. A. 1981. An introduction to the study of insects. Fifth Edition, Saunders College Publishing, USA. 928 p.
- BORROR, D. J.; WHITE, R. E. 1970. A field guide to insects America north of Mexico. Peterson Field Guide series, Houghton Mifflin Company, Boston and New York, USA. 404 p.
- BOUDREAUX, H. B. 1979. Arthropod phylogeny with special reference to insects. Wiley, New York, USA. 320 p.
- BOURGOIN, T. 1986a. Morphologie imaginaire du tentorium des Hemiptera Fulgoromorpha. International Journal of Insect Morphology & Embryology 15 (4): 237-252.
- BOURGOIN, T. 1986b. Valeur morphologique de la lame maxillaire chez les Hemiptera; remarques phylogénétiques. Annales de la Société Entomologique de France (Nouvelle série) 22 (4): 413-422.
- BOURGOIN, T. 1993. Female genitalia in Hemiptera Fulgoromorpha, morphological and phylogenetic data. Annales de la Société Entomologique de France (Nouvelle série) 29 (3): 225-244.
- BOURGOIN, T. 2007. FLOW. Fulgoromorpha lists on the web. Available at: <<http://flow.snv.jussieu.fr/cgi-bin/entomosite.pl?base=FLOW&page=home&lang=en>>. [Accessed: February 2008].
- BOURGOIN, T.; CAMPBELL, B. C. 2002. Inferring a phylogeny for Hemiptera: falling into the 'autapomorphic trap'. Denisia 4: 67-82.
- BOURGOIN, T.; HUANG, J. 1990. Morphologie comparée des genitalia mâles des Trypetermorphi et remarques phylogénétiques (Hemiptera: Fulgoromorpha: Tropiduchidae). Annales de la Société Entomologique de France (N.S.) 26 (4): 555-564.
- BOURGOIN, T.; STEFFEN-CAMPBELL, J. D.; CAMPBELL, B. C. 1997. Molecular phylogeny of Fulgoromorpha (Insecta, Hemiptera, Archaeorrhyncha). The Enigmatic Tettigometridae: Evolutionary Affiliations and Historical Biogeography. Cladistics 13: 207-224.
- BRUES, C. T.; MELANDER, A. L.; CARPENTER, F. M. 1954. Classification of insects. Keys to the living and extinct families of insects, and to the living families of other terrestrial arthropods. Bulletin of the Museum of Comparative Zoology 108: 1-917.
- BURCKHARDT, D. 2005. Biology, ecology, and evolution of gall-inducing psyllids (Hemiptera: Psylloidea), pp. 143-157. In: Raman, A.; Schaefer C.W.; Withers T.M. (eds.). Biology, ecology, and evolution of gall-inducing arthropods. Science Publishers, Enfield, USA. 2 vols., 817 p.
- BURCKHARDT, D.; AGOSTI, D. 1991. New records of South American Peloriidiidae (Homoptera: Coleorrhyncha). Revista Chilena de Entomología 19: 71-75.
- BURCKHARDT, D.; CEKALOVIC K., T. 2002. An anomalous moss-bug from southern Chile and notes on *Pantinia darwini* (Hemiptera, Coleorrhyncha, Peloriidiidae). Mitteilungen der Schweizerischen Entomologischen Gesellschaft 75 (1-2): 57-59.
- BURCKHARDT, D.; LAUTERER, P. 1989. Systematics and biology of Rhinocolinae (Homoptera: Psylloidea). Journal of Natural History 23: 643-712.
- BYRNE, D. N.; BELLOWS, T. S. Jr. 1991. Whitefly biology. Annual Review of Entomology 36: 431-457.
- CAMERON, S. L.; BECKENBACH, A. T.; DOWTON, M.; WHITING, M. F. 2006. Evidence from mitochondrial genomics on interordinal relationships in insects. Arthropod Systematics and Phylogeny 64 (1): 27-34.
- CAMPBELL, B. C.; STEFFEN-CAMPBELL, J. D.; GILL, R. J. 1994. Evolutionary origin of whiteflies (Hemiptera: Sternorrhyncha: Aleyrodidae) inferred from 18S rDNA sequences. Insect Molecular Biology 3 (2): 73-88.
- CAMPBELL, B. C., STEFFEN-CAMPBELL, J. D.; SORENSON, J. T.; GILL, R. J. 1995. Paraphyly of Homoptera and Auchenorrhyncha inferred from 18S rDNA nucleotide sequences. Systematic Entomology 20 (3): 175-194.
- CARAYON, J. 1962. Observations sur l'appareil odorifique des Hétéroptères particulièrement celui des Tingidae, Vianaididae et Piesmatidae. Cahiers des Naturalistes 18 (1): 1-16.
- CARAYON, J. 1971. Notes et documents sur l'appareil odorant métathoracique des Hémiptères. Annales de la Société Entomologique de France 7 (4): 737-770.
- CARVALHO, G. S.; WEBB, M. D. 2005. Cercopid spittle bugs of the New World (Hemiptera, Auchenorrhyncha, Cercopidae). Pensoft, Sofia, Bulgaria. 271 p.
- CARVER, M.; GROSS, G. F.; WOODWARD, T. E. 1991. Chapter 30. Hemiptera, pp. 429-509. In: CSIRO, Division of Entomology. Insects of Australia. 2nd edition, 2 volumes. Cornell University Press, Ithaca, USA. 1137 p.
- CASSIS, G.; BULBERT, M. 2004. Key to the Tingidae of Australia. FaunaNet. Accessed: February 2008. Available at: <http://www.faanet.gov.au/faunakeys/tingidae_intro.htm>.
- CASSIS, G.; GROSS, G. F. 1995. Hemiptera: Heteroptera (Coleorrhyncha to Cimicomorpha). Houston, W. W. K.; Maynard, G. V. (eds.). Zoological Catalogue of Australia. Vol. 27.3A. CSIRO, Melbourne, Australia. 506 p.
- CASSIS, G.; GROSS, G. F. 2002. Hemiptera: Heteroptera (Pentatomomorpha). Houston, W. W. K.; Wells, A. (eds.). Zoological

- Catalogue of Australia. Vol. 27. B. CSIRO, Melbourne, Australia. 737 p.
- CASSIS, G.; BETTS, E.; ELLIOTT, M. G. 2003. Key to the stink bugs of Australia [Heteroptera: Pentatomidae]. FaunaNet. Available at: <http://www.faunanet.gov.au/faunakeys/stink_intro.htm>. [Accessed: February 2008].
- CASSIS, G.; FLEMONS, P.; ELLIOTT, M. G.; DONNELLY, A.; LANE, J.; BULBERT, M.; SMITH, M.; REID, C. A. M.; HARRIS, R.; CARTER, G.; SILVEIRA, R. 2002. Key to the Heteroptera of Australia. FaunaNet. Available at: <http://www.faunanet.gov.au/faunakeys/heteroptera_intro.htm>. [Accessed: February 2008].
- CHEN, S.; YANG, C. T. 1995. The metatarsi of the Fulgoroidea (Homoptera: Auchenorrhyncha). Chinese Journal of Entomology 15: 257-269.
- CHINA, W. E. 1962. South American Peloriidiidae (Hemiptera-Homoptera: Coleorrhyncha). Transactions of the Royal Entomological Society of London 114 (5): 131-161.
- CHINA, W. E.; MILLER, N. C. E. 1959. Check-list and keys to the families and subfamilies of the Hemiptera-Heteroptera. Bulletin of the British Museum of Natural History 8 (1): 1-45.
- CHOU, I.; LEI, Z.; LI, L.; LU, X.; YAO, W. 1997. The Cicadidae of China (Homoptera: Cicadoidea). Tianze Eldoneio, Hong Kong. 380 p. [In Chinese].
- CLARIDGE, M. F. 1985. Acoustic signals in the Homoptera: behavior, taxonomy, and evolution. Annual Review of Entomology 30: 297-317.
- COBBEN, R. H. 1978. Evolutionary trends in Heteroptera. Part 2. Mouthpart-structures and feeding strategies. Mededelingen Landbouwhogeschool 78-5. H. Veeman, Wageningen, the Netherlands. 407 p.
- COOK, L. G.; GULLAN, P. J.; TRUEMAN, H. E. 2002. A preliminary phylogeny of the scale insects (Hemiptera: Sternorrhyncha: Coccoidea) based on nuclear small-subunit ribosomal DNA. Molecular Phylogenetics and Evolution 25: 43-52.
- COX, J. M.; WILLIAMS, D. J. 1987. Do the Eriococcidae form a monophyletic group? Bollettino del Laboratorio di Entomologia Agraria "Filippo Silvestri" Portici 43 (Suppl.): 13-17.
- CRANSTON, P. S.; GULLAN, P. J. 2003. Phylogeny of insects., pp. 882-898. In: Resh, V. H.; Cardé, R. T. (eds.). Encyclopedia of Insects. Academic Press, San Diego, USA. xxviii + 1266 p.
- CRESPI, B.; CARMEAN, D.; VAWTER, L.; VON DOHLEN, C. 1996. Molecular phylogenetics of Thysanoptera. Systematic Entomology 21: 79-87.
- CROCROFT, R. B.; ANDRODRÍGUEZ, R. L. 2005. The behavioral ecology of insect vibrational communication. Bioscience 55 (4): 323-334.
- CRYAN, J. A. 2005. Molecular phylogeny of Cicadomorpha (Insecta: Hemiptera: Cicadoidea, Cercopoidea and Membracoidea): adding evidence to the controversy. Systematic Entomology 30 (4): 563-574.
- CRYAN, J. R.; ROBERTSON, J. A.; DEITZ, L. L. 2003. The New World treehopper tribe Microcentrini (Hemiptera: Membracidae: Stegaspidinae): Monographic Revision and Phylogenetic Position. Thomas Say Publications, Monographs of the Entomological Society of America, Lanham, USA. 108 p.
- CRYAN, J. R.; WIEGMANN, B. M.; DEITZ, L. L.; DIETRICH, C. H. 2000. Phylogeny of the treehoppers (Insecta: Hemiptera: Membracidae): evidence from two nuclear genes. Molecular Phylogenetics and Evolution 17: 317-334.
- CRYAN, J. R.; WIEGMANN, B. M.; DEITZ, L. L.; DIETRICH, C. H.; WHITING, M. F. 2004. Treehopper trees: phylogeny of Membracidae (Hemiptera: Cicadomorpha: Membracoidea) based on molecules and morphology. Systematic Entomology 29: 441-454.
- CSIRO [Commonwealth Scientific and Industrial Research Organisation, Division of Entomology]. 1991. Insects of Australia. 2nd edition, 2 volumes. Cornell University Press, Ithaca, USA. 1137 p.
- DAMGAARD, J.; COGNATO, A. I. 2005. Phylogeny and reclassification of species groups in *Aquarius* Schellenberg, *Limnopus* Stål and *Gerris* Fabricius (Insecta: Hemiptera-Heteroptera, Gerridae). Systematic Entomology 31: 93-112.
- DAMGAARD, J.; SPERLING, F. A. H. 2001. Phylogeny of the water strider genus *Gerris* Fabricius (Heteroptera: Gerridae) based on COI mtDNA, EF-1a nuclear DNA and morphology. Systematic Entomology 26: 241-254.
- DAMGAARD, J.; ANDERSEN, N. M.; CHENG, L.; SPERLING, F. A. H. 2000a. Phylogeny of sea skaters, *Halobates* (Hemiptera, Gerridae), based on mtDNA sequence and morphology. Zoological Journal of the Linnean Society 130: 511-526.
- DAMGAARD, J.; ANDERSEN, N. M.; SPERLING, F. A. H. 2000b. Phylogeny of the water strider genus *Aquarius* Schellenberg (Heteroptera: Gerridae) based on mitochondrial and nuclear DNA and morphology. Insect Systematics and Evolution 31: 71-90.
- DAMGAARD, J.; ANDERSEN, N. M.; MEIER, R. 2005. Combining molecular and morphological analyses of water strider phylogeny (Hemiptera-Heteroptera, Gerromorpha): effects of alignment and taxon sampling. Systematic Entomology 30 (2): 289-309.
- DEITZ, L. L. 1975. Classification of the higher categories of the New World treehoppers (Homoptera: Membracidae). North Carolina Agricultural Experiment Station Technical Bulletin 225: i-iv, 1-177.
- DEITZ, L. L.; DIETRICH, C. H. 1993. Superfamily Membracoidea (Homoptera: Auchenorrhyncha). I. Introduction and revised classification with new family-group taxa. Systematic Entomology 18: 287-296.
- DIETRICH, C. H. 1999. The role of grasslands in the diversification of leafhoppers (Homoptera: Cicadellidae): a phylogenetic perspective, pp. 44-48. In: Warwick, C. (ed.). Proceedings of the Fifteenth North American Prairie Conference. Natural Areas Association, Bend, Oregon, USA.
- DIETRICH, C. H. 2004. Phylogeny of the leafhopper subfamily Evacanthinae with a review of Neotropical species and notes on related groups (Hemiptera: Membracoidea: Cicadellidae). Systematic Entomology 29: 455-487.
- DIETRICH, C. H. 2005. Keys to the families of Cicadomorpha and subfamilies and tribes of Cicadellidae (Hemiptera: Auchenorrhyncha). Florida Entomologist 88 (4): 502-517.
- DIETRICH, C. H.; DEITZ, L. L. 1991. Numerical phenetic and cladistic analyses of the treehopper tribe Aconophorini (Homoptera: Membracidae: Membracinae). Annals of Entomological Society America 84 (3): 228-238.
- DIETRICH, C. H.; DEITZ, L. L. 1993. Superfamily Membracoidea (Homoptera: Auchenorrhyncha). II. Cladistic analysis and conclusions. Systematic Entomology 18: 297-312.
- DIETRICH, C. H.; MCKAMEY, S. H.; DEITZ, L. L. 2001a. Morphology based phylogeny of the treehopper family Membracidae (Hemiptera: Cicadomorpha: Membracoidea). Systematic Entomology 26: 213-239.
- DIETRICH, C. H.; RAKITOV, R. A.; HOLMES, J. L.; BLACK, W. C. IV. 2001b. Phylogeny of the Major Lineages of Membracoidea (Insecta: Hemiptera: Cicadomorpha) Based on 28S rDNA Sequences. Molecular Phylogenetics and Evolution 18 (2): 293-305.
- DIETRICH, C. H.; WALLNER, A. M. 2002. Diversity and taxonomic composition of Cicadellidae in the Amazonian rainforest canopy (Hemiptera, Cicadomorpha, Membracoidea). Abstracts of the 11th International Auchenorrhyncha Congress, Potsdam/Berlin.
- DMITRIEV, D. A. 2006. An interactive key to tribes of Leafhoppers (Cicadellidae). Available at: <<http://ctap.inhs.uiuc.edu/dmitriev/key.asp?key=Cicnymph&i=1&lng=En>>. [Accessed: February 2008].
- DMITRIEV, D. A.; DIETRICH, C. H. 2006. An interactive key to genera of Erythroneurini. Available at: <<http://ctap.inhs.uiuc.edu/dmitriev/key.asp?key=Erythroneura&lng=En&i=1&keyN=1>>. [Accessed: February 2008].
- DOWNIE, D. A.; GULLAN, P. J. 2004. Phylogenetic analysis of mealybugs (Hemiptera: Coccoidea: Pseudococcidae) based on

- DNA sequences from three nuclear genes, and a review of the higher classification. *Systematic Entomology* 29: 238-259.
- DUFFELS, J. P.; VAN DEER LAAN, P. A. 1985. Catalogue of the Cicadoidea (Homoptera, Auchenorrhyncha) 1956 - 1980. *Series Entomologica*, vol. 34. Dr W. Junk Publishers, Kluwer Academic Publishers Group, Dordrecht, the Netherlands. 414 pp.
- DUFOUR, L. 1833. Recherches anatomiques et physiologiques sur les hemiptères: accompagnées de considérations relatives à l'histoire naturelle et à la classification de ces insectes. Impr. de Bachelier, Paris, France. xix + 333 p.
- EMELJANOV, A. F. 1990. An attempt of construction of the phylogenetic tree of the planthoppers (Homoptera, Cicadina). *Entomologicheskoe Obozrenie* 69 (2): 353-356.
- ESTÉVEZ, A. L.; DE REMES-LENICOV, A. M. M. 1989. Pelorididos de Tierra del Fuego (Homoptera: Coleorrhyncha). *Neotropica (La Plata)* 34 (92): 104.
- EVANS, J. W. 1963. The phylogeny of the Homoptera. *Annual Review of Entomology* 8: 77-94.
- EVANS, J. W. 1981. A review of the present knowledge of the family Peloridiidae and new genera and species from New Zealand and New Caledonia (Hemiptera: Insecta). *Records of the Australian Museum* 34 (5): 381-406.
- FAGUA, G. 2005. Sobre el uso poco acertado del taxón "Homoptera". *Entomólogo* 33 (99): 3-5.
- FISCHER, C.; MAHNER, M.; WACHMANN, E. 2000. The rhabdom structure in the ommatidia of the Heteroptera (Insecta), and its phylogenetic significance. *Zoomorphology* 120 (1): 1-13.
- FOLDI, I., 1997. Defense strategies in scale insects: phylogenetic inference and evolutionary scenarios (Hemiptera, Coccoidea), pp. 203-230. In: Grandcolas, P. (ed.). *The Origin of Biodiversity in Insects: Phylogenetic Tests of Evolutionary Scenarios. Mémoires du Muséum national d'Histoire naturelle* 173: 1-354.
- FREYTAG, P. H.; GAIANI, M. 2002. Saltahojas Neotropicales neotropicales / Neotropical Leafhoppers leafhoppers (Hemiptera: Cicadellidae). Available at: <<http://www.miza-fpolar.info/ve/cicadellidae/>>. [Accessed: February 2008].
- FREYTAG, P. H.; SHARKEY, M. J. 2002. A preliminary list of the leafhoppers (Homoptera: Cicadellidae) of Colombia. *Biota Colombiana* 3 (2): 235-283.
- FROESCHNER, R. C. 1999. True bugs (Heteroptera) of Panama: A synoptic catalog as a contribution to the study of Panamanian biodiversity. *Memoirs of the American Entomological Institute* 61: 1-393.
- FUNKHOUSER, W. D. 1927. General catalogue of the Hemiptera. Fascicle I. Membracidae. Smith College, Northampton, USA. 581 p.
- GOGALA, M. 2006. Vibratory signals produced by Heteroptera-Pentatomomorpha and Cimicomorpha, pp. 275-295. In: Drosopoulos, S.; Claridge, M. F. (eds.). *Insect sounds and communication. Physiology, behaviour, ecology and evolution*. CRC (Taylor & Francis), Boca Raton, USA. xvii + 532 p. + a DVD.
- GOODCHILD, A. J. P. 1966. Evolution of the alimentary canal of the Hemiptera. *Biological Review* 41: 97-140.
- GORB, R. G.; BEUTEL, R. G. 2001. Evolution of locomotory attachment pads of hexapods. *Naturwissenschaften* 88 (12): 530-534.
- GRAZIA, J.; SCHUH, R. T.; WHEELER, W. C. *in press*. Phylogenetic relationships of family groups in Pentatomoidea based on morphology and DNA sequences (Insecta: Heteroptera). *Cladistics*.
- GRIMALDI, D.; ENGEL, M. S. 2005. *Evolution of the insects*. Cambridge University Press, New York, USA. 755 pp.
- GULLAN, P. 1999 [2001]. Why the taxon Homoptera does not exist. *Entomologica* 33: 101-104.
- GULLAN, P.; COOK, L. G. 2007. Phylogeny and higher classification of the scale insects (Hemiptera: Sternorrhyncha: Coccoidea). *Zootaxa* 1668: 413-425.
- GULLAN, P. J.; KOSZTARAB, M. 1997. Adaptations in scale insects. *Annual Review of Entomology* 42: 23-50.
- HACKER, H. 1932. A new species of Peloridiidae from Queensland. *Queensland agricultural Journal* 38: 262-263.
- HAMILTON, K. G. A. 1981. Morphology and evolution of the Rhynchotan head (Insecta: Hemiptera, Homoptera). *The Canadian Entomologist* 113 (11): 953-974.
- HAMILTON, K. G. A. 1983. Classification, morphology and phylogeny of the family Cicadellidae (Rhynchota: Homoptera), pp. 15-37. In: Knight, W. J.; Pant, N. C.; Robertson, T. S.; Wilson, M. R. (eds.). *Proceedings of the 1st international workshop on leafhoppers and planthoppers of economic importance*, London, 4-7 Oct. 1982. Commonwealth Institute of Entomology, London, UK.
- HAMILTON, K. G. A. 1999. The ground-dwelling leafhoppers Myerslopiidae, new family, and Sagmatini, new tribe (Homoptera: Membracoidea). *Invertebrate Taxonomy* 13: 207-235.
- HAMILTON, K. G. A. 2001. A new family of froghoppers from the American tropics (Hemiptera: Cercopoidea: Epipygidae). *Biodiversity* 2: 15-21.
- HARDY, N. B.; GULLAN, P. J.; HODGSON, C. J. 2008. A subfamily-level classification of mealybugs (Hemiptera: Pseudococcidae) based on integrated molecular and morphological data. *Systematic Entomology* 33: 51-71.
- HEBSGAARD, M. B.; ANDERSEN, N. M.; DAMGAARD, J. 2004. Phylogeny of the true water bugs (Nepomorpha: Hemiptera-Heteroptera) based on 16S and 28S rDNA and morphology. *Systematic Entomology* 29: 488-508.
- HEIE, O. E. 1987. Palaeontology and phylogeny, pp. 367-391. In: Minks, A. K.; Harrewijn, P. (eds.). *Aphids: Their biology, natural enemies, and control*. Vol 2A. Elsevier, Amsterdam, the Netherlands.
- HEMING, B. S. 1980. Development of the mouthparts in embryos of *Haplothrips verbasci* (Osborn) (Insecta, Thysanoptera, Phlaeothripidae). *Journal of Morphology* 164: 235-263.
- HENNIG, W. 1966. *Phylogenetic systematics*. University of Illinois Press, Urbana, USA. 263 p.
- HENNIG, W. 1969. *Die Stammesgeschichte der Insekten*. Waldemar Kramer, Frankfurt am Main, Germany. 436 p.
- HENNIG, W. 1981. *Insect phylogeny*. John Wiley & Sons, New York, USA. 514 p.
- HENRY, T. J. 1997. Phylogenetic analysis of family groups within the infraorder Pentatomorpha (Hemiptera: Heteroptera), with emphasis on the Lygaeoidea. *Annals of the Entomological Society of America* 90 (3): 275-301.
- HILL, L. 1987. First record of Dipsocoridae (Hemiptera) from Australia with the description of four new species of *Cryptostemma* Herrich-Schaeffer. *Journal of the Australian Entomological Society* 26: 129-139.
- HOCH, H.; DECKERT, J.; WESSEL, A. 2006. Vibrational signalling in a Gondwanan relict insect (Hemiptera: Coleorrhyncha: Peloridiidae). *Biology Letters* 2: 222-224.
- HODGSON, C. J. 1994. *The Scale insect family Coccidae: An identification manual to genera*. CAB International, Wallingford, UK. 639 p.
- HODKINSON, I. D. 1989. The biogeography of the Neotropical jumping plant-lice (Insecta: Homoptera: Psylloidea). *Journal of Biogeography* 16 (3): 203-217.
- HODKINSON, J. D.; CASSON, D. 1991. A lesser predilection for bugs: Hemiptera (Insecta) diversity in tropical rain forests. *Biological Journal of the Linnean Society* 43: 101-109.
- HOLLIS, D. 2004. Australian Psylloidea: jumping plantlice and lerp insects. Australian Biological Resources Study (ABRS), Canberra, Australia. 232 p.
- HOY, R. R.; ROBERT, D. 1996. Tympanal hearing in insects. *Annual Review of Entomology* 41: 433-450.
- HÜNEFELD, F. 2007. The genital morphology of *Zorotypus hubbardi* Caudell, 1918 (Insecta, Zoraptera, Zorotypidae). *Zoomorphology* 126: 135-151.

- ICZN (International Commission on Zoological Nomenclature). 1999. International Code of Zoological Nomenclature. 4th edition. International Trust for Zoological Nomenclature, London, UK. xxix + 306 pp.
- JAMIESON, B. G. M.; DALLAI, R.; AFZELIUS, B. A. 1999. Insects. Their spermatozoa and phylogeny. Science Publishers Inc., Enfield, USA. 555 p.
- JOHNSON, K. P.; YOSHIZAWA, K.; SMITH, V. S. 2004. Multiple origins of parasitism in lice. Proceedings of the Royal Society of London, series B 271: 1771-1776.
- KERZHNER, I. M. 1981. Fauna SSSR. Nasekomye Khobotnye, t. 13, vyp. 2, Poluzhestkokrylye Semejstva Nabidae. [Insecta Rhynchota. Vol. 13, No.2. Hemiptera family Nabidae. Fauna of the USSR.] Nauka Publishing, Leningrad, Russia. 326 p. [In Russian].
- KITCHING, I. J.; FOREY, P. L.; HUMPHRIES, C. J.; WILLIAMS, D. M. 1998. Cladistics. The Theory and practice of parsimony analysis. 2nd edition. Oxford University Press, New York, USA. 228 p.
- KJER, K. M. 2004. Aligned 18S and Insect Phylogeny. Systematic Biology 53 (3): 506-514.
- KJER, K. M.; CARLE, F. M.; LITMAN, J.; WARE, J. 2006. A molecular phylogeny of Hexapoda. Arthropod Systematics and Phylogeny 64 (1): 35-44.
- KONDO, T. 2001. Las cochinitas de Colombia (Hemiptera: Coccoidea). Biota Colombiana 2 (1): 31-48.
- KOTEJA, J. 1974. On the phylogeny and classification of the scale insects (Homoptera, Coccinea). Acta Zoologica Cracoviensia, 19: 267-326.
- KRISTENSEN, N. P. 1975. The phylogeny of hexapod orders. A critical review of recent accounts. Zeitschrift für Zoologische Systematik und Evolutionsforschung 13: 1-44.
- KRISTENSEN, N. P. 1981. Phylogeny of insect orders. Annual Review of Entomology 26: 135-157.
- KRISTENSEN, N. P. 1991. Chapter 5. Phylogeny of extant Hexapods. Pp. 125-140. In: CSIRO, Division of Entomology. Insects of Australia. 2nd edition, 2 volumes. Cornell University Press, Ithaca, USA. 1137 p.
- KUKALOVÁ-PECK, J. 1991. Chapter 6. Fossil history and the evolution of Hexapod structures. pp: 141-179. In: CSIRO, Division of Entomology. Insects of Australia. 2nd edition, 2 volumes. Cornell University Press, Ithaca, USA. 1137 p.
- LATREILLE, P. A. 1810. Considérations générales sur l'ordre naturel des animaux composant les classes des crustacés, des arachnides, et des insectes; avec un tableau méthodique de leurs genres, disposés en familles. F. Schoell, Paris, France. 444 p.
- LENT, H.; WYGODZINSKY, P. 1979. Revision of the Triatominae (Hemiptera: Reduviidae) and their significance as vectors of Chagas' disease. Bulletin of the American Museum of Natural History 163: 123-520.
- LESTON, D.; PENDERGRAST, J. G.; SOUTHWOOD, T. R. E. 1954. Classification of terrestrial (Geocorisae). Nature 174 (4419): 91-92.
- LI, H.-M.; DENG, R.-Q.; WANG, J.-W.; CHEN, Z.-Y.; JIA, F.-L.; WANG, X.-Z. 2005. A preliminary phylogeny of the Pentatomomorpha (Hemiptera: Heteroptera) based on nuclear 18S rDNA and mitochondrial DNA sequences. Molecular Phylogenetics and Evolution 37: 313-326.
- LIANG, A. P.; FLETCHER, M. J. 2002. Morphology of the antennal sensilla in four Australian spittlebug species (Hemiptera: Cercopidae) with implications for phylogeny. Australian Journal of Entomology 41: 39-44.
- LIN, C.-P.; DANFORTH, B. N.; WOOD, T. K. 2004. Molecular phylogenetics and evolution of maternal care in Membracine treehoppers. Systematic Biology 53 (3): 400-421.
- LYAL, C. H. C. 1985. Phylogeny and classification of the Psocodea, with particular reference to the lice (Psocodea: Phthiraptera). Systematic Entomology 10: 145-165.
- MAHNER, M. 1993. Systema Cryptoceratorum Phylogenicum (Insecta, Heteroptera). Zoologica 48 (143): 1-302.
- MALDONADO C., J. 1990. Systematic catalogue of the Reduviidae of the World. Caribbean Journal of Science, Special publication No. 1, University of Puerto Rico, Mayagüez, Puerto Rico. 694 p.
- MANZARI, S.; QUICKE, D. L. J. 2006. A cladistic analysis of whiteflies, subfamily Aleyrodinae (Hemiptera: Sternorrhyncha: Aleyrodidae). Journal of Natural History 40 (44-46): 2423-2554.
- MARTIN, J. H.; MOUND, L. A. 2007. An annotated check list of the world's whiteflies (Insecta: Hemiptera: Aleyrodidae). Zootaxa 1492: 1-84.
- MARTÍNEZ-TORRES, D.; BUADES, C.; LATORRE, A.; MOYA, A. 2001. Molecular systematics of aphids and their primary endosymbionts. Molecular Phylogenetics and Evolution 20: 437-449.
- McATEE, W. L.; MALLOCH, J. R. 1925. Revision of the bugs of the family Cryptostemmatidae in the collection of the United States National Museum. Proceedings of the U.S. National Museum 67 (13): 1-42, pls. 1-4.
- McKAMEY, S.H. 1998. Taxonomic catalogue of the Membracoidea (exclusive of leafhoppers): second supplement to fascicle I - Membracidae of the general catalogue of the Hemiptera. Memoirs of the American Entomological Institute 60: 1-377.
- McKAMEY, S. H. 2001. Checklist of leafhopper species 1758-1955 (Hemiptera: Membracoidea: Cicadellidae and Myserslopiidae) with synonymy and distribution [Catalogue of the Homoptera, Fascicle 6, Abridged]. Available at: <<http://www.sel.barc.usda.gov/selhome/leafhoppers/mckpaper.htm>>. [Accessed: October 2007].
- METCALF, Z. P. 1932. General catalogue of the Homoptera. Fascicle IV Fulgoroidea. Part 1 Tettigometridae. Smith College, Northampton, USA. 74 p.
- METCALF, Z. P. 1936. General catalogue of the Homoptera. Fascicle IV Fulgoroidea. Part 2 Cixiidae. Smith College, Northampton, USA. 269 p.
- METCALF, Z. P. 1943. General catalogue of the Homoptera. Fascicle IV Fulgoroidea. Part 3 Araeopidae (Delphacidae). Smith College, Northampton, USA. 552 p.
- METCALF, Z. P. 1945. General catalogue of the Homoptera. Fascicle IV Fulgoroidea. Part 4 Derbidae, Part 5 Achilixiidae, Part 6 Meenoplidae, Part 7 Kinnaridae. Smith College, Northampton, USA. 252 p.
- METCALF, Z. P. 1946. General catalogue of the Homoptera. Fascicle IV Fulgoroidea. Part 8. Dictyopharidae. Smith College, Northampton, USA. 246 p.
- METCALF, Z. P. 1947a. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 9 Fulgoridae. Smith College, Northampton, USA. 276 p.
- METCALF, Z. P. 1947b. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 10 Achilidae. Smith College, Northampton, USA. 276 p.
- METCALF, Z. P. 1954a. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 11 Tropiduchidae. North Carolina State College, Raleigh, USA. 167 p.
- METCALF, Z. P. 1954b. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 12 Nogodinidae. North Carolina State College, Raleigh, USA. 75 p.
- METCALF, Z. P. 1954c. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 14 Acanaloniidae. North Carolina State College, Raleigh, USA. 55 p.
- METCALF, Z. P. 1955a. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 16 Ricanidae. North Carolina State College, Raleigh, USA. 199 p.
- METCALF, Z. P. 1955b. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 17 Lophopidae. North Carolina State College, Raleigh, USA. 75 p.
- METCALF, Z. P. 1956. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 18 Eurybrachidae and Gengidae. North Carolina State College, Raleigh, USA. 81 p.

- METCALF, Z. P. 1957. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 13 Flatidae and Hypochthonellidae. North Carolina State College, Raleigh, USA. 565 p.
- METCALF, Z. P. 1958. General catalog of the Homoptera. Fascicle IV Fulgoroidea. Part 15 Issidae. North Carolina State College, Raleigh, USA. 561 p.
- METCALF, Z. P. 1960a. A bibliography of the Cercopoidea (Homoptera: Auchenorrhyncha). North Carolina State College, Raleigh, USA. 262 p.
- METCALF, Z. P. 1960b. General catalog of the Homoptera. Fascicle VII Cercopoidea. Part 1. Machaerotidae. North Carolina State College, Raleigh, USA. 49 p.
- METCALF, Z. P. 1961. General catalog of the Homoptera. Fascicle VII Cercopoidea. Part 2. Cercopidae. North Carolina State College, Raleigh, USA. 607 p.
- METCALF, Z. P. 1962a. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 2. Hylicidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 18 p.
- METCALF, Z. P. 1962b. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 3. Gyponidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 229 p.
- METCALF, Z. P. 1962c. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 4. Ledridae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 147 p.
- METCALF, Z. P. 1962d. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 5. Ulopidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 101 p.
- METCALF, Z. P. 1962e. General catalog of the Homoptera. Fascicle VII Cercopoidea. Part 3. Aphrophoridae. North Carolina State College, Raleigh, USA. 600 p.
- METCALF, Z. P. 1962f. General catalog of the Homoptera. Fascicle VII Cercopoidea. Part 4. Clastopteridae. North Carolina State College, Raleigh, USA. 59 p.
- METCALF, Z. P. 1963a. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 6. Evancanthidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 63 p.
- METCALF, Z. P. 1963b. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 7. Nirvanidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 35 p.
- METCALF, Z. P. 1963c. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 8. Aphrodidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 268 p.
- METCALF, Z. P. 1963d. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 9. Hecalidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 123 p.
- METCALF, Z. P. 1963e. General catalog of the Homoptera. Fascicle VIII Cicadoidea. Part 1. Cicadidae. Section I Tibiceninae. North Carolina State College, Raleigh, USA. 585 p.
- METCALF, Z. P. 1963f. General catalog of the Homoptera. Fascicle VIII Cicadoidea. Part 1. Cicadidae. Section II Gaeninae and Cicadinae. North Carolina State College, Raleigh, USA. 586-919 p.
- METCALF, Z. P. 1963g. General catalog of the Homoptera. Fascicle VIII Cicadoidea. Part 2. Tibicinidae. North Carolina State College, Raleigh, USA. 492 p.
- METCALF, Z. P. 1964a. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 11. Coelidiidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 182 p.
- METCALF, Z. P. 1964b. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Bibliography of the Cicadelloidea. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 349 p.
- METCALF, Z. P. 1965a. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 1 Tettigellidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 730 p.
- METCALF, Z. P. 1965b. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 12. Eurymelidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 43 p.
- METCALF, Z. P. 1966a. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 13. Macropsidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 261 p.
- METCALF, Z. P. 1966b. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 14. Agalliidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 173 p.
- METCALF, Z. P. 1966c. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 15. Iassidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 229 p.
- METCALF, Z. P. 1966d. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 16. Idioceridae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 237 p.
- METCALF, Z. P. 1967. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 10. Euscelidae. Section I, II, II (three volumes). Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 2695 p.
- METCALF, Z. P. 1968. General catalog of the Homoptera. Fascicle VI. Cicadelloidea. Part 17. Cicadellidae. Agricultural Research Service, United States Dept. of Agriculture, Washington, USA. 1513 p.
- METCALF, Z. P.; WADE, V. 1963. A bibliography of the Membracoidea and fossil Homoptera (Homoptera: Auchenorrhyncha). North Carolina State College, Raleigh, USA. 200 p.
- METCALF, Z. P.; WADE, V. 1965a. A supplement to fascicle I - Membracidae of the general catalogue of the Hemiptera. Membracoidea, in two sections. Section I. Part 1 - Membracidae, Centrotinae, Platybelinae, Hoplophorioninae, Darninae. North Carolina State College, Raleigh, USA. 743 p.
- METCALF, Z. P.; WADE, V. 1965b. A supplement to fascicle I - Membracidae of the general catalogue of the Hemiptera. Membracoidea, in two sections. Section II. Part 1 - Membracidae (continued), Smilinae, Tragopinae, Membracinae. Part 2 - Aetalionidae. Part 3 - Biturritidae. Part 4 - Nicomiidae. North Carolina State College, Raleigh, USA. 745-1552 p.
- MILLER, D. R. 1984. Phylogeny and classification of the Margarodidae and related groups (Homoptera: Coccoidea), pp. 321-324. In: Kaszab, Z. (ed.). Verhandlungen des Zehnten Internationalen Symposiums über Entomofaunistik Mitteleuropas (SIEEC), Budapest, Hungary.
- MILLER, D. R.; BEN-DOV, Y. 2006. ScaleNet. United States Department of Agriculture. Available at: <<http://www.sel.barc.usda.gov/scalenet/scalenet.htm>>. [Accessed: August 2007].
- MILLER, D. R.; GIMPEL, M. E. 2000. A systematic catalogue of the Eriococcidae (felt scales) (Hemiptera: Coccoidea) of the World. Intercept, Andover, UK. 589 p.
- MILLER, D. R.; GIMPEL, M. E.; RUNG, A. 2005. A systematic catalogue of the Cercococcidae, Halimococcidae, Kermesidae, Micrococcidae, Ortheziidae, Phenacoleachiidae, Phoenicococcidae, and Stictococcidae (Hemiptera: Coccoidea) of the World. Intercept Limited, Wimborne, UK. 554.
- MILLER, D. R.; HODGSON, C. J. 1997. Phylogeny, pp. 229-250. In: Ben-Dov, Y., Hodgson, C. J. (eds.). Soft Scale Insects—Their Biology, Natural Enemies and Control, vol. A. Elsevier, Amsterdam, the Netherlands. 476 p.

- MILLER, D. R.; KOSZTARAB, M. 1979. Recent advances in the study of scale insects. *Annual Review of Entomology* 24: 1-27.
- MILLER, D. R.; RUNG, A.; VENABLE, G. L.; GILL, R. J. 2004. Scale families. An interactive key to the identification of families of scale insects (Hemiptera, Coccoidea). Available at: <<http://www.sel.barc.usda.gov/scalekeys/ScaleInsectsHome/ScaleInsectsFamilies.html>>. [Accessed: February 2008].
- MORRIS, D. C.; MOUND, L. A. 2003. Thysanoptera phylogeny - the molecular future, pp. 153-155. In: Klass, K-D. (ed.). Proceedings of the first Dresden meeting on Insect Phylogeny: "Phylogenetic relationships within insect orders" (Dresden, September 19-21, 2003). *Entomologische Abhandlungen* 61 (2): 119-172.
- MORSE, G. E.; NORMARK, B. B. 2006. A molecular phylogenetic study of armoured scale insects (Hemiptera: Diaspididae). *Systematic Entomology* 31 (2): 338-349.
- MOULDS, M. S. 2005. An appraisal of the higher classification of cicadas (Hemiptera: Cicadoidea) with special reference to the Australian fauna. *Records of the Australian Museum* 57: 375-446.
- MOUND, L. A.; HALSEY, S. H. 1978. Whitefly of the world. A systematic catalogue of the Aleyrodidae (Homoptera) with host plant and natural enemy data. *British Museum (Natural History)/John Wiley & Sons, Chichester, UK*. 340 p.
- MUIR, F. 1923. On the classification of the Fulgoroidea (Homoptera). *Proceedings of the Hawaii Entomological Society* 5 (2): 205-247.
- MURAJI, M.; TACHIKAWA, S. 2000. Phylogenetic analysis of water striders (Hemiptera: Gerroidea) based on partial sequences of mitochondrial and nuclear ribosomal RNA genes. *Entomological Science* 3 (4): 615-626.
- MYERS, J. G.; CHINA, W. E. 1929. The systematic position of the Peloridiidae as elucidated by a further study of the external anatomy of *Hemiodoecus leai* China. *Annals and Magazine of Natural History* 105 (3): 282-294.
- NIESER, N. 1975. The water bugs (Heteroptera: Nepomorpha) of the Guyana region. *Studies on the fauna of Suriname and other Guyanas* 16: 1-308, 24 pls.
- O'BRIEN, L. B.; WILSON, S. W. 1985. Planthopper systematics and external morphology, pp. 61-102. In: Nault, L. R., Rodriguez, J. G. (eds.). *The Leafhoppers and Planthoppers*. Wiley, New York, USA. 500 p.
- OMAN, P. W.; KNIGHT, W. J.; NIELSON, M. W. 1990. Leafhoppers (Cicadellidae): A bibliography, generic checklist and index to the world literature, 1956-1985. *CAB International, UK*. 368 p.
- ORTIZ-RIVAS, B.; MOYA, A.; MARTÍNEZ-TORRES, D. 2004. Molecular systematics of aphids (Homoptera: Aphididae): new insights from the long-wavelength opsin gene. *Molecular Phylogenetics and Evolution* 30: 24-37.
- OUVRARD, D. 2002. Systématique phylogénétique des Hemiptera Psylloidea: morphologie comparée du thorax et structures secondaires de l'ARNr 18S. *Bulletin de la Société zoologique de France* 127 (4): 345-357.
- OUVRARD, D.; CAMPBELL, B. C.; BOURGOIN, T.; CHAN, K. L. 2000. 18S rRNA secondary structure and phylogenetic position of Peloridiidae (Insecta, Hemiptera). *Molecular Phylogenetics and Evolution* 16 (3): 403-417.
- OUVRARD, D.; BURCKHARDT, D.; SOULIER-PERKINS, A.; BOURGOIN, T. 2008. Comparative morphological assessment and phylogenetic significance of the wing base articulation in Psylloidea (Insecta, Hemiptera, Sternorrhyncha). *Zoomorphology* 127: 37-47.
- PACKAUSKAS, R.; SCHAEFER, C. W. 1998. Revision of the Cryptocoridae (Hemiptera: Pentatomoidea). *Annals of the Entomological Society of America* 91: 363-386.
- PEDIGO, L. P. 1996. *Entomology and pest management*. Second Edition. Prentice Hall, Upper Saddle River, USA. 679 p.
- PEREIRA, D. L. V.; de MELO, A. L.; HAMADA, N. 2007. Chaves de identificação para famílias e gêneros de Gerromorpha e Nepomorpha (Insecta: Heteroptera) na Amazônia Central. *Neotropical Entomology* 36 (2): 210-228.
- POLHEMUS, J. T. 1985. Shore bugs (Heteroptera, Hemiptera; Saldidae). A world overview and taxonomy of Middle American forms. *The Different Drummer, Englewood, USA*. 252 pp.
- POPOV, Y. A.; SHCHERBAKOV, D. E. 1996. Origin and evolution of the Coleorrhyncha as shown by the fossil record, pp. 9-30. In: Schaefer, C. W. (ed.). *Studies on hemipteran phylogeny*. Proceedings, Thomas Say Publications in Entomology. Entomological Society of America, Lanham, USA. 244 p.
- QIN, T-K; GULLAN, P. J. 1995. A cladistic analysis of wax scales (Hemiptera: Coccoidea: Coccidae: Ceroplastinae). *Systematic Entomology* 20: 289-308.
- RAKITOV, R. A. 2002. Structure and function of the Malpighian tubules, and related behaviors in juvenile cicadas: evidence of homology with spittlebugs (Hemiptera: Cicadoidea & Cercopoidea). *Zoologischer Anzeiger* 241: 117-130.
- RÉDEI, D. 2007. A new species of the family Hypsipterygidae from Vietnam, with notes on the hypsipterygid fore wing venation (Heteroptera, Dipsocoromorpha). *Deutsche Entomologische Zeitschrift* 54 (1): 43-50.
- REMAUDIÈRE, G.; REMAUDIÈRE, M. 1997. *Catalogue des Aphididae du monde*. Institut National de la Recherche Agronomique, Paris, France. 473 p.
- REUTER, O. M. 1910. Neue beiträge zur phylogenie und systematik der miriden nebst einleitenden bemerkungen über die phylogenie der Heteropteren-Familien, *Acta Societatis Scientiarum Fennicae*, 37 (3): 1-171.
- RIEGER, C. 1976. Skelett und Muskulatur des Kopfes und Prothorax von *Ochterus marginatus* Laterille. *Zoomorphologie* 83: 109-191.
- ROSS, H. H. 1957. Evolutionary developments in the leafhoppers, the insect family Cicadellidae. *Systematic Zoology* 6 (2): 87-97 +69.
- ROY, L.; GUILBERT, E.; BOURGOIN, T. 2007. Phylogenetic patterns of mimicry strategies in Darnini (Hemiptera: Membracidae). *Annales de la Société Entomologique de France (Nouvelle série)* 43 (3): 273-288.
- SCHAEFER, C. W.; PANIZZI, A. R. 2000. *Heteroptera of economic importance*. CRC Press, Boca Raton, USA. 828 p.
- SCHLEE, v D. 1969a. Sperma-Übertragung (und anderen Merkmale) in ihrer Bedeutung für das phylogenetische System der Sternorrhyncha. *Phylogenetische Studien an Hemiptera I. Psylliformes (Psyllina + Aleyrodina) als monophyletische Gruppe*. *Zeitschrift für Morphologie der Tiere* 64: 95-138.
- SCHLEE, v D. 1969b. Die verwandtschaftsbeziehungen innerhalb der Sternorrhyncha aufgrund synapomorpher merkmale. *Phylogenetische studien an Hemiptera II: Aphidiformes (Aphidina + Coccina) als monophyletische Gruppe*. *Stuttgarter Beiträge zur Naturkunde* 199: 1-19.
- SCHLEE, v D. 1969c. Bau und funktion des aedeagus bei Psyllina und deren bedeutung für systematische und phylogenetische untersuchungen (Insecta, Hemiptera). *Phylogenetische Studien an Hemiptera III. Entkräftung eines Argument gegen die Monophylie der Sternorrhyncha*. *Zeitschrift für Morphologie der Tiere* 64: 139-150.
- SCHLEE, v D. 1969d. Morphologie und symbiose; ihre beweiskraft für die verwandtschaftsbeziehungen der Coleorrhyncha (Insecta, Hemiptera). *Phylogenetische studien an Hemiptera IV.: Heteropteroidea (Heteroptera + Coleorrhyncha) als monophyletische gruppen*. *Stuttgarter Beiträge zur Naturkunde* 210: 1-27.
- SCHUH, R. T. 1974. The Orhthotylinae and Phylinae (Hemiptera: Miridae) South Africa with a phylogenetic analysis of the antimimetic tribes of the two subfamilies for the world. *Entomologica Americana* 47: 1-332.
- SCHUH, R. T. 1976. Pretarsal structure in the Miridae (Hemiptera) with a cladistic analysis of relationships within the family. *American Museum Novitates* 2601: 1-39.

- SCHUH, R. T. 1979. [Review of] Evolutionary trends in Heteroptera. Part II. Mouthpart-structures and feeding strategies. *Systematic Zoology* 28 (4): 653-656.
- SCHUH, R. T. 1986. The influence of cladistics on heteropteran classification. *Annual Review of Entomology* 31: 67-93.
- SCHUH, R. T. 1995. Plant bugs of the world (Insecta: Heteroptera: Miridae). Systematic catalog, distributions, host list, and bibliography. The New York Entomological Society, New York, USA. 1329 p.
- SCHUH, R. T. 1997. [Review of] Studies on hemipteran phylogeny. *Journal of the New York Entomological Society* 104 (3-4): 231-235.
- SCHUH, R. T. 2000. Biological systematics. Principles and applications. Comstock Publishing Associates, Cornell University Press, Ithaca, USA. 236 p.
- SCHUH, R. T. 2006. Plant bugs of the World (Insecta: Heteroptera: Miridae). Systematic catalog. Available at: <<http://research.amnh.org/pbi/catalog/>>. [Accessed: July 2007].
- SCHUH, R. T.; POLHEMUS, J. T. 1980. Analysis of taxonomic congruence among morphological, ecological, and biogeographic data sets for the Leptopodomorpha (Hemiptera). *Systematic Zoology* 29 (1): 1-26.
- SCHUH, R. T.; SLATER, J. A. 1995. True bugs of the world (Hemiptera: Heteroptera). Classification and natural history. Cornell University Press, Ithaca, USA. 336 p.
- SCHUH, R. T.; ŠTYS, P. 1991. Phylogenetic analysis of cimicomorphan family relationships (Heteroptera). *Journal of the New York Entomological Society* 99 (3): 298-350.
- SCHUH, R.T.; CASSIS, G.; GUILBERT, E. 2006. Description of the first recent macropterous species of Vianaidinae (Heteroptera: Tingidae) with comments on the phylogenetic relationships of the family within the Cimicomorpha. *Journal of the New York Entomological Society* 114 (1-2): 38-53.
- SCHUH, R. T.; GALIL, B.; POLHEMUS, J. T. 1987. Catalog and bibliography of Leptopodomorpha (Heteroptera). *Bulletin of the American Museum of Natural History* 185 (3): 243-406.
- SCHUH, R. T.; WEIRAUCH, C.; HENRY, T. J.; HALBERT, S. 2008. Curaliidae, a new family of Heteroptera (Insecta: Hemiptera) from the Eastern United States. *Annals of the Entomological Society of America* 101 (1): 20-29.
- SCHUH, R. T.; WEIRAUCH, C.; WHEELER, W. C. *in press*. Phylogenetic relationships within the Cimicomorpha (Hemiptera: Heteroptera): a total evidence analysis. *Systematic Entomology*.
- SCUDDER, G. G. E. 1961. The comparative morphology of the insect ovipositor. *Transactions of the Royal Entomological Society of London* 113 (2): 25-40.
- SEEGER, W. 1975. Funktionsmorphologie an Spezialbildungen der Fühlergeißel von Psocoptera und anderen Paraneoptera (Insecta); Psocodea als monophyletische Gruppe. *Zeitschrift für Morphologie der Tiere* 81: 137-159.
- SHCHERBAKOV, D. E.; POPOV, Y. A. 2002. Superorder Cimicidea Laicharting, 1781. Order Hemiptera Linné, 1758. The bugs, cicadas, plantlice, scale insects, etc. (= Cimicida Laicharting, 1781, = Homoptera Leach, 1815 + Heteroptera Latreille, 1810), pp. 143-157. In: Rasnitsyn, A. P.; Quicke, D. L. J. (eds.). *History of Insects*. Kluwer Academic Publishers, the Netherlands. 517 p.
- SLATER, J. A. 1982. Hemiptera, pp. 417-447. In: Parker, S. P. (ed.). *Synopsis and classification of living organisms*. McGraw Hill, New York, USA. 1119 p.
- SLATER, J. A.; O'DONNELL, J. E. 1995. A catalogue of the Lygaeidae of the world (1960-1994). *New York Entomological Society*, New York, USA. 410 p.
- SORENSEN, J. T.; CAMPBELL, B. C.; GILL, R. J.; STEFFEN-CAMPBELL, J. D. 1995. Non-monophyly of Auchenorrhyncha ("Homoptera"), based upon 18S rDNA phylogeny: Eco-evolutionary and cladistic implications within pre-Heteropteroidea Hemiptera (s.l.) and a proposal for new monophyletic suborders. *Pan-Pacific Entomologist* 71 (1): 31-60.
- SOULIER-PERKINS, A. 2007. COOL: Cercopoidea organized online. Available at: <<http://rameau.snv.jussieu.fr/cool/index.php>>. [Accessed: February 2008].
- SOULIER-PERKINS, A.; SUEUR, J.; HOCH, H. 2007. Historical use of substrate-borne acoustic production within the Hemiptera: first record for an Australian Lophopid (Hemiptera, Lophopidae). *Australian Journal of Entomology* 46: 129-132.
- SPENCE, J. R.; ANDERSEN, N. M. 1994. Biology of water striders: interaction between systematics and ecology. *Annual review of Entomology* 39: 101-128.
- STONEDAHL, G. M.; DOLLING, W. R. 1991. Heteroptera identification: a reference guide, with special emphasis on economic groups. *Journal of Natural History* 25 (4): 1027-1066.
- ŠTYS, P. 1970. On the morphology and classification of the family Dipsocoridae s. lat., with particular reference to the genus *Hysipteryx* Drake (Heteroptera). *Acta Entomologica Bohemoslovaca* 67: 21-46.
- ŠTYS, P. 1983. A new family of Heteroptera with dipsocoromorphan affinities from Papua New Guinea. *Acta Entomologica Bohemoslovaca* 80: 256-292.
- ŠTYS, P. 1985. Soucasny stav beta-taxonomie radu Heteroptera. *Práce Slovenská entomologická spoločnosť SAV, Bratislava* 4: 205-235.
- ŠTYS, P. 1989. Phylogenetic systematics of the most primitive true bugs (Heteroptera, Enicocephalomorpha, Dipsocoromorpha) [in Czech]. *Práce Slovenská entomologická spoločnosť SAV, Bratislava* 8: 69-85.
- ŠTYS, P. 1998. Evolutionary origin of squamiform microsculpture on the forewing-holding devices (frenae) in Heteroptera. *Europena Journal of Entomology* 95: 307-310.
- ŠTYS, P.; JANSSON, A. 1988. Check-list of recent family-group names of Nepomorpha (Heteroptera) of the World. *Acta Entomologica Fennica* 50: 1-44.
- ŠTYS, P.; KERZHNER, I. M. 1975. The rank and nomenclature of higher taxa in recent Heteroptera. *Acta Entomologica Bohemoslovaca* 72: 65-79.
- SWEET, M. H. 1996. Comparative external morphology of the pregenital abdomen of the Hemiptera. p. 119-158. In: Schaefer, C. W. (ed.). *Studies on hemipteran phylogeny*. Proceedings, Thomas Say Publications in Entomology. Entomological Society of America, Lanham, USA. 244 p.
- SWEET, M. H. 2006. Justification for the Aradimorpha as an infraorder of the suborder Heteroptera (Hemiptera, Prosothrypa) with special reference to the pregenital abdominal structure. *Denisia* 19: 225 - 248.
- SWEET, M. H.; SCHAEFER, C. W. 2002. Parastrachiinae (Hemiptera: Cydnidae) raised to family level. *Annals of the Entomological Society of America* 95: 441-448.
- TAKIYA, D. M.; DMITRIEV, D. A. 2007. An interactive key to genera of the tribe Proconiini. Available at: <<http://ctap.inhs.uiuc.edu/takiya/key.asp?key=Proconia&lng=En&i=1&keyN=1>>. [Accessed: February 2008].
- TAKIYA, D. M.; TRAN, P. L.; DIETRICH, C. H.; MORAN, N. A. 2006. Co-cladogenesis spanning three phyla: leafhoppers (Insecta: Hemiptera: Cicadellidae) and their dual bacterial symbionts. *Molecular Ecology* 15: 4175-4191.
- TULLGREN, A. 1918. Zur Morphologie und Systematik der Hemipteren. I. Über das Vorkommen von s.g. Trichobothrien bei Hemiptera-Heteroptera und ihre mutmassliche Bedeutung für das Heteropterensystem. *Entomologisk Tidskrift* 39 (2): 113-132.
- URBAN, J. M.; CRYAN, J. R. 2007. Evolution of the planthoppers (Insecta: Hemiptera: Fulgoroidea). *Molecular Phylogenetics and Evolution* 42: 556-572.
- USINGER, R. L. 1943. A revised classification of the Reduvioidea with a new subfamily from South America (Hemiptera). *Annals of the Entomological Society of America* 36: 602-617.
- VARGAS-ROJAS, J. M.; FREYTAG, P.; SARMIENTO, C. E. 2006. Catalogo ilustrado de los cicadelinos de Colombia. Available at: <<http://www.unradio.unal.edu.co/ciencias/proyec>>

- tos/ins_cien/CATALOGO/pagina_ppal.htm>. [Accessed: September 2007].
- VON DOHLEN, C. D.; MORAN, N. A. 1995. Molecular phylogeny of the Homoptera: a paraphyletic taxon. *Journal of Molecular Evolution* 41 (2): 211-223.
- VON DOHLEN, C. D.; MORAN, N. A. 2000. Molecular data support a rapid radiation of aphids in the Cretaceous and multiple origins of host alternation. *Biological Journal of the Linnean Society* 71: 689-717.
- VON DOHLEN, C. D.; ROWE, C. A.; HEIE, O. E. 2006. A test of morphological hypotheses for tribal and subtribal relationships of Aphidinae (Insecta: Hemiptera: Aphididae) using DNA sequences. *Molecular Phylogenetics and Evolution* 38: 316-329.
- WALLACE, M. S.; DEITZ, L. L. 2004. Phylogeny and systematics of the treehopper subfamily Centrotinae (Hemiptera: Membracidae). *Memoirs on Entomology International* 19: 1-377.
- WALLACE, M. S.; DEITZ, L. L. 2005. Australian treehoppers (Hemiptera: Membracidae: Centrotinae: Terentiini): phylogeny and biogeography. *Invertebrate Systematics* 20: 163-183.
- WEBER, H. 1930. *Biologie der hemipteren*. Julius Springer, Berlin, Germany. 543 p.
- WEIRAUCH, C. 2008. Cladistic analysis of Reduviidae (Hemiptera: Cimicomorpha) based on morphological characters. *Systematic Entomology* 33: 229-274.
- WHEELER, W. C.; SCHUH, R. T.; BANG, R. 1993. Cladistic congruence among higher groups of Heteroptera: congruence between morphological and molecular data sets. *Entomologica Scandinavica* 24: 121-137.
- WHEELER, W. C.; WHITING, M.; WHEELER, Q. D.; CARPENTER, J. C. 2001. The phylogeny of the extant Hexapod orders. *Cladistics* 17: 113-169.
- WHITE, I. M.; HODKINSON, I. D. 1985. Nymphal taxonomy and systematics of the Psylloidea (Homoptera). *Bulletin of the British Museum (Natural History), Entomology series* 50 (2): 153-301.
- WHITING, M. F. 2002. Phylogeny of the holometabolous insect orders: molecular evidence. *Zoologica Scripta* 31: 3-15.
- WHITING, M. F.; CARPENTER, J. C.; WHEELER, Q. D.; WHEELER, W. C. 1997. The Strepsiptera problem: phylogeny of the holometabolous insect orders inferred from 18S and 28S ribosomal DNA sequences and morphology. *Systematic Biology* 46 (1): 1-68.
- WILLIAMS, D. J.; GRANARA DE WILLINK, M. C. 1992. *Mealybugs of Central and South America*. CAB International, UK. 635 p.
- WILLMANN, R. 2004. Phylogenetic relationships and evolution of insects, pp. 330-344. In: Cracraft, J.; Donoghue, M. J. (eds.). *Assembling the tree of life*. Oxford University Press, New York, USA. 592 p.
- WILSON, S. E. 2005. Keys to the families of Fulgoromorpha with emphasis on planthoppers of potential economic importance in the Southeastern United States (Hemiptera: Auchenorrhyncha). *Florida Entomologist* 88 (4): 464-481.
- WOOTTON, R. J. 1965. Evidence for tracheal capture in early Heteroptera. *Proceedings of the 12th International Congress of Entomology* 65-67.
- WOOTTON, R. J.; AND C. R. BETTS, C. R.. 1986. Homology and function in the wings of Heteroptera. *Systematic Entomology* 11 (3): 389-400.
- WYGODZINSKY, P. 1950. Contribution towards the knowledge of the family "Cryptostemmatidae" (Hemiptera). *Revista Brasileira de Biologia* 10: 377-392.
- WYGODZINSKY, P.; SCHMIDT, K. 1991. Revision of the new world Enicocephalomorpha (Heteroptera). *Bulletin of the American Museum of Natural History* 200: 1-265.
- YANG, C-T. 2004. The phylogeny of Hemiptera (Homoptera-Heteroptera). *Journal of Agriculture and Forestry* 53 (1): 51-78.
- YEH, W. B.; YANG, C. T. 1999. Fulgoromorpha phylogeny based on 28S rDNA nucleotide sequence. *Chinese Journal of Entomology* 11: 87-111.
- YEH, W. B.; YANG, C. T.; HUI, C. F. 1998. Phylogenetic relationships of the Tropiduchidae-group (Homoptera: Fulgoroidea) of planthoppers inferred through nucleotide sequences. *Zoological Studies* 37 (1): 45-55.
- YEH, W. B.; YANG, C. T.; HUI, C. F. 2005. A molecular phylogeny of planthoppers (Hemiptera: Fulgoroidea) inferred from mitochondrial 16S rDNA sequences. *Zoological Studies* 44 (4): 519-535.
- YOSHIZAWA, K. 2007. The Zoraptera problem: evidence for Zoraptera + Embiodea from the wing base. *Systematic Entomology* 32: 197-204.
- YOSHIZAWA, K.; JOHNSON K. P. 2005. Aligned 18S for Zoraptera (Insecta): Phylogenetic position and molecular evolution. *Molecular Phylogenetics and Evolution* 37: 572-580.
- YOSHIZAWA, K.; JOHNSON K. P. 2006. Morphology of male genitalia in lice and their relatives and phylogenetic implications. *Systematic Entomology* 31: 350-361.
- YOSHIZAWA, K.; SAIGUSA, T. 2001. Phylogenetic analysis of paraneopteran orders (Insecta: Neoptera) based on forewing base structure, with comments on monophyly of Auchenorrhyncha (Hemiptera). *Systematic Entomology* 26: 1-13.
- ZAHNISER, J. N. 2007. An interactive key to Deltocephaline-related tribes. Available at: <<http://ctap.inhs.uiuc.edu/zahniser/key.asp?key=Delt&lng=En&i=1&keyN=1>>. [Accessed: February 2008].
- ZAHNISER, J. N.; DIETRICH, C. H. 2008. Phylogeny of the leafhopper subfamily Deltocephalinae (Insecta: Auchenorrhyncha: Cicadellidae) and related subfamilies based on morphology. *Systematics and Biodiversity* 6 (1): 1-24.
- ZRZAVÝ, J. 1990. Evolution of Hemiptera: an attempt at synthetic approach., pp. 19-22. In: 6th international Symposium of Scale Insects Studies, Cracow.
- ZRZAVÝ, J. 1992. Evolution of antennae and historical ecology of the hemipteran insects (Paraneoptera). *Acta Entomologica Bohemoslovaca* 89 (2): 77-86.

Recibido: 12-dic-2007 • Aceptado: 21-mar-2008