

FIRST RECORD OF THE GENUS *Saccharosydne* Kirkaldy 1907 (HEMIPTERA - FULGOROMORPHA - DELPHACIDAE) IN ARGENTINA

María F. Rossi Batiz and Ana M. M. de Remes Lenicov

SUMMARY

A new genus, *Saccharosydne* Kirkaldy, is reported in Argentina. This is the most economically important representative of the tribe *Saccharosydnini*. The geographical distribution of all

species and some morphological and biological relevant data of this genus are added. This new record represents the southernmost limit of the tribe in the Americas.

PRIMER REGISTRO DEL GÉNERO *Saccharosydne* Kirkaldy 1907 (HEMIPTERA - FULGOROMORPHA - DELPHACIDAE) EN ARGENTINA

María F. Rossi Batiz and Ana M. M. de Remes Lenicov

RESUMEN

Se registra por primera vez el género *Saccharosydne* Kirkaldy en Argentina. Éste es económicamente el más importante representante de la tribu *Saccharosydnini*. Se agrega la distribución

geográfica de todas las especies y algunos datos morfológicos y biológicos relevantes. Este nuevo registro representa el límite más austral de la tribu en América.

PRIMEIRO REGISTRO DO GÊNERO *Saccharosydne* Kirkaldy 1907 (HEMIPTERA - FULGOROMORPHA - DELPHACIDAE) NA ARGENTINA

María F. Rossi Batiz e Ana M. M. de Remes Lenicov

RESUMO

Registra-se por primeira vez o gênero *Saccharosydne* Kirkaldy na Argentina. Este é econômica e o mais importante representante da tribo *Saccharosydnini*. Soma-se à distribuição geo-

gráfica de todas as espécies e alguns dados morfológicos e biológicos relevantes. Este novo registro representa o limite mais austral da tribo na América.

Introduction

The Delphacidae family is the largest and most economically important one amongst the Fulgoroidea, Delphacinae being the most diverse subfamily. According to Asche (1985), the subfamily groups three tribes: Delphacini, Saccharosydnini and Tropidocephalini. All the 3 genera of the Saccharosydnini tribe, *Neomalaxa* Muir, 1918, *Pseudomacrocorupha* Muir, 1930 and *Saccharosydne*

Kirkaldy, 1907, are registered in tropical and subtropical regions of America and Asia (Asche, 1985), being *Pseudomacrocorupha* the only one recorded up to date in Argentina.

The intensification of new cultural practices in Argentina for productivity improvement has greatly changed the vegetal physiognomy, leading to the establishment and increment of foreign insect pest populations. Because of the finding of three possible spe-

cies of tropical *Saccharosydne* in recent collections on garlic (*Allium sativum* L.), rye (*Secale cereale* L.) and foxtail (*Cortaderia spp.*) from Mendoza, San Juan and Neuquén provinces of Argentina, the purpose of this contribution is to record in the country this genus for the first time, adding the world distribution and some morphological and biological data of this economically important genus.

The Argentinean specimens were field collected using wa-

ter traps and nets, and hand captured. They were preserved in the Museo de La Plata collection (MLP). Materials from the collections of MLP and the British Museum (Natural History) (BM), London, were also examined.

Genus: *Saccharosydne* Kirkaldy, 1907

Genotype: *Delphax saccharivora* Westwood, 1833: 413, by original designation.

Saccharosydne is distinguished mainly by slender, elongated forms; head much

KEYWORDS / Argentina / Geographical Distribution / Morphology / *Saccharosydne* /

Received: 02/11/2008. Modified: 12/19/2008. Accepted: 12/19/2008.

María F. Rossi Batiz. Biologist, Universidad Nacional de La Plata (UNLP), Argentina. Professor, División Entomología, Facultad de Ciencias Naturales

y Museo (UNLP), Argentina. e-mail: mfrb@fcnym.unlp.edu.ar

Ana M. Marino de Remes Lenicov. Doctor in Natural Scienc-

es, UNLP, Argentina. Professor UNLP, Argentina. Researcher, CONICET, Argentina. Address: División Entomología, Facultad de Ciencias Naturales y Museo

(UNLP), Paseo del Bosque S/N (1900) La Plata, Buenos Aires, Argentina. e-mail: amarino@fcnym.unlp.edu.ar

narrower than pronotum, angular in profile; vertex much extended before eyes; frons about twice as long as width with the median carinae not forked; antennae short, reaching the base of clypeus. Anal segment of male without processes.

Although all species have elongated, slightly flattened forms, with elongated fore wings and general light green-yellowish coloration, the male genitalia is further distinctive, particularly by the large development of the aedeagus. It consists of a horizontal, slightly curved tube directed posteriorly with a stout spine, curving upward distally, joined to a long lightly sclerotized whiplike filament inward directed, strongly elongated, coiled within the pygofer reaching far cephalad into the abdomen.

Muir and Giffard (1924) separated this genus and the monotypic *Neomalaxa* from all other Delphacidae by the male genitalia, and from *Stenocranus* Fieber, 1866 by the terminalia build, as well as the vertex carinae disposition.

All the seven species of this genus are distinguished from all other Delphacidae principally by the combined characteristics of the pygofer, aedeagal complex and anal segment.

Geographical Distribution

So far, with the exception of two species in Asia, this genus is confined to America. *S. brevirostris* Muir, 1926, in Ecuador; *S. gracilis* Muir, 1926 and *S. ornatipennis* Muir, 1926, in Brazil; *S. procerus* Matsumura, 1931, in Japan, China, USSR (Maritime and Mainland Territory); *S. rostifrons* (Crawford, 1914), in Cuba; *S. saccharivora* (Westwood, 1833), in Antigua, Antillas, Barbados, Cuba, British Guiana, Grenada, Haití, Jamaica, Puerto Rico, Santo Domingo, Trinidad, USA and Venezuela (Guagliumi, 1953; Giraldo-Vanegas *et al.*, 2004), Colombia (Gómez and Lastra Borja, 1995) and Cuba (Arocha *et al.*, 2005). It causes direct damage to the plant during feeding and oviposition, and indirect damage when the populations are big and the sugary excretes allow the development of fumagina (*Fumago sacchari* Speg.), impeding normal photosynthesis and transpiration (Guagliumi, 1953; Giraldo-Vanegas *et al.*, 2004, 2005, 2006). This pest acquires more importance because of its displayed vector capacity of the Sugarcane Yellow Leaf Phytoplasma (SCYLP), caus-

bias (Gómez and Lastra Borja, 1995), Grenada (Westwood, 1833) and Cuba (Arocha *et al.*, 2005); and *S. viridis* Muir, 1926, in British Guiana (Figure 1). The collections in Argentina extends to the south the known geographic distribution.

Host Plants

Several grasses and gramineous plants have been reported as hosts of the genus *Saccharosydne*. *S. brevirostris* was recorded on grass on the shore of the Napo River, Ecuador (Muir, 1926); *S. ornatipennis* on *Paspalum intermedium* Munro, from Rezende, Rio de Janeiro, Brazil (Muir, 1926); *S. gracilis* off rank grass of Corcovado, Rio de Janeiro, Brazil (Muir, 1926); *S. viridis* on rice and rice grass of Blairmont, British Guiana (Muir, 1926); *S. procerus* on rice fields and in neighboring water-bamboo *Zizania caduciflora* L. habitats (XiaoPing *et al.*, 2005); *S. saccharivora* on sugarcane *Saccharum officinarum* L. (Westwood, 1833; Guagliumi, 1953; Giraldo-Vanegas *et al.*, 2004, 2005, 2006).

Phytosanitary Importance

Among *Saccharosydne* species, only *S. saccharivora* is well known as a sugarcane pest in Antigua, Antillas, Barbados, Cuba, British Guiana, Grenada, Haití, Jamaica, Puerto Rico, Santo Domingo, Trinidad, USA and Venezuela (Guagliumi, 1953; Giraldo-Vanegas *et al.*, 2004), Colombia (Gómez and Lastra Borja, 1995) and Cuba (Arocha *et al.*, 2005). It causes direct damage to the plant during feeding and oviposition, and indirect damage when the populations are big and the sugary excretes allow the development of fumagina (*Fumago sacchari* Speg.), impeding normal photosynthesis and transpiration (Guagliumi, 1953; Giraldo-Vanegas *et al.*, 2004, 2005, 2006). This pest acquires more importance because of its displayed vector capacity of the Sugarcane Yellow Leaf Phytoplasma (SCYLP), caus-




Figure 1. Distribution map of genus *Saccharosydne* in America. The countries where *Saccharosydne* species have been registered are 1: United States of America, 2: Bermudas, 3: Cuba, 4: Jamaica, 5: Haití, 6: Santo Domingo, 7: Puerto Rico, 8: Antigua, Antillas, Santa Lucía, 9: Barbados, 10: Grenada, 11: Trinidad, 12: Venezuela, 13: Colombia, 14: British Guiana, 15: Ecuador, 16: Brazil, 17: Argentina.

ing agent of the Sugarcane Yellow Leaf Syndrome (YLS) in Cuba. This infection, characterized by a yellowing of the midrib and lamina, was first reported in the 1960s in East Africa and later in Hawaii, South Africa and Cuba. It is now widely distributed in most sugarcane-growing countries from all continents (Arocha *et al.*, 2005).

Natural Enemies

As natural enemies of *S. saccharivora* are mentioned several species of insects: egg parasitoids (two species of Hymenoptera); nymphs and adults parasitoids (one species of Hymenoptera and one species of Strepsiptera) (Guagliumi, 1953; Terán, 1980; Giraldo-Vanegas *et al.*, 2004); predators of eggs, nymphs and adults (one species

of Diptera, six of Coleoptera, five of Neuroptera, three of Hemiptera, one species each of Dermaptera and Orthoptera), and the pathogen fungi *Metarrhizium* sp. and *Fusarium* sp. (Guagliumi, 1953). *Anagrus* spp. (Hymenoptera) has been mentioned as eggs parasitoid of *S. procerus* and *S. rostifrons* (XiaoPing, 2000; Triapitsyn, 2002). No predator or evident parasitoid has been found to date in the material from Argentina.

Remark

Among the Argentinean specimens three morphospecies can be distinguished taking into account the relative body measurements, the shape of the posterior margin of pygofer, parameres and aedeagus. These are very similar to *S. gracilis*, but differ

in the anal angles of pygofer caudally produced and the apical shape of the parameres. Further studies are required in order to clarify the identity of these local specimens.

Reference Specimens

11 ♂♂ and 10 ♀♀, "La Consulta", Mendoza, Argentina, with water trap on garlic crop, 22/VIII, 5/IX, 26/IX and 1/X/03; 23-27/VIII, 27/IX-01/X and 04-08/X/04; 14/IX, 21/IX and 20/X/05; 20/IX and 12/X/06. Lanatti leg.

Other examined materials from Argentina: 1 ♀, Dto. Pocito, San Juan, 23/I/64, Torres-Ferreyyra leg.; 1 ♂, 1 ♀, Dto. Pocito, San Juan, with net on rye, 13-17/IX/04, Meneguzzi leg; 9 ♂, 5 ♀ 30/IX/02; 18/VII, 22/VIII, 05/IX, 12/IX, 26/IX and 07/XI/03, 6 ♀ 9-13/VIII, 20-24/IX, 27/IX-1/X, 04-08/X, 29-03/XII and 13-17/XII/04, 4 ♂, 1 ♀ 13/IX, 15/IX, 18/IX, 02/X and 12/X/06, "La Consulta", Mendoza, with water trap on garlic, Lanatti leg.; 5 ♂, 9 ♀, 1 nymph, hand captured on foxtail, San Martín de los Andes, Neuquén, 26/II/07, Logarzo leg.

Examined materials from the BM collection: 1 ♂ *Oxycranus procerus* Matsumura, Nagasaki, Japan (1913), F. Muir coll., BM 1932-279; 1 ♂ *S. procerus* (Matsumura), Foochow, China (1935), MS Yang leg., Pres. by Com. Inst. Ent. BM 1948-548; 1 ♀ *S. procerus* (Matsumura), Nantou, Taiwan

(1984), CT Yang col. and det.; 1 ♂ *S. viridis* Muir, paratype, (1923), on rice Blairmont, FX Williams leg., BM 1929-293; 1 ♂, 1 ♀ *S. gracilis* Muir, paratypes, Río de Janeiro, Brazil (1924), FX Williams leg., BM 1929-293; 1 ♀ *S. brevirostris* Muir, paratype, ex beach grass, Napo River, Ecuador (1923), FX Williams leg., BM 1929-293; 1 ♀ *S. ornatipennis* Muir, paratype, on grass, Rezende, Brazil (1924), FX Williams leg., BM 1929-293; 1 ♀ *S. cf. rostifrons*, Jacareacanga, Brazil (1984), M Alvarenga leg., M Asche det., BM 1971-165; 1 ♂ *S. rostifrons* on *Paspalum conjugatum*, Cayo District Central Farm, Honduras, E García leg., Pres. by Com. Inst. Ent., BM 1988-26; 1 ♂ *S. saccharivora* (Westwood), Bermuda Sandys P, Fort Scaur (1988), MR Wilson & D. J Hilburn leg., MR Wilson det.; 1 ♂ *S. saccharivora* (Westwood) on sugarcane, Grenada (1961), FD Bennett leg., MSK Ghauri det., Pres. by Com. Inst. Ent., BM 1961-6.

ACKNOWLEDGMENTS

The authors thank Silvio Lanatti, Natalia Meneguzzi, and Guillermo Logarzo for collecting and providing specimens for this study. Special thanks to Mick Webb for the access to the material housed in the collection of the British Museum (Natural History), London. The work was supported by FONCyT (PICT/03 08-15219), CONICET and UNLP.

REFERENCES

- Arocha Y, López M, Fernández M, Piñol B, Horta D, Peralta EL, Almeida R, Carvajal O, Picornell S, Wilson MR, Jones P (2005) Transmission of a sugarcane yellow leaf phytoplasma by the delphacid planthopper *Saccharosydne saccharivora*, a new vector of sugarcane yellow leaf syndrome. *Plant Pathol.* 54: 634-642.
- Asche M (1985) Zur Phylogenie der Delphacidae Leach, 1815 (Homoptera Cicadina Fulgoromorpha). *Marburg. Entomol. Publik.* 2: 1-910.
- Giraldo-Vanegas H, Vargas A, Lindarte JO (2004) El saltahojas verde de la caña de azúcar en el Estado Táchira. *INIA Divulga N° 3*. Venezuela. pp. 25-27.
- Giraldo-Vanegas H, Nass H, Hernández E, Amaya F, Vargas A, Ramírez M, Ramírez F, Ramón M, Lindarte JO (2005) Incidencia del saltahojas verde de la caña en siete cultivares de caña de azúcar, Estado Táchira, Venezuela. *Agron. Trop.* 56: 553-567.
- Giraldo-Vanegas H, Vargas A, Sarmiento A, Hernández E, Amaya F, Ramírez M, Ramírez F, Contreras EJ (2006) Evaluación de bioplaguicidas para el manejo del saltahojas verde de la caña de azúcar *Saccharosydne saccharivora* (Westwood) (Hemiptera: Delphacidae), en el valle San Antonio-Ureña, Táchira, Venezuela. *Agron. Trop.* 56: 253-276.
- Gómez LA, Lastra Borja LA (1995) Insectos asociados con la caña de azúcar en Colombia. In *El Cultivo de la Caña en la Zona Azucarera de Colombia. CENICANA*. Cali, Colombia. pp. 237-263.
- Guagliumi P (1953) *El saltahojas de la caña de azúcar Saccharosydne saccharivora Westwood y la funagrina en Venezuela*. Boletín Técnico N° 7. Instituto Nacional de Agricultura. Ministerio de Agricultura y Cría. Caracas, Venezuela. 82 pp.
- Muir F (1926) *Contributions to Our Knowledge of South American Fulgoroidea (Homoptera). Part I. The Family Delphacidae*. Bulletin N° 18. Hawaiian Sugar Planters' Association. Honolulu, HI, USA. 51 pp.
- Muir F, Giffard WM (1924) *Studies in North American Delphacidae (Homoptera)*. Bulletin N° 15. Hawaiian Sugar Planters' Association. Honolulu, HI, USA. 50 pp.
- Terán BJ (1980) Lista preliminar de Hymenoptera parásitos de otros insectos en Venezuela. *Rev. Fac. Agron.* II: 283-389.
- Triapitsyn SV (2002) Descriptive notes on a new and other little known species of *Anagrus* Haliday, 1833 (Hymenoptera: Mymaridae) from the New World tropics and subtropics. *Entomotropica - Bol. Entomol. Venez.* 17: 213-223.
- Westwood JO (1833) Additional observations upon the insect which infests the sugar canes in Grenada. *Mag. Nat. Hist. - J. Zool., Bot., Mineral., Geol. Meteorol.* 6: 409-413.
- XiaoPing Y (2000) Role of *Saccharosydne procerus* on *Zizania caduciflora* as an alternate host for *Anagrus nilaparvatae*, the egg parasitoid of the brown planthopper *Nilaparvata lugens*, which attacks temperate rice. In *Exploiting Biodiversity for Sustainable Pest Management*. Proc. Impact Symposium on Exploiting Biodiversity for Sustainable Pest Management. Kunming, China. ISBN 971-22-0156-2.
- XiaoPing Y, JianMing C, ZhongXian L, XuSong Z, HongXin X, Jue-Feng Z (2005) Habitat manipulation in sustainable pest management in the rice ecosystem of the Yangtze River Delta. In *Rice is Life: Scientific Perspectives for the 21st Century*. Proc. World Rice Research Conference. Tsukuba, Japan. pp. 470-472.