

# Biological Studies of Insects Feeding on the Kudzu Plant, *Pueraria lobata* (Leguminosae) I. List of Feeding Species

Jiraporn TAYUTIVUTIKUL\* and Kanetosi KUSIGEMATI

(Laboratory of Plant Pathology and Entomology)

Received for Publication September 9, 1991

## Introduction

Kudzu, *Pueraria lobata* (Willd.) Ohwi, is a vigorous perennial vine which is widely distributed in the temperate and tropical regions in Asia<sup>24</sup>). The plant has both useful and harmful phases from an applied point of view. The tuber containing starch can be made into flour for food or utilized for medical purposes, and the leaves are used for fodder. In ancient times, textiles were made from the stalk<sup>24</sup>). The Chinese crude drug “Gehua”, used as a folk medicine in China, Taiwan, Korea and Japan, is considered to be the flower of *Pueraria* spp.<sup>27</sup>). Several chemical substances isolated from the callus tissues induced on the stem are also utilized in various medical treatments<sup>56,60</sup>).

The plant, however, has now run wild, becoming a serious weed on fallow land, riverbanks and roadsides in Japan. The plant, introduced in the United States about a century ago from Japan to prevent soil erosion<sup>19,49,54</sup>), has been both of benefit and a nuisance and is now out of control there also<sup>58</sup>). Kudzu was used as a food and fuel<sup>61</sup>) in addition to its benefits as an erosion protective agent, but now has become a noxious weed that displaced local flora in several areas<sup>5,6,49</sup>).

Van Emden<sup>65,66</sup>) aroused interest in wild plants for crop pest management, and the Kudzu plant may play an important role in this since it is widely distributed around agroecosystems.

Despite its important status, the Kudzu plant has not been studied for its associated insects, nor for a means of biological control<sup>17</sup>) which has not been succeeded owing to its large tuber roots though chemical control of the plant has been tried also in both countries mentioned<sup>37</sup>). It was thus believed valuable to conduct fundamental studies on the insects associated with the plant for the purposes mentioned above.

Following a preliminary survey on insects of the plant made in Yamaguchi City in 1984<sup>18</sup>), the first author began biological studies of its dominant phytophagous bugs in the same area in 1986 and 1987 and reported the biological data of two species<sup>62,63</sup>).

The present study was carried on in Kagoshima City to accumulate further biological data on Kudzu-feeding species for future application. Among feeders found in the Kagoshima area, 18 insect species and 1 mite species were selected, and their seasonal abundance and habitat or within-plant distribution were studied.

The insect species belong to the following 4 orders: Hemiptera (5 species), Coleoptera (5

---

\* Present address: National Biological Control Research Center, Kasetsart University and National Research Council of Thailand, Bangkok 10900, Thailand

species), Diptera (3 species) and Lepidoptera (5 species). Among those, four leaf-miner species, i.e.,—*Japanagromyza tristella*, *Liocrobyla lobata*, *Neolithocolletis hikomonticola* and *Spulerina dissotoma*—were intensively studied for their biology and distribution within the leaf since they are expected candidates as biological control agents.

Three other species—*Homoeocerus unipunctatus*, *Hedylepta misera* and *Matsumuraeses falcana*—were studied for their development since they were found to be dominant in the survey area and to attack other leguminous plants.

This study was thus designed to elucidate the following: (1) species composition of Kudzu-feeding insects, (2) seasonal abundance and habitat on the plant of the selected species, (3) biology and distribution within the leaf of the four leafminers, (4) biological data including developmental period, developmental zero, total effective temperature, adult longevity and fertility of the three dominant species.

In this paper, species composition of Kudzu-feeding species is enumerated, and remaining topics together with the methods and materials used will be reported in a second paper as well as general remarks about the overall studies.

## Materials and Methods

### Sampling procedures

Samplings of phytophagous insects were conducted during 1988–1989 at three Kudzu growing sites in Kagoshima City: Korimoto graveyard, Shimota and Toso. When immature and mature insects were discovered, they were hand-picked or aspirated from Kudzu plants on the field at one week intervals, with 1 or 2 hours usually spent at each site. The affected plant parts were collected together with the suspected feeding insects, placed in plastic bags and brought back to the laboratory for rearings. Insects resting on the plants and pollinators were not included in this study.

Another field investigation was conducted at two sites, Korimoto graveyard and Toso, from early May to early December in 1988 and from late March to early December in 1989 to monitor seasonal abundance of the feeding species. Data on the feeding species composition obtained in this investigation are also referred to here, though the method used together with that for samplings to determine seasonal abundance, rearing and development will be described in Part 2.

## Results and Discussion

### 1. List of feeding species

A total of 71 species of Insecta and 2 species of Arachnoidea have been previously recorded from the Kudzu plant, *Pueraria lobata* (Willd.) Ohwi from Japan and USA so far as we are aware. In the present study, a total of 61 species of Insecta and one species of Arachnoidea feeding on Kudzu plants, including newly found species were observed in Kagoshima City. Among them, a total of 38 species belonging to 26 families of 5 insect orders are recorded for the first time from this plant.

Those known and the newly found species, a total of 109 species belonging to 46 families of 6 insect orders and 2 species of Acarina, are listed and shown in Table 1 and Fig. 1.

The number of species in each order is as follows: Orthoptera (5%)—6 species of 4 families, Hemiptera (Homoptera) (10%)—11 species of 10 families, Hemiptera (Heteroptera) (23%)—26






Fig. 1. Percentage of phytophagous species of Kudzu plant belonging to different orders, A: Known feeding species based on published records and those found in Kagoshima City, B: Feeding species found in Kagoshima City.

species of 8 families, Thysanoptera (4%)—4 species of 1 family, Coleoptera (22%)—24 species of 8 families, Diptera (5%)—6 species of 2 families, Lepidoptera (29%)—32 species of 13 families, Acarina (2%)—2 species of 1 family. Hemiptera, Coleoptera and Lepidoptera are dominant among the order recorded.

Most of the species listed in Table 1 are polyphagous and attack other plants such as soybean, *Glycine max* Merr., adzuki bean, *Vigna angularis* (Willd.) Ohwi et Ohashi var. *angularis*, wisteria, *Wistaria brachybotrys* Sieb. et Zucc., and other leguminous plants. Of the 62 species found in Kagoshima City, 38 are foliage-feeding including gall-makers and leafminers, 23 are sap-feeding and only 1 species is endophagous on stored seed. Nineteen species of which biology is described in Part 2 are comparatively dominant. Only 14 among the Kudzu-feeding species listed below are monophagous on Kudzu plant so far as we know.

#### Hemiptera

Miridae: *Proboscidoecoris varicornis* (Jakovlev)

#### Thysanoptera

Thripidae: *Salpingothrips aimatofus* Kudo

#### Coleoptera

Buprestidae: *Trachys auricollis* E. Saunders

Curculionidae: *Mesalcidodes trifidus* (Pascoe)

#### Diptera

Cecidomyiidae: *Lasioptera puerariae* Shinji; *Pitydiplosis* sp.; Genus sp.

Agromyzidae: *Ophiomyia puerarivora* Sasakawa

#### Lepidoptera

Lyonetiidae: *Leucoptera puerariella* Kuroko

Gracillariidae: *Liocrobyla lobata* Kuroko; *Neolithocolletis hikomonticola* Kumata

Pyrilidae: *Hedylecta misera* (Butler); *Pleuroptya punctimarginalis* (Hampson)

Noctuidae: *Microxylla confusa* (Wileman)

## 2. Newly recorded species from Kudzu plant

A total of 38 species belonging to 26 families of 5 insect orders were among those collected and are recorded here for the first time from Kudzu plant; data on collection follows. The words—Korimoto, Shimota and Toso—identify the locality in Kagoshima City where collection was made, and all specimens were collected by the first author (JT).

### Insecta

#### Orthoptera

##### Tettigoniidae

###### *Gampsocleis buergeri* (de Haan)

Specimen examined. Toso: 1 ex., 9. v. 1988.

##### Pyrgomorphidae

###### *Atractomorpha lata* (Motschulsky)

Specimens examined. Korimoto: 1 ex., 5. ix. 1988. Shimota: 1 ex., 8. ix. 1988; 1 ex., 22. ix. 1988. Toso: 1 ex., 2. ix. 1988; 1 ex., 9. ix. 1988; 2 exx., 22. ix. 1988.

##### Acrididae

###### *Acrida cinerea* (Thunberg)

Specimen examined. Korimoto: 1 ex., 5. viii. 1988.

###### *Oxya japonica japonica* (Thunberg)

Specimen examined. Toso: 1 ex., 22. ix. 1988.

###### *Patanga japonica* (Bolivar)

Specimens examined. Korimoto: 1 ex., 11. vi. 1988; 1 ex., 4. vii. 1988. Toso: 1 ex., 4. vi. 1988; 1 ex., 17. vi. 1988.

##### Tetrigidae

###### *Tetrix japonica* (Bolivar)

Specimen examined. Toso: 1 ex., 9. v. 1988.

### Hemiptera

#### Flatidae

###### *Geisha distinctissima* (Walker)

Specimens examined. Korimoto: 3 exx., 13. vi. 1988. Toso: 1 ex., 24. vi. 1988. Shimota: 2 exx., 27. vi. 1988.

#### Ricaniidae

###### *Orosanga japonica* (Melichar)

Specimens examined. Korimoto: 5 exx., 27. vi. 1988. Shimota: 3 exx., 1. vi. 1988. Toso: 2 exx., 19. viii. 1988.

#### Cercopidae

###### *Eoscartopis assimilis* (Uhler)

Specimens examined. Korimoto: 1 ex., 20. v. 1988. Shimota: 2 exx., 15. vi. 1988

#### Penthimiidae

###### *Penthimia nitida* Lethierry

Specimen examined. Shimota: 1 ex., 24. viii. 1988.

#### Cicadellidae

###### *Bothrogonia ferruginea* (Fabricius)

Specimens examined. Shimota: 2 exx., 15. vi. 1988.

###### *Cicadella viridis* (L.)

Specimens examined. Korimoto: 2 exx., 22. viii. 1988. Shimota: 3 exx., 24. viii. 1988.

#### Typhlocybidae

*Empoasca* sp.

Specimens examined. Shimota: 3 exx., 1. vi. 1988.

#### Miridae

*Adelphocoris triannulatus* (Stål)

Specimens examined. Shimota: 1 ex., 8. ix. 1988. Toso: 1 ex., 1. v. 1988.

#### Berytidae

*Metacanthus pulchellus* Dallas

Specimens examined. Shimota: 1 ex., 19. x. 1988. Toso: 1 ex., 11. vi. 1988; 2 exx., 24. vi. 1988; 1 ex., 28. x. 1988.

#### Lygaeidae

*Nysius plebejus* Distant

Specimens examined. Shimota: 1 ex., 15. iv. 1988; 1 ex., 8. ix. 1988; 2 exx., 22. ix. 1988; 2 exx., 5. x. 1988; 1 ex., 19. x. 1988. Toso: 1 ex., 11. vi. 1988.

#### Alydidae

*Riptortus clavatus* (Thunberg)

Specimens examined. Shimota: 3 exx., 8. ix. 1988; 1 ex., 22. ix. 1988.

#### Coreidae

*Stictopleurus punctatonevosus* Goeze

Specimens examined. Toso: 1 ex., 27. v. 1988. Shimota: 1 ex., 15. vi. 1988.

*Cletus trigonus* (Thunberg)

Specimen examined. Shimota: 1 ex., 1. vi. 1988.

#### Pentatomidae

*Eysarcoris ventralis* (Westwood)

Specimens examined. Toso: 1 ex., 11. vi. 1988; 1 ex., 9. ix. 1988; 1 ex., 21. x. 1988.

#### Thysanoptera

##### Thripidae

*Thrips hawaiiensis* (Morgan)

Specimens examined. Korimoto: 213 exx., 3. x. 1988.

*Megaleurothrips distalis* (Karny)

Specimens examined. Korimoto: 12 exx., 3. x. 1988.

#### Coleoptera

##### Scarabaeidae

*Anomala lucens* Ballion

Specimens examined. Korimoto: 2 exx., 13. vi. 1988; 2 exx., 20. vi. 1988; 3 exx., 27. vi. 1988; 2 exx., 11. vii. 1988; 1 ex., 25. vii. 1988; 1 ex., 8. viii. 1988; 2 exx., 15. vii. 1988; 2 exx., 22. viii. 1988; 1 ex., 29. viii. 1988; 1 ex., 5. ix. 1988. Shimota: 3 exx., 8. ix. 1988; 1 ex., 10. viii. 1988; 1 ex., 24. viii. 1988. Toso: 2 exx., 19. viii. 1988; 2 exx., 26. viii. 1988; 1 ex., 2. ix. 1988; 1 ex., 9. ix. 1988.

*Anomala cuprea* (Hope)

Specimens examined. Korimoto: 3 exx., 27. v. 1988. Shimota: 3 exx., 15. vi. 1988.

*Blitopertha orientalis* (Waterhouse)

Specimens examined. Korimoto: 2 exx., 4. vii. 1988. Shimota: 2 exx., 15. vi. 1988.

*Mimela splendens* (Gyllenhal)

Specimens examined. Shimota: 2 exx., 31. v. 1990.

*Popillia japonica* Newman

Specimens examined. Shimota: 3 exx., 8. ix. 1988.

#### Dermestidae

*Anthrenus verbasci* (Linnaeus)

Specimens examined. Emerged in the laboratory from seeds obtained from a seed dealer: 5 exx., 23. vii. 1989; 3 exx., 13. viii. 1989; 2 exx., 15. ix. 1989.

#### Nitidulidae

*Haptoncurina paulula* (Reiter)

Specimens examined. Shimota: 3 exx., 3. x. 1988.

#### Chrysomelidae

*Pagria signata* (Motschulsky)

Specimens examined. Korimoto: 5 exx., 20. v. 1988. Toso: 3 exx., 4. vi. 1988. Shimota: 5 exx., 24. vi. 1988.

#### Curculionidae

*Episomus turritus* (Gyllenhal)

Specimens examined. Toso: 1 ex., 9. v. 1988; 1 ex., 19. vii. 1988. Shimota: 3 exx., 31. v. 1988; 2 exx., 15. vi. 1988; 2 exx., 24. vi. 1988.

*Sympiezomias lewisi* (Roelofs)

Specimens examined. Toso: 3 exx., 4. vi. 1988; 1 ex., 24. vi. 1988.

*Heterarthrus lewisi* (Pascoe)

Specimen examined. Shimota: 1 ex., 31. v. 1990.

#### Lepidoptera

##### Tortricidae

*Matsumuraeses falcana* (Walsingham)

Specimens examined. Shimota: 1 ex., 3. v. 1988; 2 exx., 18. v. 1988.

##### Psychidae

*Acanthopsyche nigraplaga* (Wileman)

Specimens examined. Shimota: 1 ex., 5. x. 1988; 1 ex., 8. x. 1988; 1 ex., 19. x. 1988.

##### Pyralidae

*Hedylepta misera* (Butler)

Specimens examined. Korimoto: 1 ex., 8. v. 1988. Toso: 1 ex., 30. iv. 1988; 1 ex., 3. v. 1988.

*Maruca testulalis* (Hübner)

Specimens examined. Shimota: 4 exx., 8. ix. 1988; 3 exx., 12. ix. 1988; 4 exx., 22. ix. 1988; 2 exx., 30. ix. 1988; 2 exx., 3. x. 1988. Toso: 1 ex., 3. ix. 1988; 1 ex., 8. ix. 1988; 2 exx., 30. ix. 1988.

##### Noctuidae

*Paragabara flavomaculata* (Oberthür)

Specimens examined. Toso: 2 exx., 11. vi. 1988; 1 ex., 19. viii. 1988.

Table 1-a. List of phytophagous insects of Kudzu plant, *Pueraria lobata* (Willd.) Ohwi<sup>1)</sup>

| Species                                    | Host plant <sup>2)</sup>   | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>  | Reference <sup>5)</sup> |
|--|--|-----------------------------------|---|-------------------------|
| <b>I INSECTA</b>                           |  |                                   |   |                         |
| <b>ORTHOPTERA</b>                          |  |                                   |   |                         |
| Tettigoniidae                              |  |                                   |   |                         |
| * <i>Gampsocleis buergeri</i> (de Haan)    | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Gladiolus hybridus</i> Hort.   | Leaf                              | Japan (Honshu, Shikoku, Kyushu, Tsu-shima)  | 46<br>Present data      |
| Pyrgomorphidae                             |  |                                   |   |                         |
| * <i>Atractomorpha lata</i> (Motschulsky)  | <i>Glycine max</i> Merrill<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>Phaseolus vulgaris</i> L.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Dahlia jurezii</i> Hort.<br><i>Chrysanthemum morifolium</i> Ram. | Leaf                              | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Yaku-shima, Ryukyus, Izu-oshima), Taiwan, Korea, Oriental region | 14, 46<br>Present data  |
| Acrididae                                  |  |                                   |   |                         |
| * <i>Acrida cinerea</i> (Thunberg)         | ** <i>Pueraria lobata</i> (Willd.) Ohwi  | Leaf                              | Japan (Honshu, Shikoku, Kyushu, Tsu-shima, Ryukyus, Izu-oshima), Europe, Africa, Oriental region, Australian region   | Present data            |
| * <i>Oxya japonica japonica</i> (Thunberg) | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Oryza sativa</i> L.<br><i>Saccharum officinarum</i> L.<br>etc.   | Leaf                              | Japan (Honshu, Shikoku, Kyushu, Tanega-shima, Yaku-shima, Ryukyus), Taiwan, Oriental region                           | 46<br>Present data      |
| * <i>Patanga japonica</i> (Bolivar)        | <i>Glycine max</i> Merrill<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>Phaseolus vulgaris</i> L.   | Leaf                              | Japan (Honshu, Shikoku, Kyushu, Tsu-shima, Ryukyus), Taiwan, Korea, China, India, Himalaya,                           | 46<br>Present data      |

1) Published records and the present data.

2) Species names after Makino<sup>37)</sup>.

3) Plant part of Kudzu affected. That of some species is not available in the published records.

4) After Hirashima<sup>12)</sup>.

5) References which recorded Kudzu feeding and other host plant feeding. Numbers indicate those in the literature list of text.

\* Feeding on Kudzu in Kagoshima city.

\*\* New host plant recorded.

Table 1-a. (Continued)

| Species                                 | Host plant <sup>2)</sup>  | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>   | Reference <sup>5)</sup> |
|---|---|-----------------------------------|--|-------------------------|
|   | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Oryza sativa</i> L.<br><i>Triticum aestivum</i> L.<br><i>Setaria</i> spp.<br>etc.   |                                   | Sikkim   |                         |
| Tetrigidae                              |   |                                   |  |                         |
| * <i>Tetrix japonica</i> (Bolivar)      | <i>Brassica</i> spp.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br>Leguminous plants<br><i>Solanum tuberosum</i> L.<br><i>Zea mays</i> L.<br><i>Zoysia japonica</i> Steud.<br>etc.  | Leaf                              | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Iki, Tsu-shima, Ryukyus), Saghalien, China, Taiwan, Mongolia, USSR | 46<br>Present data      |
| HEMIPTERA-HOMOPTERA                     |   |                                   |  |                         |
| Flatidae                                |   |                                   |  |                         |
| * <i>Geisha distinctissima</i> (Walker) | <i>Castanea crenata</i> Sieb. et Zucc.<br><i>Morus bombycis</i> Koidz.<br><i>Paeonia suffruticosa</i> Andr.<br><i>Prunus mume</i> Sieb. et Zucc.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Camellia sinensis</i> (L.) O. Kuntze<br><i>Diospyros kaki</i> Thunb.<br>etc.    | Stem, petiole                     | Japan (Honshu, Shikoku, Kyushu, Tsu-shima, Yaku-shima, Ryukyus), Taiwan, China                               | 40, 46<br>Present data  |
| Ricaniidae                              |   |                                   |  |                         |
| * <i>Orosanga japonica</i> (Melichar)   | <i>Morus bombycis</i> Koidz.<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>Phaseolus vulgaris</i> L.<br><i>Glycine max</i> Merrill<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Camellia sinensis</i> (L.) O. Kuntze | Stem, petiole                     | Japan (Honshu, Shikoku, Kyushu, Ryukyus), Taiwan, Korea  | 40, 46<br>Present data  |

|              |  |                  |   |                        |  |
|--------------|--|------------------|---|------------------------|--|
|              | <i>Diospyros kaki</i> Thunb,<br>etc.   |                  |   |                        |  |
| Cercopidae   | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Chrysanthemum</i> spp.   | Stem,<br>petiole | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu, Tsu-shima, Yaku-shima),<br>Taiwan, Korea, USSR | 40<br>Present data     |  |
| Penthimiidae | <i>Castanea crenata</i> Sieb. et Zucc.<br><i>Pyrus pyrifolia</i> (Burm.) var. <i>culta</i> Nakai<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Diospyros kaki</i> Thunb.  | Leaf             | Japan (Honshu, Shikoku, Kyushu,<br>Ryukyus), Taiwan, Korea, China,<br>USSR                  | 46<br>Present data     |  |
| Cicadellidae | <i>Morus bombycis</i> Koidz.<br><i>Ficus carica</i> L.<br><i>Arachis hypogaea</i> L.<br><i>Glycine max</i> Merrill<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>Phaseolus vulgaris</i> L.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Vitis vinifera</i> L.<br><i>Camellia sinensis</i> (L.) O. Kuntze<br><i>Diospyros kaki</i> Thunb.<br>etc. | Leaf             | Japan (Honshu, Shikoku, Kyushu,<br>Tsu-shima)   | 46<br>Present data     |  |
|              | <i>Morus</i> spp.<br><i>Camellia sinensis</i> (L.) O. Kuntze<br><i>Pyrus</i> spp.<br><i>Rosa</i> spp.<br><i>Prunus</i> spp.<br><i>Glycine max</i> Merrill<br><i>Vigna angularis</i> (Willd.)<br><i>V. unguiculata</i> (L.)<br><i>Lablab purpureus</i> (L.)<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Diospyros kaki</i> Thunb.<br><i>Gladiolus gandavensis</i> van Houtt      | Leaf             | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu), Taiwan, Palearctic region                     | 40, 46<br>Present data |  |
|              | * <i>Cicadella viridis</i> (Linnaeus)  |                  |   |                        |  |

Table 1-a. (Continued)

| Species                        | Host plant <sup>2)</sup>  | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>                              | Reference <sup>5)</sup> |
|--------------------------------|---|-----------------------------------|---|-------------------------|
| Typhlocybiidae                 | <i>Oryza sativa</i> L.<br><i>Triticum aestivum</i> L.<br><i>Hordeum vulgare</i> L. var. <i>vulgare</i><br><i>Secale cereale</i> L.<br><i>Saccharum officinarum</i> L.<br><i>Zea mays</i> L.<br>etc.   |                                   |   |                         |
| * <i>Empoasca</i> sp.          | ** <i>Pueraria lobata</i> (Willd.) Ohwi   | Leaf                              |   | Present data            |
| Aphididae                      | <i>Astragalus sinicus</i> L.<br><i>Robinia pseudo-acacia</i> L.<br><i>Vicia faba</i> L.<br><i>V. hirsuta</i> (L.)<br><i>Pisum sativum</i> L.<br><i>Vigna angularis</i> (Wild.) Ohwi et Ohashi<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc. | Leaf,<br>stem                     | Japan (Hokkaido, Honshu, Shikoku, Kyushu), cosmopolitan | 11, 40, 46              |
| Ortheziidae                    | <i>Salix</i> spp.<br><i>Polygonum Blumei</i> Meisn.<br><i>Akebia quinata</i> Decne<br><i>Lespedeza</i> spp.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc.   | Leaf                              | Japan (Honshu, Shikoku, Kyushu), Korea, Taiwan          | 21                      |
| Eriococcidae                   | <i>Indigofera pseudo-inctoria</i> Matsum.<br><i>Desmodium podocarpum</i> DC. subsp. <i>fallax</i> Schindl.<br><i>Lespedeza buergeri</i> Miq.<br><i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc.                    | Stem                              | Japan (Honshu, Shikoku, Kyushu)                         | 21, 46                  |
| <i>Eriococcus sojae</i> Kuwana |   |                                   |   |                         |



|  |   |      |  |        |              |
|--|---|------|--|--------|--------------|
| Diaspidae  |   |      |  |        |              |
| * <i>Pseudaulacaspis pentagona</i><br>(Targioni) | <i>Salix</i> spp.<br><i>Juglans mandshurica</i> Maxim.<br>var. <i>sachalinensis</i> (Miyabe et Kudo)<br><i>Morus bombycis</i> Koidz.<br><i>Pyrus pyrifolia</i> (Burm.) var. <i>culta</i> Nakai<br><i>Prunus persica</i> Batsch<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Zanthoxylum piperitum</i> (L.) DC.<br><i>Camellia sinensis</i> (L.) O. Kuntze.<br><i>Diospyros kaki</i> Thunb.<br><i>Cycas revoluta</i> Thunb.<br>etc. | Stem | Japan (Honshu, Shikoku, Kyushu,<br>Nansei-shoto), cosmopolitan         | 21, 46 |              |
| <b>HEMIPTERA-HETEROPTERA</b>                     |   |      |  |        |              |
| Miridae  |   |      |  |        |              |
| * <i>Adelphocoris triannulatus</i><br>(Stål)     | <i>Indigofera pseudo-tinctoria</i> Matsum.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Cucumis melo</i> L.<br><i>Chrysanthemum</i> spp.<br><i>Oryza sativa</i> L.<br><i>Lolium multiflorum</i> Lam.<br><i>Allium tuberosum</i> Rottl.  | Leaf | Japan (Hokkaido, Shikoku, Kyushu),<br>Korea, China, Saghalien, Kuriles | 22     | Present data |
| * <i>Ectometopterus micanulus</i><br>(Horváth)   | <i>Indigofera pseudo-tinctoria</i> Matsum.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Cucumis melo</i> L.<br><i>Chrysanthemum</i> spp.<br><i>Oryza sativa</i> L.<br><i>Lolium multiflorum</i> Lam.<br><i>Allium tuberosum</i> Rottl.   | Leaf | Japan (Honshu, Shikoku, Kyushu)  | 22     |              |
| <i>Proboscidoecoris varicornis</i><br>(Jakovlev) | <i>Pueraria lobata</i> (Willd.) Ohwi  |      | Japan (Honshu, Kyushu, Tsu-shima),<br>Korea, Oriental region           | 22     |              |
| * <i>Halticillus insularis</i> (Usinger)         | <i>Raphanus sativus</i> L.<br><i>Arachis hypogaea</i> L.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Ipomoea batatas</i> Lam.<br><i>Cucumis melo</i> L.<br>etc.   | Leaf | Japan (Tsu-shima, Ryukyus),<br>Taiwan, Micronesia                      | 22, 38 |              |
| <i>Pilophorus typicus</i> (Distant)              | <i>Pueraria lobata</i> (Willd.) Ohwi  | Leaf | Japan (Honshu, Shikoku, Kyushu,  | 22     |              |

Table 1-a. (Continued)

| Species                                | Host plant <sup>2)</sup>  | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>  | Reference <sup>5)</sup> |
|--|---|-----------------------------------|---|-------------------------|
|  | <i>Oryza sativa</i> L.  |                                   | Ryukyus, Taiwan, China, Oriental region                                   |                         |
| Berytidae                              |   |                                   |   |                         |
| * <i>Metacanthus pulchellus</i> Dallas | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Paulownia tomentosa</i> Steudel<br><i>Sesamum indicum</i> L.  | Leaf                              | Japan (Hokkaido, Honshu, Shikoku, Kyushu), China, India                   | 22<br>Present data      |
| <i>Yemma exilis</i> Horváth            | <i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Paulownia tomentosa</i> Steudel<br><i>Sesamum indicum</i> L.<br><i>Setaria viridis</i> Beauv.<br>etc.  | Leaf, stem, ear                   | Japan (Honshu, Shikoku, Kyushu)   | 22                      |
| Lygaeidae                              |   |                                   |   |                         |
| * <i>Nysius plebejus</i> Distant       | <i>Rumex Acetosa</i> L.<br><i>Fragaria</i> spp.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Erigeron philadelphicus</i> L.<br><i>Chrysanthemum morifolium</i> Ram.<br><i>Oryza sativa</i> L.<br><i>Agropyron kamoji</i> Ohwi<br><i>Lolium multiflorum</i> Lam.<br><i>Eleusine indica</i> (L.) Gaertn.<br><i>Setaria viridis</i> Beauv.<br>etc. | Leaf                              | Japan (Honshu, Shikoku, Kyushu, Ryukyus), Midway Is.                      | 22<br>Present data      |
| * <i>Piocoris varius</i> (Uhler)       | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus junos</i> Sieb. ex Tanaka<br><i>Erigeron annuus</i> L.<br><i>Chrysanthemum</i> spp.<br>Aphids, scale insects etc.   |                                   | Japan (Honshu, Shikoku, Kyushu, Ryukyus), Korea                           | 22                      |
| * <i>Chauliops fallax</i> Scott        | <i>Desmodium</i> spp.<br><i>Glycine max</i> Merrill<br><i>Amphicarpaea edgeworthii</i> Benth.   | Leaf, flower                      | Japan (Honshu, Shikoku, Kyushu, Ryukyus), Taiwan, China, Sri Lanka, India | 22, 25, 50, 62          |

|   |  |                           |   |                        |  |  |
|---|--|---------------------------|---|------------------------|--|--|
| Alydidae                                      |  |                           |   |                        |  |  |
| * <i>Riptortus clavatus</i> (Thunberg)        | <i>Vigna angularis</i> (Wild.) Ohwi et Ohashi<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Oryza sativa</i> L.  | Stem,<br>flower,<br>pod   | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu, Nansei-shoto), Taiwan,<br>Korea        | 22, 25<br>Present data |  |  |
|   | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br>Leguminous plants<br><i>Citrus</i> spp.<br><i>Ipomoea batatas</i> Lam.<br><i>Sesamum indicum</i> L.<br>etc.   |                           |   |                        |  |  |
| Coreidae                                      |  |                           |   |                        |  |  |
| * <i>Cletus trigonus</i> (Thunberg)           | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Oryza sativa</i> L.<br><i>Setaria viridis</i> L.<br><i>Digitaria ciliaris</i> (Retz.)<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Glycine max</i> Merrill<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc.                             | Leaf,<br>stem,<br>petiole | Japan (Honshu, Shikoku, Kyushu,<br>Nansei-shoto), Taiwan, China,<br>Oriental region | 22<br>Present data     |  |  |
|   | <i>Homoeocerus dilatatus</i> Horváth   |                           | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu), Korea, China, E. Siberia              | 22                     |  |  |
| * <i>H. unipunctatus</i> (Thunberg)           | <i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Lepedeza cuneata</i> G. Don<br><i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Oryza sativa</i> L.<br><i>Morus bombycis</i> Koidz.<br><i>Rumex japonicus</i> Houtt.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Oryza sativa</i> L.<br>etc. | Leaf,<br>stem,<br>petiole | Japan (Honshu, Shikoku, Kyushu,<br>Ryukyus), Taiwan, China,<br>Oriental region      | 22                     |  |  |
|   | <i>Hygia (Hygia) opeca</i> (Uhler)   |                           | Japan (Honshu, Shikoku, Kyushu,<br>Ryukyus), Korea, Taiwan, China                   | 22                     |  |  |
| * <i>Sitctopleurus punctatonevus</i><br>Goeze | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Oryza sativa</i> L.  | Leaf                      | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu)  | 22<br>Present data     |  |  |
| Plataspidae                                   |  |                           |   |                        |  |  |
| <i>Coptosoma biguttulum</i> Motschulsky       | <i>Lepedeza</i> spp.<br><i>Glycine max</i> Merrill<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi   |                           | Japan (Hokkaido, Shikoku, Kyushu),<br>Korea, China, E. Siberia                      | 22                     |  |  |

Table 1-a. (Continued)

| Species   | Host plant <sup>2)</sup>   | Plant part affected <sup>3)</sup>   | Distribution <sup>4)</sup>   | Reference <sup>5)</sup> |
|---|--|-------------------------------------|--|-------------------------|
| * <i>Megacopta punctatissima</i><br>(Montandon) | <i>Pueraria lobata</i> (Willd.) Ohwi<br>etc.<br><i>Deutzia crenata</i> Sieb. et Zucc.<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Astragalus sinicus</i> L.<br><i>Lespedeza</i> spp.<br><i>Glycine max</i> Merrill<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Cirrus</i> spp.<br><i>Oryza sativa</i> L.<br><i>Triticum aestivum</i> L.<br>etc. | Stem,<br>petiole,<br>flower,<br>pod | Japan (Honshu, Shikoku, Kyushu,<br>Tsu-shima), China                             | 10, 22, 63              |
| Dinidridae                                      |  |                                     |  |                         |
| <i>Megynenum gracilicorne</i> Dallas            | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Cucumis melo</i> L.<br><i>C. sativus</i> L.<br><i>Trichosanthes cucumeroides</i><br>Maxim.<br>etc.  |                                     | Japan (Honshu, Shikoku, Kyushu,<br>Ryukyus), Korea, Taiwan, China                | 22                      |
| Pentatomidae                                    |  |                                     |  |                         |
| <i>Dolycoris baccalum</i> (L.)                  | <i>Oryza sativa</i> L.<br><i>Glycine max</i> Merrill<br><i>Phaseolus vulgaris</i> L.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc.   |                                     | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu), Korea, China, Palearctic<br>region | 22, 25                  |
| * <i>Eysarcoris ventralis</i> (Westwood)        | <i>Glycine max</i> Merrill<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Oryza sativa</i> L.<br><i>Agropyron kamoji</i> Ohwi<br><i>Lolium multiflorum</i> Lam.<br><i>Eleusine indica</i> Gaertn.<br><i>Setaria italica</i> Beauv.<br>etc.   | Leaf                                | Japan (Honshu, Kyushu, Nansei-<br>shoto), Oriental region                        | 22<br>Present data      |

|                                      |   |   |        |
|--------------------------------------|---|---|--------|
| <i>Halyomorpha picus</i> (Fabricius) | <i>Robinia pseudo-acacia</i> L.<br><i>Vigna angularis</i> (Wild.) Ohwi et Ohashi<br><i>Phaseolus vulgaris</i> L.<br><i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Vitis vinifera</i> L.<br><i>Triticum aestivum</i> L.<br>etc. | Japan (Honshu, Shikoku, Kyushu, Ryukyus), Taiwan, Korea, China, Oriental region                 | 22     |
| <i>Homalagonia obtusa</i> (Walker)   | <i>Malus</i> spp.<br><i>Prunus persica</i> Batsch<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Pisum sativum</i> L.<br><i>Glycine max</i> Merrill<br><i>Phaseolus vulgaris</i> L.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc.                 | Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, China, E. Siberia                             | 22     |
| <i>Menida violacea</i> Motschulsky   | <i>Morus bombycis</i> Koidz.<br><i>Oryza sativa</i> L.<br><i>Robus crataegifolius</i> Bunge<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc.  | Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, China, E. Siberia                             | 22     |
| <i>Nezara antennata</i> Scott        | <i>Vicia faba</i> L.<br><i>Glycine max</i> Merrill<br><i>Vigna angularis</i> (Wild.) Ohwi et Ohashi<br><i>Phaseolus vulgaris</i> L.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Triticum aestivum</i> L.<br>etc.                  | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Nansei-shoto), Taiwan, China, Korea, India, Indochina | 22, 25 |
| <i>Piezodorus hybneri</i> (Gmelin)   | <i>Medicago polymorpha</i> L.<br><i>Laspedeza</i> spp.<br><i>Vicia faba</i> L.<br><i>Glycine max</i> Merrill<br><i>Phaseolus vulgaris</i> L.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Veronica</i> spp.<br><i>Oryza sativa</i> L.<br>etc.            | Japan (Honshu, Shikoku, Kyushu, Nansei-shoto), Taiwan, China, India, Australia                  | 22, 25 |

Table 1-a. (Continued)

| Species                                  | Host plant <sup>2)</sup>  | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>   | Reference <sup>5)</sup> |
|--|---|-----------------------------------|--|-------------------------|
| <b>THYSANOPTERA</b>                      |   |                                   |  |                         |
| <b>Thripidae</b>                         |   |                                   |  |                         |
| <i>Salpingothrips aimotofus</i> Kudo     | <i>Pueraria lobata</i> (Willd.) Ohwi  | Flower                            | Japan (Honshu), Taiwan, Kainan-to  | 28                      |
| <i>Thrips coloratus</i> Schmutz          | <i>Ficus carica</i> L.<br><i>Vicia faba</i> L.<br><i>Pisum sativum</i> L.<br><i>Vigna angularis</i> (Wild.) Ohwi et Ohashi<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Vitis vinifera</i> L.<br><i>Camellia sinensis</i> (L.) O. Kuntze<br><i>Diospyros kaki</i> Thunb.<br><i>Nicotiana tabacum</i> L.<br>etc.  |                                   | Japan (Honshu, Shikoku, Kyushu),<br>China, S. E. Asia, India, Sri<br>Lanka, New Guinea   | 64                      |
| * <i>Thrips hawaiiensis</i> (Morgan)     | <i>Camellia sinensis</i> (L.) O. Kuntze<br><i>Brassica campestris</i> L.<br><i>Glycine max</i> Merrill<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp<br><i>Cucumis sativa</i> L.<br><i>Lycopersicon esculentum</i> Mill.<br><i>Nicotiana tabacum</i> L.<br><i>Allium fistulosum</i> L.<br><i>Oryza sativa</i> L.<br><i>Saccharum officinarum</i> L.<br>etc. | Flower                            | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu, Ryukyus), Taiwan, China,<br>S. E. Asia, India, New Guinea,<br>Micronesia, Hawaii, Europe. | 41, 64<br>Present data  |
| * <i>Megalurothrips distalis</i> (Karny) | <i>Vicia faba</i> L.<br><i>Glycine max</i> Merrill<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Luffa aegyptiaca</i> Mill.<br><i>Diospyros kaki</i> Thunb.<br><i>Ipomaea batatas</i> Lam.   | Flower                            | Japan (Honshu, Shikoku, Kyushu,<br>Ryukyus), Taiwan, China, S. E.<br>Asia, India, Sri Lanka.   | 41, 64<br>Present data  |

*Chrysanthemum morifolium* Ramat.  
*Allium cepa* L.  
 etc.

#### COLEOPTERA

##### Scarabaeidae

##### \* *Anomala cuprea* (Hope)

|   |      |   |              |
|---|------|---|--------------|
| <i>Quercus</i> sp.                      | Leaf | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Tanegashima, Yaku-shima) | 46           |
| <i>Fragaria x-magna</i> Thuill          |      |   | Present data |
| <i>Rosa</i> spp.                        |      |   |              |
| <i>Prunus</i> spp.                      |      |   |              |
| <i>Albizia julibrissin</i> Durazz.      |      |   |              |
| <i>Glycine max</i> Merrill              |      |   |              |
| <i>Vigna angularis</i> (Willd.)         |      |   |              |
| <i>V. unguiculata</i> (L.)              |      |   |              |
| ** <i>Pueraria lobata</i> (Willd.) Ohwi |      |   |              |
| <i>Citrus</i> spp.                      |      |   |              |
| <i>Diospyros kaki</i> Thunb.            |      |   |              |
| <i>Ipomoea batatas</i> Lam.             |      |   |              |
| <i>Arcium lappa</i> L.                  |      |   |              |
| <i>Iris</i> spp.                        |      |   |              |
| etc.                                    |      |   |              |

##### \* *A. lucens* Ballion

|  |              |   |              |
|--|--------------|---|--------------|
| <i>Glycine max</i> Merrill                     | Leaf, flower | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Yaku-shima) | 46           |
| <i>Vigna angularis</i> (Willd.) Ohwi et Ohashi |              |   | Present data |
| <i>Phaseolus vulgaris</i> L.                   |              |   |              |
| ** <i>Pueraria lobata</i> (Willd.) Ohwi        |              |   |              |
| <i>Zea mays</i> L.                             |              |   |              |
| <i>Picea jezoensis</i> Carr.                   |              |   |              |
| <i>Abies sachalinensis</i> (Fr. Schm.)         |              |   |              |
| <i>Pinus thunbergii</i> Parl.                  |              |   |              |

##### \* *Blitopertha orientalis* (Waterhouse)

|   |      |   |              |
|---|------|---|--------------|
| <i>Rosa</i> sp.                         | Leaf | Japan (Hokkaido, Honshu, Shikoku, Kyushu) | 46           |
| <i>Prunus</i> spp.                      |      |   | Present data |
| ** <i>Pueraria lobata</i> (Willd.) Ohwi |      |   |              |
| <i>Citrus</i> spp.                      |      |   |              |
| <i>Diospyros kaki</i> Thunb.            |      |   |              |
| <i>Zoysia</i> spp.                      |      |   |              |

##### \* *Mimela splendens* (Gyllenhal)

|  |      |   |              |
|--|------|---|--------------|
| <i>Rosa</i> spp.                         | Leaf | Japan (Hokkaido, Honshu, Shikoku, Kyushu) | 46           |
| ** <i>Pueraria lobata</i> (Willd.) Ohwi  |      |   | Present data |
| <i>Camellia sinensis</i> (L.) O. Kuntze. |      |   |              |
| <i>Diospyros kaki</i> Thunb.             |      |   |              |

Table 1-a. (Continued)

| Species                                 | Host plant <sup>2)</sup>   | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>   | Reference <sup>5)</sup> |
|---|--|-----------------------------------|--|-------------------------|
| * <i>Popillia japonica</i> Newman       | <i>Castanea crenata</i> Sieb. et Zucc.<br><i>Malus</i> spp.<br><i>Rosa</i> spp.<br><i>Prunus</i> spp.<br><i>Glycine max</i> Merrill<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Vitis vinifera</i> L.<br><i>Diospyros kaki</i> Thunb.<br><i>Dahlia juorezii</i> Hort.<br><i>Asparagus officinalis</i> L.<br>var. <i>atilis</i> L.<br>etc. | Leaf                              | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Yaku-shima)   | 46<br>Present data      |
| Buprestidae                             |  |                                   |  |                         |
| * <i>Trachys auricollis</i> E. Saunders | <i>Pueraria lobata</i> (Willd.) Ohwi   | Leaf                              | Japan (Honshu, Shikoku, Kyushu, Sado, Tanega-shima, Yaku-shima)  | 34, 35                  |
| <i>T. reitteri</i> Obenberger           | <i>Glycine max</i> Merrill<br><i>Amphicarpaea edgeworthii</i> Benth.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc.   | Leaf                              | Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, China  | 34, 35                  |
| Dermeestidae                            |  |                                   |  |                         |
| * <i>Anthrenus verbasci</i> (Linnaeus)  | Dried specimen, woolen cloth,<br>etc.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi   | Stored seed                       | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Ryukyus), cosmopolitan   | 45, 46<br>Present data  |
| Nitidulidae                             |  |                                   |  |                         |
| * <i>Haptoncurina paulula</i> (Reitter) | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Poncirus trifoliata</i> (L.)<br><i>Fortunella</i> spp.   | Flower                            | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Sado, Izu-shoto, Tsu-shima, Goto-retto, Tanega-shima, Yaku-shima, Ryukyus), Taiwan, China, Bhutan, USSR. | 46<br>Present data      |
| Cerambycidae                            |  |                                   |  |                         |
| <i>Falsomesosella gracilior</i> (Bates) | <i>Pterocarya rhoifolia</i> Sieb. et Zucc.<br><i>Alnus hirsuta</i> (Turcz.)<br><i>Zelkova serrata</i> Makino   | Stem                              | Japan (Honshu, Shikoku, Kyushu, Tsu-shima)   | 26                      |



|   |      |  |    |  |
|---|------|--|----|--|
| <i>Morus bombycis</i> Koidz.                |      |  |    |  |
| <i>Prunus</i> spp.                          |      |  |    |  |
| <i>Wistaria brachybotrys</i> Sieb. et Zucc. |      |  |    |  |
| <i>Robinia pseudo-acacia</i> L.             |      |  |    |  |
| <i>Pueraria lobata</i> (Willd.) Ohwi        |      |  |    |  |
| <i>Abies firma</i> Sieb. et Zucc.           |      |  |    |  |
| etc.  |      |  |    |  |
| <i>Castanea crenata</i> Sieb. et Zucc.      | Stem | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Iki, Tanega-shima, Yakushima, Ryukyus) | 26 |  |
| <i>Quercus</i> spp.                         |      |  |    |  |
| <i>Zelkova serrata</i> Makino               |      |  |    |  |
| <i>Morus bombycis</i> Koidz.                |      |  |    |  |
| <i>Prunus</i> spp.                          |      |  |    |  |
| <i>Wistaria brachybotrys</i> Sieb. et Zucc. |      |  |    |  |
| <i>Pueraria lobata</i> (Willd.) Ohwi        |      |  |    |  |
| <i>Citrus</i> spp.                          |      |  |    |  |
| <i>Pinus densiflora</i> Sieb. et Zucc.      |      |  |    |  |
| etc.  |      |  |    |  |
| <i>Castanea crenata</i> Sieb. et Zucc.      | Stem | Japan (Tsu-shima), Korea, Saghalien  | 26 |  |
| <i>Quercus</i> spp.                         |      |  |    |  |
| <i>Zelkova serrata</i> Makino               |      |  |    |  |
| <i>Morus bombycis</i> Koidz.                |      |  |    |  |
| <i>Malus</i> spp.                           |      |  |    |  |
| <i>Wistaria brachybotrys</i> Sieb. et Zucc. |      |  |    |  |
| <i>Pueraria lobata</i> (Willd.) Ohwi        |      |  |    |  |
| <i>Tsuga sieboldii</i> Carr.                |      |  |    |  |
| etc.  |      |  |    |  |
| <i>Quercus</i> spp.                         | Stem | Japan (Okushiri, Honshu, Shikoku, Kyushu, Tsu-shima), Palawan, Vietnam           | 26 |  |
| <i>Zelkova serrata</i> Makino               |      |  |    |  |
| <i>Wistaria brachybotrys</i> Sieb. et Zucc. |      |  |    |  |
| <i>Pueraria lobata</i> (Willd.) Ohwi        |      |  |    |  |
| <i>Acer rufinerve</i> Sieb. et Zucc.        |      |  |    |  |
| <i>Trichosanthes cucumeroides</i> Maxim.    |      |  |    |  |
| <i>Phyllostachys</i> spp.                   |      |  |    |  |
| <i>Pleiblastus</i> spp.                     |      |  |    |  |
| etc.  |      |  |    |  |
| <i>Juglans mandshurica</i> Maxim.           | Stem | Japan (Honshu, Shikoku, Kyushu, Tsu-shima, Iki)                                  | 26 |  |
| var. <i>sieboldiana</i> Makino              |      |  |    |  |
| <i>Alnus hirsuta</i> (Turcz.)               |      |  |    |  |
| <i>Castanea crenata</i> Sieb. et Zucc.      |      |  |    |  |
| <i>Mesosa longipennis</i> Bates             |      |  |    |  |
| <i>M. myops myops</i> (Dalman)              |      |  |    |  |
| * <i>Atimura japonica</i> Bates             |      |  |    |  |
| <i>Mesosella simiola</i> Bates              |      |  |    |  |

Table 1-a. (Continued)

| Species                                 | Host plant <sup>2)</sup>  | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>   | Reference <sup>5)</sup> |
|---|---|-----------------------------------|--|-------------------------|
|   | <i>Zelkova serrata</i> Makino<br><i>Morus bombycis</i> Koidz.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Melia azedarach</i> L. var. <i>subiripinnata</i> Mig.<br><i>Vitis coignetiae</i> Pulliat<br><i>Abies firma</i> Sieb. et Zucc.<br>etc.   |                                   |  |                         |
| <i>Pterolophia annulata</i> (Chevrolat) | <i>Castanea crenata</i> Sieb. et Zucc.<br><i>Quercus</i> spp.<br><i>Zelkova serrata</i> Makino<br><i>Morus bombycis</i> Koidz.<br><i>Ficus carica</i> L.<br><i>Prunus yedoensis</i> spp.<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Vitis vinifera</i> L.<br>etc. | Stem                              | Japan (Honshu, Shikoku, Kyushu, Iki, Tsu-shima, Ogasawara, Ryukyus), Taiwan, China, Korea, Burma | 26                      |
| <i>P. caudata caudata</i> Bates         | <i>Castanea crenata</i> Sieb. et Zucc.<br><i>Quercus</i> spp.<br><i>Zelkova serrata</i> Makino<br><i>Morus bombycis</i> Koidz.<br><i>Ficus carica</i> L.<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Aesculus turbinata</i> Blume<br>etc.    | Stem                              | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Iki, Tanega-shima)                          | 26                      |
| <i>P. zonata</i> (Bates)                | <i>Castanea crenata</i> Sieb. et Zucc.<br><i>Quercus</i> spp.<br><i>Morus bombycis</i> Koidz.<br><i>Ficus carica</i> L.<br><i>Hydrangea macrophylla</i> (Thunb.)<br><i>Prunus persica</i> Batsh   | Stem                              | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tanega-shima, Yaku-shima)                              | 26                      |

|               |  |  |               |   |                        |
|---------------|--|--|---------------|---|------------------------|
|               | <i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Cryptomeria japonica</i> (L.) D. Don<br>etc. |  |               |   |                        |
| Bruchidae     |  |  |               |   |                        |
|               | <i>Callosobruchus ademptus</i> (Sharp)   | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Torilis japonica</i> (Houtt.) DC.   | Seed          | Japan (Honshu, Kyushu)  | 43                     |
| Chrysomelidae |  |  |               |   |                        |
|               | * <i>Pagria signata</i> (Motschulsky)  | <i>Glycine max</i> Merrill<br>** <i>Pueraria lobata</i> (Willd.) Ohwi  | Leaf          | Japan (Honshu, Shikoku, Kyushu, Ryukyus, Ogasawara), Korea, China, USSR, Taiwan, Indochina, Philippines, Guam, Indonesia, India | 23, 46<br>Present data |
| Curculionidae |  |  |               |   |                        |
|               | * <i>Episomus turritus</i> (Gyllenhal)   | <i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Lespedeza</i> spp.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br>Leguminous plants | Leaf, stem    | Japan (Honshu, Shikoku, Kyushu, Tsu-shima), Korea, China  | 43, 46<br>Present data |
|               | * <i>Sympiezomias lewisi</i> (Roelofs)   | <i>Morus bombycis</i> Koidz.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br>etc.                                | Leaf          | Japan (Honshu, Shikoku, Kyushu)   | 43<br>Present data     |
|               | * <i>Eugnathus distinctus</i> Roelofs  | <i>Lespedeza</i> spp.<br><i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>Leguminous plants                     | Leaf, stem    | Japan (Honshu, Shikoku, Kyushu, Ryukyus), China, Taiwan, Korea  | 43, 46                 |
|               | <i>Mesalcidodes trifidus</i> (Pascoe)  | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Pyrus</i> spp.<br><i>Citrus</i> spp.  | Stem, petiole | Japan (Honshu, Shikoku, Kyushu, Tsu-shima), Taiwan, China   | 42, 43, 46             |
|               | * <i>Heterarthrus lewisi</i> Wollaston   | <i>Wistaria brachybotrys</i> Sieb. et Zucc.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi   | Withered stem | Japan (Honshu, Shikoku, Kyushu, Tsu-shima)  | 43<br>Present data     |
| DIPTERA       |  |  |               |   |                        |
| Cecidomyiidae |  |  |               |   |                        |
|               | <i>Lasioptera puerariae</i> Shinji<br>* <i>Pitydiplosis</i> sp.  | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Pueraria lobata</i> (Willd.) Ohwi   | Stem<br>Leaf  | Japan (Honshu), Korea<br>Japan (Honshu, Kyushu, Nansei-shoto, Ryukyus)  | 70<br>69, 71, 72       |

Table 1-a. (Continued)

| Species                                       | Host plant <sup>2)</sup>   | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>  | Reference <sup>5)</sup> |
|---|--|-----------------------------------|---|-------------------------|
| * Genus sp.                                   | <i>Pueraria lobata</i> (Willd.) Ohwi   | Leaf                              | Japan (Honshu, Kyushu, Nansei-shoto, Ryukyus)   | 69, 71, 72              |
| Agromyzidae                                   |  |                                   |   |                         |
| * <i>Japanagromyza tristella</i> (Thomson)    | <i>Glycine max</i> Merrill<br><i>G. soja</i> Sieb. et Zucc.<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>Phaseolus vulgaris</i> L.<br><i>Vigna unguiculata</i> (L.) Walp.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc. | Leaf                              | Japan (Honshu), Taiwan, China   | 46, 48                  |
| <i>Japanagromyza variihalterata</i> (Malloch) | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Glycine soja</i> Sieb. et Zucc.   | Leaf                              | Japan (Honshu)  | 52                      |
| <i>Ophiomyia puerarivora</i> Sasakawa         | <i>Pueraria lobata</i> (Willd.) Ohwi   | Leaf                              | Japan (Honshu)  | 53                      |
| LEPIDOPTERA                                   |  |                                   |   |                         |
| Tortricidae                                   |  |                                   |   |                         |
| * <i>Matsumuraeses falcana</i> (Walsingham)   | <i>Robinia pseudo-acacia</i> L.<br><i>Glycine max</i> Merrill<br>** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Zoysia japonica</i> Steud.  | Leaf, young stem                  | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Izu-oshima, Yaku-shima, Okinawa), Taiwan, China, Nepal | 20, 51<br>Present data  |
| <i>M. ussuriensis</i> (Caradja)               | <i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi  | Leaf, young stem                  | Japan (Honshu, Shikoku), USSR   | 20, 39                  |
| <i>M. vicina</i> Kuznetsov                    | <i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi   | Leaf, young stem                  | Japan (Honshu, Tsu-shima, Hachijo-jima), China  | 20, 39, 47              |
| Psychidae                                     |  |                                   |   |                         |
| * <i>Acanthopsyche nigraplaga</i> (Wileman)   | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Diospyros kaki</i> Thunb.<br><i>Citrus</i> spp.  | Leaf                              | Japan (Honshu, Shikoku, Kyushu), China, India   | 13, 46<br>Present data  |
| Lyonetiidae                                   |  |                                   |   |                         |
| <i>Leucoptera puerariella</i> Kuroko          | <i>Pueraria lobata</i> (Willd.) Ohwi   | Leaf                              | Japan (Honshu, Kyushu)  | 32, 33                  |
| <i>Microthauma glycinella</i> (Hübner)        | <i>Glycine max</i> Merrill   | Leaf                              | Japan (Kyushu, Okinawa)   | 32, 33, 39              |

|  |   |                 |  |                    |  |
|--|---|-----------------|--|--------------------|--|
|  | <i>Pueraria lobata</i> (Willd.) Ohwi  |                 |  |                    |  |
| Gracillariidae                                 |   |                 |  |                    |  |
| * <i>Liocrobyla lobata</i> Kuroko              | <i>Pueraria lobata</i> (Willd.) Ohwi  | Leaf            | Japan (Honshu, Shikoku, Kyushu)  | 31, 33, 39         |  |
| * <i>Neolithocolletis hikomonticola</i> Kumata | <i>Pueraria lobata</i> (Willd.) Ohwi  | Leaf            | Japan (Honshu, Kyushu)   | 29, 33, 39         |  |
| * <i>Spulerina dissotoma</i> (Meyrick)         | <i>Lespedeza cyrtobotrya</i> Miq.<br><i>Pueraria lobata</i> (Willd.) Ohwi   | Leaf            | Japan (Honshu, Kyushu), India  | 30, 33             |  |
| <i>Hylaconis puerariae</i> Kumata              | <i>Amphicarpaea edgeworthii</i> Benth.<br><i>Pueraria lobata</i> (Willd.) Ohwi  | Leaf            | Japan (Hokkaido, Honshu)   | 29, 33, 39         |  |
| <i>Porphyosela dorinda</i> Meyrick             | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Desmodium heterocarpon</i> (L.) DC.  | Leaf            | Japan (Yaku-shima), Taiwan, India  | 29                 |  |
| Pyralidae                                      |   |                 |  |                    |  |
| <i>Charema noctescens</i> Moore                | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Tripterygium regelii</i> Sprague et Takeda   | Leaf            | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Tanega-shima, Yaku-shima), Korea, Taiwan, China, India-Malaysia                                     | 13, 15, 39         |  |
| * <i>Hedylecta misera</i> (Butler)             | ** <i>Pueraria lobata</i> (Willd.) Ohwi   | Leaf            | Japan (Honshu, Shikoku, Kyushu, Tanega-shima, Yaku-shima), Korea, China  | Present data       |  |
| * <i>Maruca testulalis</i> (Hübner)            | <i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>V. unguiculata</i> (L.) Walp.<br>** <i>Pueraria lobata</i> (Willd.) Ohwi   | Flower, petiole | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Yaku-shima, Ryukyus), Taiwan, China, Korea, India, Malaysia, Hawaii, Oceanic Is., Australia, Africa | 13<br>Present data |  |
| * <i>Pleuroptya punctimarginalis</i> (Hampson) | <i>Pueraria lobata</i> (Willd.) Ohwi  | Leaf            | Japan (Honshu, Kyushu, Okinawa), India, Malaysia   | 39                 |  |
| Lycaenidae                                     |   |                 |  |                    |  |
| * <i>Lampides boeticus</i> (Linnaeus)          | <i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Vicia faba</i> L.<br><i>Pisum sativum</i> L.<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>Phaseolus vulgaris</i> L.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>P. montana</i> Merr.<br>etc. | Flower, pod     | Japan (Honshu, Shikoku, Kyushu, Ryukyus), Europe, Africa, Asia, New Guinea, Australia, Hawaii  | 7, 46, 55          |  |
| <i>Celastrina argiolus ladonides</i>           | <i>Sophora flavescens</i> Ait.  | Leaf,           | Japan (Hokkaido, Honshu, Shikoku,  | 9                  |  |

Table 1-a. (Continued)

| Species  | Host plant <sup>2)</sup>  | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>   | Reference <sup>5)</sup> |
|--|---|-----------------------------------|--|-------------------------|
| (de l'Orza)                                    | <i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Desmodium podocarpum</i> DC.<br>subsp. <i>oxyphyllum</i> (DC.)<br>Ohashi var. <i>oxyphyllum</i> Maxim.<br><i>Lespedeza thunbergii</i> (DC.) Nakai<br><i>L. bicolor</i> Turcz.<br><i>Vicia cracca</i> L.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Cornus controversa</i> Hemsl.<br>etc. | flower                            | Kyushu, Tsu-shima, Tanega-shima,<br>Yaku-shima, Amami-oshima),<br>Korea, China, Ussuri   |                         |
| <i>Everes argiades helloitia</i> (Ménétrières) | <i>Medicago lupulina</i> L.<br><i>Sophora flavescens</i> Ait. f. <i>purpurascens</i><br>(Makino)<br><i>Indigofera pseudotinctoria</i> Matsum.<br><i>Vicia cracca</i> L.<br><i>V. sativa</i> L.<br><i>Astragalus sinicus</i> L.<br><i>Lotus corniculatus</i> L. var. <i>japonica</i> Regel<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>etc.     | Flower                            | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu, Tsu-shima, Tanega-<br>shima, Yaku-shima, Amami-<br>oshima), Amur, Ussuri, Korea,<br>China | 8                       |
| <i>Rapala arata</i> (Bremer)                   | <i>Deutzea crenata</i> Sieb. et Zucc.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Wistaria floribunda</i> (Willd.) DC.<br><i>Vicia cracca</i> L.<br><i>Lespedeza thunbergii</i> Nakai<br><i>Sophora flavescens</i> Ait. f. <i>purpurascens</i><br>(Makino)  | Seed                              | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu, Tsu-shima), Amur, China,<br>Korea   | 8                       |
| Curetidae                                      | <i>Caesalpinia decapetala</i> (Roth) Ait.<br>var. <i>japonica</i> Ohashi<br><i>Sophora flavescens</i> Ait. f. <i>purpurascens</i><br>(Makino)<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Robunia pseudo-acacia</i> L.   | Leaf,<br>stem,<br>pod,<br>bud     | Japan (Honshu, Shikoku, Kyushu,<br>Tsu-shima, Tanega-shima,<br>Amami-oshima)   | 7, 9, 55                |

## Curetidae

\* *Curetis acuta paracuta* de Niceville

|  |   |      |  |                |
|--|---|------|--|----------------|
| Nymphalidae                                    |   |      |  |                |
| <i>Neptis sappho intermedia</i><br>W. B. Pryer | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>P. montana</i> Merr.   | Leaf | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Yaku-shima), Amur, Ussuri, China, Korea | 7, 55          |
|  | <i>Zelkova serrata</i> Makino<br><i>Celtis sinensis</i> Pers.<br>var. <i>japonica</i> (Planch.)<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Robinia pseudo-acacia</i> L.<br><i>Lespedeza buergeri</i> Miq.<br><i>Vicia unijuga</i> Al. Br.<br><i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Firmiana simplex</i> (L.)<br>etc.   |      |  |                |
| Geometridae                                    |   |      |  |                |
| <i>Ascotis selenaria cretacea</i> (Bultner)    | <i>Castanea crenata</i> Sieb. et Zucc.<br><i>Malus</i> spp.<br><i>Pyrus pyrifolia</i> (Brum.) var. <i>culta</i> Nakai<br><i>Prunus persica</i> Batsch<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Vigna angularis</i> (Willd.) Ohwi et Ohashi<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Camellia sinensis</i> (L.)<br><i>Aster ageratoides</i> Turcz.<br>subsp. <i>ovatus</i> (Franch. et Savat.)<br><i>Chrysanthemum</i> spp.<br>etc. |      | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Tanega-shima, Yaku-shima)               | 39, 46, 47     |
| <i>Nyssiodes lefuarius</i> (Erschoff)          | <i>Rumex acetosa</i> L.<br><i>R. japonicus</i> Houtt.<br><i>Rosa multiflora</i> Thunb.<br><i>Chrysanthemum morifolium</i> Ram.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br>Leguminous plants<br>etc.   |      | Japan (Hokkaido, Honshu, Kyushu), USSR, China  | 13, 39         |
| Sphingidae                                     |   |      |  |                |
| <i>Clanis bilineata tsingtauica</i> Mell       | <i>Wistaria floribunda</i> (Willd.) DC.   | Leaf | Japan (Honshu, Shikoku, Kyushu)  | 13, 16, 39, 46 |


Table 1-a. (Continued)

| Species   | Host plant <sup>2)</sup>  | Plant part affected <sup>3)</sup> | Distribution <sup>4)</sup>   | Reference <sup>5)</sup> |
|---|---|-----------------------------------|--|-------------------------|
|   | <i>Robinia pseudo-acacia</i> L.<br><i>Lespedeza</i> spp.<br><i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi  |                                   | Tsu-shima, Yaku-shima, Okinawa),<br>Korea, China   |                         |
| Lymantriidae  |   |                                   |  |                         |
| * <i>Lymantria dispar japonica</i><br>(Motschulsky) | <i>Alnus hirstia</i> (Turcz.)<br><i>Castanea crenata</i> Sieb. et Zucc.<br><i>Malus</i> spp.<br><i>Rosa</i> spp.<br><i>Prunus</i> spp.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Diospyros kaki</i> Thunb.<br>etc.  | Leaf                              | Japan (Honshu, Shikoku, Kyushu)  | 13, 39, 46              |
| Arctiidae   |   |                                   |  |                         |
| <i>Thanatarctia imparilis</i> (Butler)              | <i>Salix</i> spp.<br><i>Castanea crenata</i> Sieb. et Zucc.<br><i>Quercus serrata</i> Thunb.<br><i>Pyrus pyrifolia</i> (Burm.) var. <i>culta</i> Nakai<br><i>Prunus persica</i> Batsch<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.                    | Leaf                              | Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsu-shima, Yaku-shima),<br>Taiwan, China                                   | 39, 46                  |
| Noctuidae   |   |                                   |  |                         |
| <i>Heliothis maritima adaucta</i> Butler            | <i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Linum usitatissimum</i> L.<br>Leguminous plants  | Leaf                              | Japan (Hokkaido, Honshu, Shikoku, Kyushu)  | 39, 46                  |
| * <i>Mocis undata</i> (Fabricius)                   | <i>Cytisus scoparius</i> (L.) Link<br><i>Wistaria brachybotrys</i> Sieb. et Zucc.<br><i>Desmodium podocarpum</i> DC.<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Medicago deniculata</i> Willd.<br><i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi | Leaf                              | Japan (Honshu, Kyushu, Izu-oshima, Ryukyus), Taiwan,<br>India, Australia,<br>Oceanic Is.<br>USA (Florida, Louisiana) | 39, 46, 57<br>1, 3      |
| <i>Anticarsia gemmatilis</i> Hübner                 |   | Leaf                              |  |                         |
| <i>Egryia apicalis</i> Guenée                       | <i>Wistaria brachybotrys</i> Sieb. et Zucc.   | Leaf                              | Japan (Honshu, Shikoku, Kyushu,  | 39, 57                  |



|   |   |      |   |                    |
|---|---|------|---|--------------------|
| * <i>Microxyla confusa</i> (Wileman)            | <i>Pueraria lobata</i> (Willd.) Ohwi<br>Leguminous plants   | Leaf | Tsu-shima, Yaku-shima,<br>Ryukyus), India, S.E. Asia      | 39, 57             |
| * <i>Paragabara flavomaculata</i><br>(Oberthür) | ** <i>Pueraria lobata</i> (Willd.) Ohwi<br>Leguminous plants  | Leaf | Japan (Honshu, Shikoku, Kyushu,<br>Tsu-shima, Yaku-shima) | 68<br>Present data |
| II ARACHNOIDEA                                  |   |      |   |                    |
| ACARINA   |   |      |   |                    |
| Tetranychidae                                   |   |      |   |                    |
| * <i>Panonychus citri</i> (McGregor)            | <i>Pyrus pyrifolia</i> (Brum.) var. <i>culta</i> Nakai<br><i>Eriobotrya japonica</i> Lindl.<br><i>Prunus persica</i> Batsch<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Poncirus trifoliata</i> (L.)<br><i>Citrus</i> spp.<br><i>Fortunella</i> spp.<br><i>Diospyros kaki</i> Thunb.<br>Leguminous plants   | Leaf | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu)              | 2, 4               |
| * <i>Tetranychus kanzawai</i> Kishida           | <i>Malus</i> spp.<br><i>Pyrus serotina</i> Rehder var. <i>culta</i> Nakai<br><i>Prunus persica</i> Batsch<br><i>Fragaria</i> spp.<br><i>Fortunella</i> spp.<br><i>Glycine max</i> Merrill<br><i>Pueraria lobata</i> (Willd.) Ohwi<br><i>Citrus</i> spp.<br><i>Vitis vinifera</i> L.<br><i>Camellia sinensis</i> (L.) D. Kuntze<br><i>Solanum melongena</i> L.<br><i>Cucumis melo</i> L.<br><i>Zea mays</i> L.<br><i>Oryza sativa</i> L.<br>etc. | Leaf | Japan (Hokkaido, Honshu, Shikoku,<br>Kyushu, Okinawa)     | 4, 59              |











### Acknowledgements

The authors would like to express their sincere appreciation to Prof. A. Nagatomi and Prof. J. Yukawa (Kagoshima University) and to Prof. N. Ishibashi and Prof. S. Tojo (Saga University) for their kind guidance and criticism of this study.

The first author would like to thank Prof. K. Yano (Yamaguchi University) for his interest and encouragement throughout the study. Her hearty thanks are also expressed to the following entomologists who identified the species studied: Dr. T. Kumata (Hokkaido University) for Gracillariidae, Dr. M. Miyazaki (National Institute of Sericultural and Entomological Science) for Thysanoptera, Dr. Y. Yoshiyasu (Kyoto Prefectural University) for Pyralidae, Dr. F. Komai (Osaka University of Arts) for Tortricidae and Mr. H. Takizawa (Japan Tobacco Ind.) for Chrysomelidae.

Field accommodations and facilities at the Toso Orchid Station, Faculty of Agriculture, Kagoshima University, were provided through the courtesy of Mr. T. Kushima.

Further thanks are extended to the following colleagues of the Entomological Laboratory, Kagoshima University, for their field assistance and cooperation: Mr. S. Hayashikawa, Mr. S. Shobu and Mr. Y. Hirata.

Finally, the first author wishes to express her appreciation to the Ministry of Education, Science and Culture of Japan for providing her with a scholarship.

### References

- 1) Anazonwu, D. L. and Johnson, S. J.: Effects of host and density on larval color, size and development of the velvetbean caterpillar, *Anticarsia gemmatilis* (Lepidoptera: Noctuidae). *Env. Ent.* **15**, 779–783 (1986)
- 2) Ashihara, W.: Infestation and reproduction of the Citrus red mite, *Panonychus citri* (McGregor) (Acarina: Tetranychidae) on leguminous plants. *Appl. Ent. Zool.* **22**, 512–518 (1987)
- 3) Buschman, L. L., Whitcomb, W. H., Neal, T. M. and Mays, D. L.: Winter survival and hosts of the velvetbean caterpillar in Florida, USA. *Fla. Entomol.* **60**, 267–273 (1977)
- 4) Ehara, S. and Shinkaji, N.: An introduction to agricultural acarology. 328 pp. Zenkoku Noson Kyoiku Kyokai, Tokyo (1975) (in Japanese)
- 5) Forseth, I. N. and Teramura, A. H.: Field photosynthesis microclimate and water relations of an exotic temperate liana *Pueraria lobata* Kudzu. *Oecologia* **71**, 262–267 (1987)
- 6) Frankel, E.: Distribution of *Pueraria lobata* in and around New York City, New York, U.S.A. *Bull. Torrey Bot. Club* **116**, 390–394 (1989)
- 7) Fukuda, H., Kubo, K., Kuzuya, T., Takahashi, A., Takahashi, M., Tanaka, B. and Wakabayashi, M.: Insects' life in Japan, 3. Butterflies. xvi+278 pp., 52 pls. Hoikusha, Osaka (1972) (in Japanese)
- 8) Fukuda, H., Hama, E., Kuzuya, T., Takahashi, A., Tanaka, B., Tanaka, H., Wakabayashi, M. and Watanabe, Y.: The life histories of butterflies in Japan, II. xxii+325 pp., 64 pls., Hoikusha, Osaka (1983) (in Japanese)
- 9) Fukuda, H., Hama, E., Kuzuya, T., Takahashi, A., Tanaka, B., Tanaka, H., Wakabayashi, M. and Watanabe, Y.: The life histories of butterflies in Japan, III. xxii+373 pp., 72 pls., Hoikusha, Osaka (1984) (in Japanese)
- 10) Hibino, Y. and Ito, Y.: Mating aggregation of a stink bug, *Megacopta punctissimum* (Montandon) (Heteroptera: Plataspidae). *Res. Popul. Ecol.* **25**, 180–188 (1983)
- 11) Higuchi, H. and Miyazaki, M.: A tentative catalogue of host plants of Aphidoidea in Japan. *Ins. Mats. Suppl.* **5**: 1–66 (1969)
- 12) Hirashima, Y. (ed.): A check list of Japanese insects. Ent. Lab., Fac. Agr., Kyushu Univ., Fukuoka. 1767 pp. (1989)

- 13) Inoue, H.: Psychidae, Pyralidae, Geometridae, Sphingidae, Lymantridae and Arctiidae. in Inoue, H. et al., Moths of Japan, I: 158–162, 307–404, 412–579, 590–604 & 628–659. Kodansha, Tokyo (1982) (in Japanese)
- 14) Ishihara, T.: Orthoptera and Homoptera. in Asahina, S. et al. (eds.), Iconographia Insectorum Japonicorum Colore naturali edita, III: 45–54, pls. 23–25 & 107–136, pls. 55–68. Hokuryukan Pub. Co., Ltd., Tokyo (1965) (in Japanese)
- 15) Issiki, S.: Early stages of Japanese moths on colour, II. 237 pp., Hoikusha, Osaka (1969) (in Japanese)
- 16) Iwata, K.: Kontyu o mitsumete 50 nen, IV. 232 pp., Asahi Shinbunsha, Tokyo (1980) (in Japanese)
- 17) Julien, M. H. (ed.): Biological control of weeds. A world catalogue of agents and their target weeds. x+144 pp., CAB Int. (1987)
- 18) Kaneko, K.: Fundamental studies on the insects associated with *Pueraria thunbergiana* Benth. (Leguminosae). Bachelor thesis, Yamaguchi Univ. 100 pp. (1986, Unpublished) (in Japanese)
- 19) Kasahara, Y.: Weeds of Japan Illustrated. 11th ed., 518 pp., Yokendo, Tokyo (1982) (in Japanese)
- 20) Kawabe, A.: Tortricidae. in Inoue, H. et al. Moths of Japan, I: 62–158. Kodansha, Tokyo (1982) (in Japanese)
- 21) Kawai, S.: Scale insects of Japan in colors. 455 pp. Zenkoku Noson Kyoiku Kyokai, Tokyo (1980) (in Japanese)
- 22) Kawasaki, T. and Kawamura, M.: Hundred species of Heteroptera. 311 pp. Zenkoku Noson Kyoiku Kyokai, Tokyo (1977) (in Japanese)
- 23) Kimoto, S.: Chrysomelidae. in Hayashi, M. et al. (eds.), The Coleoptera of Japan in color, IV: 147–222, pls. 29–43. Hoikusha, Osaka (1984) (in Japanese)
- 24) Kitamura, S. and Murata, G.: Coloured illustrations of Herbaceous plants of Japan (Choripetalae). 390 pp., 72 pls. Hoikusha, Osaka (1979) (in Japanese)
- 25) Kobayashi, T.: Kamemushi-ru. In: Genshoku sakumotsu byogaichu hyakka, 2: 229–235. Nousan-gyoson Bunka Kyokai, Tokyo (1987) (in Japanese)
- 26) Kojima, K. and Nakamura, S.: Food plants of Cerambycid beetles (Cerambycidae, Coleoptera) in Japan. 336 pp., Hiba Soc. Nat. Hist., Hiroshima (1986) (in Japanese)
- 27) Kubo, M., Nagao, T., Mizuno, M. and Ohashi, H.: Pharmacognostical studies on the Chinese crude drugs gehua. Syoyakugaku Zasshi 31: 136–144 (1977) (in Japanese with English summary)
- 28) Kudo, I.: A new species of the genus *Salpingothrips* from Japan (Thysanoptera: Thrypidae). *Kontyû* 40, 230–233 (1972)
- 29) Kumata, T.: Taxonomic studies on the Lithocolletinae of Japan (Lepidoptera: Gracillariidae), Part II. *Ins. Mats.* 26, 1–48 (1963)
- 30) Kumata, T., Kuroko, H. and Ermolaev, V. P.: Japanese species of the *Acrocercops*-group (Lepidoptera: Gracillariidae), Part II. *Ins. Mats., N.S.* 40, 1–133 (1988)
- 31) Kuroko, H.: Notes on the *Liocrobyla* of Japan, with descriptions of two new species (Lepidoptera: Gracillariidae). *Esakia* 1, 1–7, 3 pls. (1960)
- 32) Kuroko, H.: Revisional studies on the family Lyonetiidae of Japan (Lepidoptera). *Esakia* 4, 1–61, 17 pls. (1964)
- 33) Kuroko, H.: Lyonetiidae and Gracillariidae. in Inoue, H. et al., Moths of Japan, I: 172–202. Kodansha, Tokyo (1982) (in Japanese)
- 34) Kurosawa, Y.: Buprestidae. in Nakane, T. et al., Iconographia Insectorum Japonicorum Colore naturali edita, II: 147–156, pls. 74–78. Hokuryukan, Tokyo (1963) (in Japanese)
- 35) Kurosawa, Y.: Buprestidae. in Kurosawa, et al. (eds.), Colored illustrations of the Coleoptera of Japan, III: 2–38, pls. 1–7. Hoikusha, Osaka (1985) (in Japanese)
- 36) Makino, T.: Makino's new illustrated flora of Japan. 906 pp. Hokuryukan, Tokyo (1989) (in Japanese)
- 37) Miller, J. H. and Edwards, B.: Kudzu *Pueraria lobata* where did it come from and how can we stop it. *South J. Appl. For.* 7, 165–169 (1983)
- 38) Miyamoto, S.: Miridae. in Asahina, S. et al. (eds.), Iconographia Insectorum Japonicorum Colore naturali edita, III: 97–100, pls. 49–50. Hokuryukan, Tokyo (1965) (in Japanese)
- 39) Miyata, A.: Handbook of the moth ecology, I, II: 1451 pp., Shouwado, Nagasaki (1983) (in Japanese with English summary)
- 40) Miyatake, Y.: Homoptera. in Ito, S. et al. (eds.), Colored illustrations of the insects of Japan, II: 34–40,



- pls. 130–174. Hoikusha, Osaka (1977) (in Japanese)
- 41) Miyazaki, M. and Kudo, I.: Bibliography and plant catalogue of Thysanoptera of Japan. Miscellaneous publication of the National Institute of Agro-environmental Sciences No. 3, 246 pp. (1988) (in Japanese)
  - 42) Morimoto, K.: Ojiroashinagazou no chuei. *Gensei*, 3, 26 (1954) (in Japanese)
  - 43) Morimoto, K.: Apionidae and Curculionidae. in Hayashi, M. et al. (eds.), The Coleoptera of Japan in color, IV: 265–345, pls. 53–68. Hoikusha, Osaka (1984) (in Japanese)
  - 44) Moritsu, M.: Aphids of Japan in colors. 545 pp. Zenkoku Noson Kyoiku Kyokai, Tokyo (1983) (in Japanese)
  - 45) Ohbayashi, N.: Dermestidae. in Kurosawa, Y. et al. (eds.), Colored illustrations of the Coleoptera of Japan, III: 125–130, pl. 21. Hoikusha, Osaka (1985) (in Japanese)
  - 46) Oho, N. (ed.): Major insect and other pests of economic plants in Japan. 307 pp. Jap. Pl. Prot. Assoc., Tokyo (1980) (in Japanese)
  - 47) Oku, T., Miyahara, T., Fujimura, T. and Toki, A.: Preliminary note on *Matsumuraeses* species (Lepidoptera: Tortricidae) injuring soybeans in Tohoku district. *Jap. J. appl. Ent. Zool.*, 27, 28–34 (1983) (in Japanese with English summary)
  - 48) Okuno, T., Tanaka, T., Kimura, Y. and Yoneyama, S.: Diseases and pests of flowers and vegetables in colour, vii + 366 pp., 64 pls. Hoikusha, Osaka (1978) (in Japanese)
  - 49) Patterson, D. T.: The history and distribution of 5 exotic weeds in North-Carolina, U.S.A. *Castanea* 41, 177–180 (1976)
  - 50) Rawat, R. R. and Sahu, H. R.: New record of *Chauliops fallax* Scott (Hemiptera: Lygaeidae) on moong and urid (*Phaseolus* spp.). *Ind. J. Ent.* 30, 179 (1968)
  - 51) Razowski, J. and Yasuda, T.: On the Laspeyresiiini genus *Matsumuraeses* Issiki (Lepidoptera: Tortricidae). *Acta zool. Cracov.* 22, 89–106 (1975)
  - 52) Sasakawa, M.: A study of the Japanese Agromyzidae (Diptera), Part 2. *Pac. Ins.* 3, 307–473 (1961)
  - 53) Sasakawa, M.: Descriptions of three new leaf-mining pests (Diptera: Agromyzidae). *Appl. Ent. Zool.* 16, 149–155 (1981)
  - 54) Sasek, T. W. and Strain, B. R.: Implications of atmospheric carbon dioxide enrichment and climatic change for the geographical distribution of two introduced vines in the U.S.A. *Clim. Change.* 16, 31–52 (1990)
  - 55) Shirozu, T. and Hara, A.: Early stages of Japanese butterflies in colour, II. 139 pp., 120 pls., Hoikusha, Osaka (1962) (in Japanese)
  - 56) Song, X. P., Chen, P. P. and Chai, X. S.: Effects of puerarin on blood pressure and plasma renin activity in spontaneously hypertensive rats. *Acta Pharmacol. Sin.* 9, 55–58 (1988) (in Chinese with English summary)
  - 57) Sugi, S.: Noctuidae. in Inoue, H. et al., Moths of Japan, I: 669–913. Kodansha, Tokyo (1982) (in Japanese)
  - 58) Sweet, M. H. and Schaefer, C. W.: Systematic status and biology of *Chauliops fallax* Scott, with a discussion of the phylogenetic relationships of the Chauliopinae (Hemiptera: Malcidae). *Ann. Ent. Soc. Amer.* 78, 526–536 (1985)
  - 59) Takafuji, A. and Ishii, T.: Inheritance of sex ratio in the Kanzawa spider mite *Tetranychus kanzawai* Kishida. *Res. Pop. Ecol.* 31, 123–128 (1989)
  - 60) Takeya, K. and Itokawa, H.: Iso flavonoids and the other constituents in callus tissues of *Pueraria lobata*. *Chem. Pharm. Bull.* 30, 1496–1499 (1982)
  - 61) Tanner, R. D. and Hussain, S. S.: Kudzu *Pueraria lobata* root starch as a substrate for the lysine enriched bakers yeast and ethanol fermentation process. *J. Agric. Food Chem.* 27, 22–27 (1979)
  - 62) Tayutivutikul, J. and Yano, K.: Biology of insects associated with the Kudzu plant, *Pueraria lobata* (Leguminosae) 1. *Chauliops fallax* (Hemiptera, Lygaeidae). *Jpn. J. Ent.* 57, 831–842 (1989)
  - 63) Tayutivutikul, J. and Yano, K.: Biology of insects associated with the Kudzu plant, *Pueraria lobata* (Leguminosae) 2. *Magacopta punctissimum* (Hemiptera, Plataspidae). *Jpn. J. Ent.* 58, 533–539 (1990)
  - 64) Umeya, K., Kudo, I. and Miyazaki, M.: Pest thrips in Japan. Zenkoku Noson-Kyoiku Kyokai, Tokyo, 422 pp. (1988) (in Japanese)
  - 65) van Emden, H. F.: The role of uncultivated land in the biology of crop pests and beneficial insects. *Sci. Hort.* 17, 121–136 (1965)

- 66) van Emden, H. F.: Wild plants in the ecology of insect pests. in Thresh, J. M. (ed.), Pests, pathogens and vegetation. 251–261. Pitman Advances Pub. Program, Boston, London, Melbourne (1981)
- 67) Yamamoto, Y.: Geometridae. in Issiki, S. et al., Early stages of Japanese moths in colour, II: 29–50, 10 pls. (1969) (in Japanese)
- 68) Yamamoto, M.: Noctuidae. in Sugi, S. (ed.), Larvae of larger moths in Japan. Kodansha, Tokyo. 453 pp. (1987) (in Japanese with English summary)
- 69) Yamauchi, S., Ikenaga, H. and Yukawa, J.: Midge galls collected from the south-west islands of Japan. *Satsuma* 31, 1–32 (1982) (in Japanese)
- 70) Yukawa, J.: A revision of the Japanese gall midges (Diptera: Cecidomyiidae). *Mem. Fac. Agr. Kagoshima Univ.* 8, 1–203 (1971)
- 71) Yukawa, J.: Midge galls of Kagoshima Prefecture (Diptera: Cecidomyiidae). *Satsuma* 37, 175–205 (1988) (in Japanese)
- 72) Yukawa, J. and Sunose, T.: Midge galls of Niigata Prefecture (Diptera: Cecidomyiidae). *Trans. Essa ent. Soc., Niigata* 66, 45–58 (1988) (in Japanese with English summary)